PASJ: Publ. Astron. Soc. Japan , 1–**??**,
© 2008. Astronomical Society of Japan.

arXiv:astro-ph/0310209v1 8 Oct 2003

Variable Star Network: World Center for Transient Object Astronomy and Variable Stars

Taichi Kato, Makoto Uemura, Ryoko Ishioka
Department of Astronomy, Kyoto University, Sakyo-ku, Kyoto 606-8502
tkato@kusastro.kyoto-u.ac.jp, ishioka@kusastro.kyoto-u.ac.jp, uemura@kusastro.kyoto-u.ac.jp
Daisaku Nogami
Hida Observatory, Kyoto University, Kamitakara, Gifu 506-1314
nogami@kwasan.kyoto-u.ac.jp
Chatief Kunjaya
Institute of Technology, Bandung, Indonesia
kunjaya@as.itb.ac.id
Hajime Baba
Center for Planning and Information Systems, The Institute of Space and Astronautical Science (ISAS),
Sagamihara, Kanagawa 229-8510
baba@plain.isas.ac.jp
and
Hitoshi Yamaoka
Faculty of Science, Kyushu University, Fukuoka 810-8560
yamaoka@rc.kyushu-u.ac.jp

(Received ; accepted)

Abstract

Variable Star Network (VSNET) is a global professional-amateur network of researchers in variable stars and related objects, particularly in transient objects, such as cataclysmic variables, black hole binaries, supernovae and gamma-ray bursts. The VSNET has been playing a pioneering role in establishing the field of *transient object astronomy*, by effectively incorporating modern advance in observational astronomy and global electronic network, as well as collaborative progress in theoretical astronomy and astronomical computing. The VSNET is now one of the best-featured global networks in this field of astronomy. We review on the historical progress, design concept, associated technology, and a wealth of scientific achievements powered by the VSNET.

Key words: accretion, accretion disks — astronomical data bases: miscellaneous — catalogs — galaxies: BL Lacertae objects: general — gamma rays: bursts — methods: data analysis — stars: binaries: eclipsing — stars: binaries: symbiotic — stars: dwarf novae — stars: emission-line, Be — stars: novae, cataclysmic variables — stars: oscillations — stars: pre-main sequence — stars: supernovae — stars: variables: other — stars: Wolf-Rayet — surveys — X-rays: bursts

1. Introduction

1.1. *Transient Object Astronomy*

With the progress of modern physics and astronomy, our outlook on the universe is dramatically changing from the stationary universe to the dynamic, ever changing, universe. The dynamical phenomena in the universe appear as variations at various time-scales ranging from the cosmological evolutionary time scale to less than a milisecond. These time-variations are becoming actually observable with the advent of the modern observing equipment and technology. Among them, time-variations arising from extreme gravity as best exemplified by black holes, and from degenerate objects, such as white dwarfs and neutron stars, have been receiving extreme attention from various fields of modern science, as the natural laboratory of general relativity and quantum mechanics, which best represent the glorious success of the "century of physics".

As can be easily expected from the extreme conditions, astronomical phenomena under strong gravity or in degenerate conditions have extremely short time-scales, and are known to be usually very unpredictable. These astronomical phenomena are now generally called "transient phenomena", or referred to as "transient objects". The concept of *transient object astronomy* appeared very late in the history of astronomy and now flourishing as a new modality of astrophysical research.[1] This success greatly owed to the recent great advancement of observing modalities, information technology and computational astrophysics. The Variable Star Network (VSNET), the objective of this review, is one of the earliest and most successful international groups that led to the modern success of transient object astronomy.

---

[1] As shown in Appendix 1, the VSNET is the earliest group which began using the term *transient objects* in the present context of astronomical significance.

1.2. New Window to Transient Object Astronomy

In the research history of *transient object astronomy*, there were two major breakthroughs in the early 1990's. The one is the development of easy availability of CCDs and personal computers, and the other is the advent of the internet. These two breakthroughs played a key role in establishing *transient object astronomy* as one of the most popular contemporary astronomy topics.

From the traditional viewpoint, CCDs were usually used as a "faint-end" extension of the former photon detection methods, e.g. photoelectric and photographic observations. This naturally led to a pursuit of observing fainter stars on long-exposure CCD images (cf. Howell, Szkody 1988; Szkody et al. 1989; Howell et al. 1990; Howell et al. 1991).

The founder of the VSNET was one of the first to break this tradition, and was virtually the first person who systematically turned modern CCD equipment to bright, transient objects, such as classical novae and outbursting dwarf novae (the best examples being Kato, Hirata 1991; Kato 1991b; Kato, Takata 1991, see the later sections for their scientific achievements). The traditional time-resolved observations of classical novae and outbursting dwarf novae were almost restricted to so-called target-of-opportunity (TOO) observations. The best traditional examples include the 1978 outburst of WZ Sge (Patterson, McGraw 1978; Mattei 1980; Patterson et al. 1981), and the 1986 outburst of SW UMa (Robinson et al. 1987). This kind of observations was usually severely limited by the telescope time allocation, and many important transient phenomena (e.g. the 1985 historical long outburst of U Gem: Cannizzo et al. 2002) faded away without receiving sufficient observational coverage. Traditional proposals for telescope time were also limited because of the transient and unpredictable nature of these phenomena; there is no guarantee that there is a suitable transient target at the time of allocated observation. For this reason, systematic observational research in these objects was severely restricted to short-period, less unpredictable objects, with an enormous effort of world-wide coordination (e.g. VW Hyi: Schwarzenberg-Czerny et al. 1985; Pringle et al. 1987; YZ Cnc: van Paradijs et al. 1994).

Timely circulation of alerts on transient objects or phenomena is also crucially important, particularly for TOO-type observation. Before the wide availablity of the internet, the typical way of communicating such alerts was a phone call from an observer (usually an amateur astronomer watching variable stars) to a variable star organization, which was typically relayed (with some delay) to local observers for confirmation. The information, if it was recognized as particularly important, was then distributed to world-wide observers usually from the Central Bureau of Astronomical Telegrams (CBAT) via telegrams, direct phone calls, or slow postcards. It usually took, even in best cases, a day or more before this crucial information was relayed to the actual observer undertaking a TOO observation. The early stage of transient objects was usually missed because of this delay. For example, the detection of the 1986 historical outburst of SW UMa was relayed via an astronomical telegram only when the object reached a historical brightness of $V \sim 9$, although the outburst was initially reported ~ 1.5 below the peak brightness. There had been very few early stage observations (i.e. within a day of the event detection) of transient objects before the 1990's.

1.3. Early Public Electronic Communication

This situation drastically changed with the public availability of the internet. In the early times (around 1990–1991), there were only sporadic internet communications on observations, mainly via personal e-mails and on public bulletin board systems. This strategy worked slightly better than in the past, the situation was basically unchanged in that most of observers had to rely on occasional communications or a slow access to news materials.

From the necessity of publicly and electronically disseminating urgent astronomical phenomena, there appeared e-mail exploders (mailing lists). The Scandinavian *varstars* list and the (mainly) professional *novanet* by the Arizona State University team played an early important role in publicly relaying information on transient objects.[2] The early-time progress of these electronic communications is summarized in the *vsnet-history* list.[3]

The scientific role of wide-availability of these e-mail exploders was recognized upon the appearance of SN 1993J in M 81 (Ripero et al. 1993; Pease 1993). This supernova showed an unusual early-time light curve and a spectral transition from a type-II to type-Ib supernova (Nomoto et al. 1993; Swartz et al. 1993; Filippenko et al. 1993a; Podsiadlowski et al. 1993). In communicating nightly rapid changes and distributing most up-to-date observation strategies, the e-mail exploders played a more crucial role than ever. Another advantage of e-mail exploders as a *standardization tool* of observations became evident (figure 1). Early-time non-standard observations were quickly corrected using the updated photometric comparison stars, and questionable observations were examined real-time to clarify the cause. This led to a huge world compilation of SN 1993J photometry updates (see figure 2) contributed by a number of volunteers, including the VSNET founder.[4] This high-quality, uniform compilation of real-time observations greatly contributed to real-time theoretical modeling of this object (e.g. Nomoto et al. 1993), spectroscopy (e.g. Baron et al. 1993; Taniguchi et al. 1993; Clocchiatti et al. 1995; Trammell et al. 1993a) and photometry (e.g. van Driel et al. 1993; Wheeler et al. 1993). We published our own results in Okyudo et al. (1993). We also contributed to a number of International Astronomical Union Circulars (IAUCs) (Kato, Hanzl 1993; Zimmermann et al. 1993; King et al. 1993; Filippenko et al. 1993b; Trammell

---

[2] Both networks do not exist at present.

[3] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-history/maillist.html⟩.

[4] The final version of the "photometry update" is publicly available at ⟨ftp://vsnet.kusastro.kyoto-u.ac.jp/pub/vsnet/SNe/sn1993j/sn.mag⟩



Fig. 1. Circulated electronic chart of SN 1993J (issued on 1993 April 13) showing V-band comparison stars for standardization.

et al. 1993b; Hu et al. 1993). The complete history of this SN 1993J story can be also seen in the *vsnet-history* archive.

1.4. *Opening of the Electronic Era of Transient Object Astronomy*

Upon the recognition of the importance of e-mail exploders on the occasion of SN 1993J, more systematic efforts were taken to standardize the communication and data reporting method. In relation to reporting observations, we started widely disseminating observations of regular variable star observations, mainly submitted to the Variable Star Observers League in Japan (VSOLJ),[5] and those personally reported to us. People started recognizing the scientific importance of widely disseminating regular observations, which can be readily reflected on scheduling new observations. New findings based on widely reported observations (e.g. superhump detection of a dwarf nova) were also relayed real-time, which worked as a positive feedback to original observers. The prototype of VSNET-type e-mail exploders was thus established in 1993. The next major astronomical event at this stage of the history was the discovery of Nova Cas 1993 (V705 Cas). This nova showed considerable degree of early-time fluctuations, as well as a later dust-forming episode. During all of the stages of evolution of the nova explosion, the data circulating strategy established at the time of SN 1993J played an impressive role: the comprehensive compilation of V705 Cas by Yasuto Takenaka (see figure 3) was cited in a Nature paper (Shore et al. 1994) as best authenticated optical record of this nova. The nova was later even symbolically called *an electronic nova* (Pepin 1995), representing the opening of new electronic era of transient object astronomy.

1.5. *Establishment of VSNET*

The information of these transient objects and regular variable star observations was initially relayed manually, or relayed on existing less specified e-mail exploder systems. In 1994, our own e-mail exploder system (VSNET) started working. This service smoothly took over the past manual e-mail announcement systems, and immediately received wide attention both from amateur and professional communities. The establishment of the VSNET thus became the “prototype” of world-wide amateur-professional collaborations based on public e-mail communication. This initiative later led to a flourishing VSNET Collaboration (section 4). The early history was reviewed by D. Nogami et al. (1997) in “Electronic Publishing, Now and the Future”, Joint Discussion 12 of the 23rd IAU General Assembly. Considering the historical significance in the advent of *transient object astronomy* and the current unavailability of this document in a solid publication, we reproduce the presented contents in Appendix 1 (in order to preserve the original contents, we only corrected minor typographical errors). The VSNET mailing list system now has more than 1300 subscribers from more

---

[5] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/VSOLJ/vsolj.html⟩ and see ⟨http://vsolj.cetus-net.org/⟩ for the VSOLJ Variable Star Bulletin page.



Fig. 2. Light curve of SN 1993J, drawn from the "SN 1993J photometry update" (see text). The Large and small dots represent (nearly) V-band and visual observations.



Fig. 3. Light curve of V705 Cas (Nova Cas 1993), covering the rise and the "great fade", and rebrightening phases, drawn from the observations circulated through the early development of the VSNET. The dots and open circles represent visual and photographic/CCD observations, respectively.

than 50 countries all over the world.

2. VSNET as a Modern-Day "Electronic Variable Star Organization"

2.1. *Historical Development of the Electronic Reporting System*

During the very initial stage of the development of the VSNET, we simply relayed observations to those who (potentially) need the data. However, it soon became evident, from the experiences with SN 1993J and V705 Cas (subsection 1.3), that there is a need for a newly designed reporting system adapted for electronic data exchanges. Since we already had sufficient experience with relaying VSOLJ reports to the world-wide variable star observers, it was a natural solution to extend the VSOLJ format to an international version. This changes were minimal, by introducing Universal Time (UT)-based system and the extension of coding system of observers. The details of the reporting system is described in Appendix 2.

By globally collecting data, we soon recognized the necessity for setting up a dedicated e-mail list for reporting observations, *vsnet-obs*.[6] The alert list *vsnet-alert*[7] was prepared at the same time, which has been one of the most renowned and reliable sources of noteworthy phenomena of variable stars and transient objects, and the messages in *vsnet-alert* have been frequently cited in the professional literature as the primary source of information (e.g. Downes et al. 1997; Downes et al. 2001; Ritter, Kolb 1998; Ritter, Kolb 2003; Kube et al. 2003).

---

[6] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-obs/maillist.html⟩. Because of the large number of articles, the online archive is subdivided; see ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/index.html⟩. for the complete message archive.

[7] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-alert/maillist.html⟩.

2.2. Standardization

The VSNET played an important role in standardization of variable star observing. There are several steps in standardization: (1) standardization of reporting format (see subsection 2.1), (2) standardization of comparison stars, and other minor steps. In standardizing comparison stars, the VSNET group also took an initiative in the history of transient object astronomy. The earliest example includes SN 1993J and classical novae, for which reliable CCD-based (or sometimes photoelectric) comparison star sequences were determined and distributed through the VSNET lists. Before this standardizating efforts were taken, the nova researchers had to cope with often unreliable early reports in IAUCs, which were often based on various sources of comparison stars with notoriously diverse photometric quality. With the advent of the VSNET, the modern-day nova observations have now become as reliable as those of other variable stars with well-established comparison star sequences. The VSNET group has been paying attention to the quality of the original discovery reports, and have issued several magnitude updates superseding the IAUC announcements. The same effort has been taken for supernova photometry, although the faintness and the large number of the target objects have made it a more difficult task than in classical novae.

In observation of (non-transient) ordinary variable stars, the VSNET took the initiative to standardize the comparison star magnitudes to the modern V system, from various old systems including the traditional Harvard visual photometric system. The effort was initially taken to revise the faint-end magnitudes for cataclysmic variables (CVs) and peculiar variables using the CCD camera at Ouda Station, Kyoto University (Ohtani et al. 1992). These results were continuously released as "VSNET charts" through the VSNET lists (see figure 4). Several independent contributed efforts, by Rafael Barbera (Grup d'Estudis Astronomics)[8], and Fraser Farrell (Astronomy Society of South Australia),[9] as well as those by the VSNET administrator team, were made to write software packages to graphically display this VSNET chart format. Brian Skiff has been continuously contributing to the VSNET in photoelectrically standardizing sequences for selected variable stars; this initiative was globally taken over with the CCD works, notably by Arne Henden and Bruce Sumner.[10] These standardized sequences and charts have own lists *vsnet-sequence* and *vsnet-chart*.[11]

For the bright end, we were the group to first extensively use Hipparcos and Tycho magnitudes at the earliest epoch (1997) of the public release of these catalogs. We immediately made public variable star charts based on Hipparcos and Tycho magnitudes.[12] Since then, this adoption of the standard V-band system (selected for $B-V$ colors) for visual photometry has been gradually becoming the global standard. Since the public release of Tycho-2 catalogue, this standard was extended to a slightly fainter magnitude.[13]

For poorly observed faint objects, we also took an initiative (vsnet-chat 700, in 1998)[14] to calibrate USNO catalog magnitudes since the early release of the USNO A1.0 catalog (Monet et al. 1998a). This calibration has been widely used, when an alternative sequence is not readily available.

2.3. Collaboration with the VSOLJ

From the very beginning of its history, the VSNET has been in continuous collaboration with the VSOLJ. The activity includes hosting the VSOLJ alert and discussion mailing lists, distributing the VSOLJ reports and prediction of variable stars, and hosting the public VSOLJ database and light curves (see subsection 3.2). A part of standardization schemes (subsection 2.2) has been developed in collaboration with the VSOLJ administrator group (see also Appendix 2) and through a discussion with world-wide variable star leaders at the International VSOLJ Meeting held in conjunction with the 23rd IAU General Assembly.

The other pioneering aspect of the VSNET–VSOLJ collaboration was the introduction of CCDs for CV photometry. This work was mainly done in collaboration with Makoto Iida, who took the world initiative to monitor faint CVs with a commercial CCD. This collaboration led to fruitful scientific results (NY Ser: Iida et al. 1995b; DV Dra: Iida et al. 1995a; BC UMa, see ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/DNe/bcuma.html⟩). These successful results were demonstrated in a number of international conferences, including the Padova CV Conference in 1995 and the Keele CV Conference in 1997, and the results became gradually digested by the professional community. This pioneering amateur-professional collaboration in transient objects and CVs finally led to the most successful VSNET Collaboration (section 4), and the strategy was taken over, albeit with a lesser degree of the original "flavor" of amateur-professional relations and publicity policy, by a number of following world-wide groups with the similar strategies.

2.4. Collaboration with the Southern Hemisphere Observers

Since the VSNET service initially started with the coordinating role among mostly northern hemisphere observers, there were initially few reports from the southern observers. With the outstanding activity of the VSNET, there arose a number of requests from professional astronomers who were planning for TOO observations of southern dwarf novae. This situation has been gradu-

---

[8] ⟨http://www.astrogea.org/⟩.

[9] ⟨http://www.assa.org.au/⟩.

[10] ⟨ftp://ftp.nofs.navy.mil/pub/outgoing/aah/sequence/sumner/⟩.

[11] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-sequence/maillist.html⟩ and ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-chart/maillist.html⟩.

[12] ⟨ftp://vsnet.kusastro.kyoto-u.ac.jp/pub/vsnet/charts/HIPTYC/⟩

[13] ⟨ftp://vsnet.kusastro.kyoto-u.ac.jp/pub/vsnet/charts/Tycho-2/⟩

[14] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-chat/msg00700.html⟩



Fig. 4. Example of VSNET chart issued in 1996. The object is GM Sgr (currently renamed as V4641 Sgr; note that this chart provided the correcteed identification against the previous publication). The availablity of a chart with the correctly identified object enabled the discovery of the giant outburst in 1999 (see subsection 6.4.1).

ally and progressively improved by increasing contributions from the southern observers, particularly by the members of Royal Astronomical Society of New Zealand (RASNZ).[15] By now, these contributions, notably by Rod Stubbings, have enabled a number of rare outburst detections and early circulation of these phenomena. Together with the collaboration with southern CCD observers, the scientific achievements of transient object astronomy in the southern hemisphere is explosively growing (e.g. microquasar V4641 Sgr: Stubbings, Pearce 1999; Uemura et al. 2002d; Uemura et al. 2002h, southern SU UMa-type dwarf novae: Kato et al. 2003h; Kato et al. 2003b; Kato et al. 2003d), which will be reviewed in later sections.

Most recently, we started collaborating with the Nagoya University team for identifying X-ray transients (see subsection 6.4) with the Simultaneous-3color InfraRed Imager for Unbiased Survey (SIRIUS) camera installed at the Infrared Survey Facility (IRSF)[16] situated at the South African Astronomical Observatory, Southerland, South Africa. Several scientific achievements have already been issued (Kato et al. 2003a; Baba et al. 2003a; Nagata et al. 2003). A work on V359 Cen in collaboration with the Microlensing Observations in Astrophysics (MOA) project[17] team has been also published (Kato et al. 2002k).

3. World-Wide Web and Anonymous FTP Service

3.1. *Historical Development*

From the very beginning of the VSNET, all messages posted to VSNET public mailing lists are opened to the public. A message can be relayed (unless the sender otherwise specifies the usage) to third-party members, or can be posted to a different mailing list or a public news service. In the earliest times, these messages were archived (they were sometimes made public) at individual receivers.

From the necessity of publicly providing standardized charts and related materials (see subsection 2.2), we initially used private anonymous ftp service operated at the Department of Astronomy, Kyoto University. This was replaced by the official VSNET anonymous ftp service in 1995 July. Contributed programs, notably VSNCHT written by Rafael Barbera, to graphically display VSNET-format charts were made public from the start of the service. Magnitude summaries of selected objects such as new novae, our own and contributed CCD images of outbursting dwarf novae were soon made available through the anonymous ftp service.

We set up the official VSNET World-Wide Web (WWW) service[18] in 1995 June. The web pages have been continuously updated, particularly in announcing newly discovered transient object phenomena. Even during these updates, we have paid all our efforts to preserve the original URLs for future reference; almost all pages that existed in the past can be tracked with the original URLs even now. The other design of the VSNET WWW system was in the combined usage of the WWW and ftp services. In the earliest times, not all internet users were able to use window-based browsers. We have thus set up two ways (WWW and anonymous ftp) of access to the desired data. With this feature, a user is able to get necessary images or programs even without a browser or a fast internet connection. This feature became, however, less important with the wide availablity of the window-based browsers. The ftp service soon included a complete archive of the VSNET mailing list messages.

In 1997 April, we started public WWW service of all archival messages posted to the VSNET lists. The WWW archive is automatically updated by using the MHonArc system.[19] We once implemented the Namazu-based[20] full-text search engine on the VSNET WWW service, but this was discontinued because publicly available search engines now have equivalent functions.

3.2. *Public Light Curve Archive*

With the development of the VSNET WWW service, we started public light curve archive service in 1996 August.[21] These light curves are drawn and regularly updated from *vsnet-obs* reports, which were incorporated into the VSNET database (Appendix 2). At present, the regular updates of these (static) light curves archive are performed by a Java$^{\rm TM}$-based light curve generator engine wrapping the Linux-based VSNET database system.

The VSNET light curve archive also hosts light curves drawn from the VSOLJ and Association Française des Observateurs d'Etoiles Variables (AFOEV)[22] public database, by courtesy of the respective organizations. These light curves can be easily reached from the VSNET top page, as well as from the AFOEV website.

We also implemented individual variable star pages[23] generated from the GCVS electronic edition. These pages provide handy links to the light curves (VSNET, VSOLJ, AFOEV) and links to the relevant pages and charts on the VSNET.

3.3. *Data Search Engine and Light Curve Generator*

Soon after the establishment of the data reporting system, we also implemented a Common Gateway Interface (CGI)-based data search engine in 1996 June.[24] This service returns the observations of a specified variable star selected from the VSNET public database, which is coherently updated from the regular variable star reports to the VSNET. This type of interactive variable star data browser was the first one in the world, and became a prototype of the subsequent similar services. This interactive data search engine has been widely used and frequently referred in the professional literature. A number of WWW services, including the well-known "Latest Supernovae" page[25] by David Bishop, provides links to most up-to-date compilations of variable star reports by directly referring to the VSNET data search engine. This WWW-based data search engine provides the modern-day extension of the early-time magnitude summaries (subsection 1.4).

---

[15] ⟨http://www.rasnz.org.nz/⟩.

[16] ⟨http://www.saao.ac.za/facilities/irsf/irsf.html⟩.

[17] ⟨http://moa.scitec.auckland.ac.nz/⟩.

[18] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/⟩.

[19] ⟨http://www.mhonarc.org/⟩.

[20] The Namazu full-text WWW-based search engine was developed by one of the VSNET administrators, Hajime Baba. This is one of the best used systems among search engines for non-commercial, personal or academic uses. See ⟨http://www.namazu.org/⟩ for the Namazu project.

[21] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/LCs/index.html⟩.

[22] ⟨http://cdsweb.u-strasbg.fr/afoev/⟩.

[23] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/gcvs/index.html⟩.

[24] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/etc/searchobs.html⟩.

In 1998 March, this VSNET data search engine furthermore started providing a Java$^{\mathrm{TM}}$ applet-based interactive light curve interface,[26] with which a user can freely browse the data (both VSNET and VSOLJ observations) with a graphical user interface (GUI). This was one of the earliest Java$^{\mathrm{TM}}$ applications in public astronomical service.

4. VSNET Collaboration

4.1. VSNET as World Center of Time-Series Observations of CVs

As described in section 2.3, the VSNET played a pioneering and essential role in establishing a new modality of internet-based amateur-professional collaboration. This direction was one of the main aims of the VSNET from the very start of its internet presence.

Among various kinds of variable stars, CVs are the "canonical" class of transient objects, as briefly reviewed in subsection 1.2. With the common interest in CVs and related systems, the VSNET amateur-professional collaboration originally mainly focused on CVs, especially on unpredictable outbursts of dwarf novae. The actual collaborative studies were done on the existing VSNET list, most frequently on *vsnet-alert* and *vsnet-obs*. In most cases, real-time reports of visual detections of outbursts in dwarf novae, usually after some verification process involving the VSNET administrator team, triggered the actual observing campaigns. This process was usually performed within several hours and a day of the detection. This prompt reaction to event triggers of transient objects later enabled an efficient reaction to gamma-ray burst (GRB) triggers (section 6.15). The unique feature of the VSNET campaigns on transient objects is that they are a collaborative effort between visual observers and CCD observers. The VSNET is historically the first organization that realized the high productivity involving traditional visual variable star observations, although this importance in transient object astronomy had long been stressed and had been a dream among researchers.

The early-time successful cooperative works include: recurrent nova V3890 Sgr (Kato et al. 1990, phone call), detection of superhumps in AQ Eri (Kato 1991a, phone call and e-mail), detection of superhumps in V1251 Cyg (Kato 1991b; Kato 1995a, phone call and e-mail), WX Cet in 1991 (Kato 1995f independent detection and e-mail), EF Peg (Kato, Takata 1991; Kato 2002c, phone call and e-mail), HV Vir in 1992 (Kato et al. 2001l, e-mail), SW UMa in 1992 (Kato et al. 1992a, phone call). Since then, most information is relayed by e-mail and e-mail exploder systems (most of the works were conducted upon response to real-time outburst detection alerts reported through the VSNET): V344 Lyr (Kato 1993), HY Lup = Nova Lup 1993 (Kato et al. 1993b), AK Cnc (Kato 1994; Mennickent et al. 1996), T Leo (Kunjaya et al. 1994), KV And (Kato et al. 1994), AY Lyr (Nogami et al. 1994), CY UMa (Kato 1995b), KV And in 1994 (Kato 1995c), FO And (Kato 1995d), TT Boo (Kato 1995e), discovery of ER UMa (Kato, Kunjaya 1995), which will be described in subsection 6.1, PU Per (Kato, Nogami 1995), HS Vir (Kato et al. 1995d), GO Com (Kato et al. 1995c), DH Aql (Nogami, Kato 1995b), V1159 Ori (Nogami et al. 1995a), HV Aur (Nogami et al. 1995b), V725 Aql (Nogami et al. 1995d, own detection), RZ LMi (Nogami et al. 1995c), V1028 Cyg in 1995 (Baba et al. 2000). This continuous stream of scientific reports based on the VSNET amateur-professional cooperations brought a great impact on the community (Liller 1996).

During this early-epoch collaborative works through the VSNET, several noteworthy rare (typically once in a decade to several decades) phenomena occurred and were studied in unprecedented detail through the VSNET: AL Com in 1995 (Mattei et al. 1995; Nogami, Kato 1995a: the outburst was detected by an AAVSO member, soon relayed to the VSNET, which enabled the detection of "early superhumps", which are observed only for several nights after the start of the outburst. See Kato et al. 1996b; Nogami et al. 1997a for our summaries of this event).[27] This information on the VSNET further enabled extensive follow-up studies by different groups (Patterson et al. 1995a; Patterson et al. 1996; Pych, Olech 1995a; Pych, Olech 1995b; Howell et al. 1996; Szkody et al. 1996).

The 1996–1997 outburst of EG Cnc (Schmeer et al. 1996) was a great surprise. This outburst was detected by Patrick Schmeer, a VSNET member, and immediately relayed through the VSNET alert system. Together with the early detection of early superhumps (Matsumoto, Schmeer 1996; Matsumoto et al. 1998), the object showed an unexpected sequence of post-superoutburst rebrightenings (Kato et al. 1997). This unexpected phenomenon was discovered by ourselves and the collaborative efforts through the VSNET.[28] This detection of the spectacular rebrightening phenomenon, which most clearly illustrated the power of real-time information exchange, also led to a number of observational and theoretical papers (Osaki et al. 1997; Osaki et al. 2001; Patterson et al. 1998; Szkody et al. 2000; Szkody et al. 2002a). This impressive phenomenon exhibited to the public, through the VSNET, the "real-time science" in the making.

---

[25] ⟨http://www.rochesterastronomy.org/snimages/⟩.

[26] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/etc/drawobs.html⟩.

[27] The full details of the phenomenon, made public in the Keele CV Conference in 1996 (Kato et al. 1996c), can be seen at ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/DNe/alcom1.html⟩. The star was nominated as the "star of the year" in this conference.

[28] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/DNe/egcnc.html⟩. It was our greatest pleasure that a number of speakers in the Wyoming CV Conference in 1997 presented this VSNET webpage as representing the most unusual activity of a dwarf nova.

The 1999 outburst of the recurrent nova U Sco (Schmeer et al. 1999; figure 5) again illustrated the ability of the international alert network system provided by the VSNET.[29] The outburst detection was made by Patrick Schmeer, and the alert was immediately disseminated through the VSNET. With this quick notification, a very stringent upper limit became immediately available, which was obtained only less than four hours before the outburst detection. The early news enabled an American observer to catch the real optical maximum ($m_{\rm v} = 7.6$), which was more than one magnitude brighter than had been supposed for this recurrent nova. It is very likely the delays in delivering information were partly responsible for the underestimate of the maximum magnitudes in past outbursts. Almost all important findings near the maximum light (within one day of detection) were obtained before the relevant IAUC was issued. The VSNET collaborative study on this outburst additionally led to the first-ever detection of eclipses during the outburst (Matsumoto et al. 2003b; Hachisu et al. 2000a; Hachisu et al. 2000b). This outburst again produced a rich scientific outcome from various researchers (Kahabka et al. 1999; Lépine et al. 1999; Munari et al. 1999; Kiyota 1999; Ikeda et al. 2000; Anupama, Dewangan 2000; Evans et al. 2001; Iijima 2002): when compared to the results from previous outbursts (1979: Barlow et al. 1981; Warner 1979, 1987: Rosino, Iijima 1988; Sekiguchi et al. 1988), the 1999 result could be even referred to as the "victory in the electronic era" of transient object astronomy.

After experiences with these, and other spectacular transient phenomena (which will be reviewed in section 6), we set up new lists (*vsnet-campaign*[30] and other sublists), which primarily deal with campaigns on selected targets. With the establishment of these dedicated lists, we started referring to our amateur-professional world-wide collaboration group as VSNET Collaboration. This (not too tightly bound) group has become harboring subsequent intensive studies undertaken in the VSNET. Figure 6 shows the global distribution of contributors.

4.2. VSNET Campaign Lists and Summaries

The VSNET campaign list and sublists were thus progressively established since 2000, encapsulating a wide range of transient astronomical phenomena. The VSNET campaign lists are subdivided into categories based on object classes (e.g. *vsnet-campaign-dn* for dwarf novae, *vsnet-campaign-xray* for X-ray binaries). Most recently, following the discovery of very unusual objects or phenomena of public interest, we sometimes set up a separate list focused on single objects (e.g. *vsnet-campaign-v4641sgr*[31] for the microquasar V4641 Sgr (subsection 6.4), *vsnet-campaign-v838mon*[32] for the most unusual stellar explosion (V838 Mon) with an astounding light echo, *vsnet-campaign-sn2002ap*[33] for the nearest hypernova SN 2002ap (see subsection 6.5). These new features enabled interested theoreticians to share real-time information of these most unusual objects. The number of VSNET campaign individual lists is 36 (2003 August).

Summaries of the activities of the VSNET Collaboration have been compiled by Makoto Uemura and issued on a weekly basis (VSNET campaign news).[34] This information has been summarized as a yearly review of the activity of the VSNET Collaboration.[35] These VSNET campaign news and summaries comprise an important (authorized) part of the nearly real-time contribution from the VSNET to the other organizations (section 4.3), and are sometimes cited in themselves as a convenient record of the activities of particular objects.

Nowadays the VSNET campaigns are more or less continuously undertaken. In order to inform the current targets of interest, we have recently set up a notification list *vsnet-campaign-target* to campaign contributors.

4.3. Contribution to Other World-Wide Alert Systems

From the very start of the AstroAlert system[36] maintained by the Sky Publishing Co., the VSNET has been nominated as one of the authorized information providers. We now regularly contribute "News from VSNET" to this alert system, primarily notifying the transient astronomical phenomena of current interest. We also occasionally issue alerts on particularly urgent phenomena and nova discoveries. The news have been relayed to the AAVSO, the *astro-l* mailing list, and are widely distributed as the primary source of information on transient astronomical phenomena.

5. Variable Star Related Activities

In addition to main scientific achievements (section 6), the VSNET group has been engaged in various activities in the field of variable stars and variable star-related matters. Here we show our representative activities.

5.1. Variable Star Identifications

The VSNET group has been historically engaged in providing updated information to the General Catalogues of Variable Stars (GCVS: Kholopov et al. 1985a; Samus et al. 2000; Samus' et al. 2002; Samus' et al. 2003; Kazarovets et al. 1998) team,[37] and provided a large amount of variable star identifications and suggestions, mainly based on examination of the historical literature with the modern technology. The present most relevant

---

[29] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Novae/usco.html⟩.

[30] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-campaign/maillist.html⟩.

[31] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-campaign-v4641sgr/maillist.html⟩.

[32] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-campaign-v838mon/maillist.html⟩.

[33] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-campaign-sn2002ap/maillist.html⟩.

[34] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-campaign-news/maillist.html⟩.

[35] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Summary/⟩.

[36] ⟨http://skyandtelescope.com/observing/proamcollab/astroalert/⟩.

[37] ⟨http://www.sai.msu.su/groups/cluster/gcvs/gcvs/⟩.



Fig. 5. Light curve of the 1999 outburst of the recurrent nova U Sco from VSNET observations.



Fig. 6. Global distribution of observers of VSNET Collaboration. The large and small "burst" marks represent regular and occasional participants to CCD photometric campaigns. The number of filled squares relative numbers of represent (primarily) visual or photographic observations from various regions of the earth.

lists are *vsnet-id* and *vsnet-gcvs*[38], which deal with variable star identifications and the GCVS revision project, respectively. The identifications reported to *vsnet-id* are also relayed to the SIMBAD[39] office. There is a series of solid papers on systematic variable star identifications (Kato 1999b; Kato 1999c; Kato 1999d; Kato 1999g; Kato 1999h; Kato 1999i; Kato 1999j; Kato 1999k; Kato 2001j). There have also been collaborations with the Multitudinous Image-based Sky-survey and Accumulative Observations (MISAO) project,[40] which used their own CCD images to identify variable objects near the cataloged positions (Yoshida et al. 1999a; Yoshida et al. 2000b; Yoshida et al. 2000c; Yoshida et al. 2000a; Yoshida et al. 1999b). These collaborative works were conducted also in collaboration with the VSOLJ.

In identifying variable stars, our own chart-plotting system (Appendix 4) has played an important role.

5.2. *New Variable Stars*

The VSNET has been providing the role of a world center of reporting newly discovered variable stars. There is a dedicated mailing list *vsnet-newvar*.[41] The reported new variable stars are usually checked by the VSNET administrator group for identification with other astronomical sources (e.g. X-ray sources, infrared sources, and emission-line stars). These identification processes are also conducted on the public mailing list. These reports on new variable stars on *vsnet-newvar* are now regarded as the primary source of original information by the GCVS team (e.g. Kazarovets et al. 2000; Kazarovets et al. 2001; Kazarovets et al. 2003).

---

[38] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-id/maillist.html⟩ and ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-gcvs/maillist.html⟩.

[39] ⟨http://simbad.u-strasbg.fr/Simbad⟩

[40] ⟨http://www.aerith.net/misao/⟩.

[41] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-newvar/maillist.html⟩.

A preliminary list (newvar.cat) of newly reported, or objects with some peculiarity (which can be candidate variable stars) has been updated, and made public[42] as a reference for those who are looking for new variable stars and variable star identifications. Examination of the properties of poorly known variable stars are also our regular works (many results have been reported through *vsnet-gcvs* lists, as well as through some solid publications such as Kato 2003d; Kato 2003e; Kato 2003f, which were based on the ASAS-3 public data; see subsection 5.3).

5.3. Confirmation of Novae

Confirmation of reported nova candidates is another regular work conducted by the VSNET administrator team, as well as scientific work in novae (subsection 6.2). This work takes advantage of the established VSNET software and catalog systems in variable star identifications (subsection 5.1) and in confirming new variable stars (subsection 5.2). The VSNET system in screening and confirming newly reported nova candidates is one of the most reliable and efficient among all the existing nova confirmation systems. In particular, the VSNET team has even succeeded in recognizing novae from otherwise dismissed new variable star reports (e.g. V1548 Aql: Uemura et al. 2001e; Kato, Takamizawa 2001). The recent discovery of V463 Sct (Haseda 2000; Uemura et al. 2000g; Kato et al. 2002l) is another example of the reliability of the VSNET nova identification system.[43] The reverse case in that the VSNET played an important role in disqualifying a suspected nova. The most recent example was with V4006 Sgr, which was originally reported as a probable nova in IAUC (Nakano et al. 2003a). This object was soon identified with a known variable star through the public discussion in the VSNET.[44] The initial discovery announcement was subsequently readily corrected within a day of the original announcement (Green 2003).

Most recently, we refer to the public photometric real-time database provided by the ASAS-3 team (Pojmanski 2002)[45] for confirming southern nova suspects. The most recent successful example includes V2573 Oph = Nova Oph 2003 (Takao et al. 2003). This application of ASAS-3 public data is also true for confirming new variable stars (e.g. V2552 Oph = Had V98: Kato, Haseda 2003) and better characterizing already known variable stars (e.g. Kato, Stubbings 2003). The individual variable star pages on VSNET website provides convenient links to the ASAS-3 pages (see subsection 3.2).

Upon recognition of new novae, the VSNET team makes an announcement to nova researchers and the variable star community to enable early-time confirmation and follow-up observations. The VSNET is now recognized as one of the most powerful media for disseminating such a kind of alerts to the open community, and is relied on by many professional nova researchers.

For this purpose, VSNET takes an "open policy" of any nova (and supernova, variable star etc.) discovery announcements, i.e. such announcements will be immediately released and made public. The full reasoning of the policy and the actual recommended reporting procedure is described in ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/etc/discovery.html⟩ and the links from this page. The VSNET has a dedicated list for announcing on new novae *vsnet-discovery-nova*,[46] although this kind of information is usually also posted to *vsnet-alert* from the convention. In any case, readers are strongly recommended to refer to the above page before attempting to make an actual nova discovery report.

5.4. Confirmation of Supernovae

As well as confirmation of nova candidates, confirmation of suspected supernovae is also undertaken from the very beginning of the VSNET. The first example was SN 1995D in NGC 2962 (Nakano et al. 1995), which was discovered by Reiki Kushida. The object was confirmed on the same night in Japan by three individual observers (vsnet-alert 30, 31, 32, 1995 February)[47], and the discovery announcement led the multicolor photometry starting soon after the discovery (Sadakane et al. 1996).

Spectroscopic confirmation of several supernovae has also been reported on the VSNET lists. The earliest one was for SN 1995al (vsnet-alert 266, 1995 November),[48] which was distributed well earlier than the formal publication in IAUC (Ayani, Kawakita 1995). Such a quick distribution of spectral type of supernova made it possible to schedule larger telescopes or space-borne instruments to observe it. The nearest type-Ic hypernova SN 2002ap (vsnet-alert 7120, 2002 January)[49] was the most successful recent example (Kinugasa et al. 2002a, Kinugasa et al. 2002b), which was followed by many instruments including the Subaru telescope (Kawabata et al. 2002) or the XMM-Newton (Soria, Kong 2002). Plentiful information, including prediscovery observations (Yamaoka, Rogers 2002), has been posted to the specially made sub-list *vsnet-campaign-sn2002ap*.

These contributions to early-time observation are summarized in subsection 6.5, as well as the scientific results drawn from them.

---

[42] ⟨ftp://vsnet.kusastro.kyoto-u.ac.jp/pub/vsnet/others/newvar.cat⟩.

[43] See ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Novae/hadv46.html⟩ for the full story.

[44] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Novae/v4006sgr.html⟩

[45] ⟨http://www.astrouw.edu.pl/~gp/asas/asas_asas3.html⟩

[46] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-discovery-nova/maillist.html⟩.

[47] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-alert/msg00030.html⟩ and so on.

[48] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-alert/msg00266.html⟩.

[49] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/alert7000/msg00120.html⟩.

5.5. *Hosting Announcements of External Organizations*

The VSNET has been playing a role providing an electronic medium for announcements made by other variable star-related organizations. The examples include the early announcements of AAVSO Newsletters (vsnet-alert 14, 1995 January), Alert Notices (vsnet-alert 58, 1995 March), News Flashes (vsnet-alert 344, 1996 February), website announcement (vsnet-alert 178, 1995 August), ALEXIS[50] transients (vsnet-alert 321, 1996 February), International TOAD Watch website[51] (vsnet-alert 439, 1996 June), website for Hungarian Astronomical Association – Variable Star Section[52] (HAA/VSS, vsnet 756, 1996 September), website for Center for Backyard Astrophysics[53] (CBA, vsnet 872, 1996 November), website for British Astronomical Association, Variable Star Section[54] (BAAVSS, vsnet 893, 1996 November), website for Group of Amateur Astronomers, Czechia, (GAMA, vsnet-alert 596, 1996 November), IAU Commission 42, Bibliography on Close Binaries[55] (BCB, vsnet 981, 1997 January), and numerous announcements on call for observations, electronic publications, and international conferences.[56] The VSNET website and mailing lists thus have been *a world center of variable star-related announcements*.

6. Scientific Achievements of VSNET

6.1. *Dwarf Novae*

Dwarf novae (cf. Osaki 1996a) have historically been the best-studied class of objects by the VSNET Collaboration and its precedent mailing list-based cooperation (hereafter referred to the VSNET Collaboration throughout this section). Figure 7 shows a comparison of long-term light curves of three representative classes of dwarf novae (cf. Osaki 1996a) drawn from the VSNET observations.[57]

The early work (up to 1996) almost immediately doubled the number of SU UMa-type dwarf novae, by the detection of superhumps (Vogt 1980; Warner 1985; Warner 1995c) during their superoutbursts. In typical SU UMa-type dwarf novae, superoutbursts only occur approximately once per year, and last for ~10 d. The intervals of superoutbursts (supercycles) are not constant, making it difficult to plan a scheduled observation. The observations of SU UMa-type dwarf novae in the past were more or less TOO-type observations (subsection 1.2). We first prepared a comprehensive list of candidates of SU UMa-type dwarf novae, selected by various criteria from the literature and from our own observations. This strategy was in some way similar to the other projects such as:

- High-galactic latitude CV search: Howell, Szkody (1990); Szkody, Howell (1992); Drissen et al. (1994); Augusteijn, Stehle (1995)
- Its descendant concept of “Tremendous Outburst Amplitude Dwarf Novae” (TOADs): Howell et al. (1995); Howell, Szkody (1995); Szkody, Howell (1995); Howell, Skidmore (2000)
- Recurrent Object Programme:[58] e.g. Poyner (1996), and the related extensive work by Vanmunster, Howell (1996)

Although all of these other projects or works basically relied on catalog selections (e.g. Howell et al. 1995) or were based on the (usually poorly known) past activities. Our strategy was different in that: (1) our selection was based on comprehensive and extensive search through all available literature (this work was conducted in collaboration with the VSOLJ), (2) we had examined the Palomar Observatory Sky Survey (POSS) plates and our own systematic CCD survey to get unique identifications and more reliable outburst amplitudes of the objects, (3) we applied a theoretical background (e.g. Osaki 1985; Osaki 1989; Ichikawa, Osaki 1994) for selecting the objects, and (4) we had amateur collaborators to share our interest in monitoring faint dwarf novae with a CCD (subsection 2.3). The combination of these factors brought an unprecedented success in discovering new SU UMa-type dwarf novae. Since the early success story was already described in subsection 4.1, we mainly focus on the recent contributions of the VSNET Collaboration to this field.

6.1.1. *Discovery of ER UMa Stars*

In the canonical picture of SU UMa-type dwarf novae, the mass-transfer (driven by angular momentum loss from the binary) in these systems was believed to be the result of gravitational wave radiation (GWR: cf. Paczynski 1967; Faulkner 1971; Tutukov, Yungelson 1979; Taam et al. 1980; Paczynski, Sienkiewicz 1981), since the fully convective secondary at this ($\leq$2 hr) orbital period ($P_{\rm orb}$) is generally considered improbable to sustain magnetic braking, which plays a major role in longer period ($P_{\rm orb}$ $\geq$3 hr) CVs (Ritter 1976; Ritter 1980; Whyte, Eggleton 1980; Paczynski 1981; Rappaport et al. 1982; Rappaport et al. 1983; Spruit, Ritter 1983; Paczynski, Sienkiewicz 1983; Verbunt 1984; Patterson 1984; Ritter 1985). Since the angular momentum loss by GWR is a unique function of component

---

[50] ⟨http://nis-www.lanl.gov/~asocops/alxhome/⟩.

[51] ⟨http://www-astro.physics.ox.ac.uk/~erik/itw/⟩.

[52] ⟨http://vcssz.mcse.hu/⟩.

[53] ⟨http://cba.phys.columbia.edu/⟩.

[54] ⟨http://www.britastro.com/vss/⟩.

[55] ⟨http://a400.sternwarte.uni-erlangen.de/ftp/bcb/⟩.

[56] We now have a dedicated list for conference announcements: ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-conference/maillist.html⟩.

[57] These data of various kinds of variable stars are freely available (from the URL in subsection 3.3), or upon request to the VSNET administrator (vsnet-adm@kusastro.kyoto-u.ac.jp). There is no specially requested authorization for the usage of these data in scientific publications other than the usual scientific manner of acknowledgement (please refer to the message from the data search engine for a recommended form of reference). The data usually cover observations since 1995, and are particularly convenient for making correlation analysis with other modern observations (e.g. spectroscopy and multiwavelength observations).

[58] ⟨http://www.theastronomer.org/recurrent.html⟩.



Fig. 7. Long-term light curve of three representative classes of dwarf novae (SS Cyg, Z Cam, and SU UMa) drawn from VSNET observations. Portions of the light curves show variation in outburst cycle lengths and quiescent magnitude (SS Cyg), a standstill (Z Cam), and two superoutbursts (SU UMa).

masses and orbital separation, and the masses of the primary white dwarfs and the main-sequence secondary stars only have a small degree of diversity, the SU UMa-type stars were historically considered as an "one parameter system", i.e. the basic properties are determined by a single parameter, P_{orb}.

This concept so widely prevailed that statistics and classifications were usually given following this concept (Warner 1995c; Vogt 1993), and that numerical simulations implicitly assumed this canonical picture (Osaki 1989). In observations, a search for new SU UMa-type dwarf novae was usually restricted to dwarf novae with long recurrence times (the best example being the Recurrent Object Programme). Our discovery, however, of the ER UMa stars completely changed this picture.

ER UMa (=PG 0943+521) was originally classified as a novalike object (Green et al. 1982) selected from its ultraviolet excess (Palomar-Green Survey: Green et al. 1986, see also Ringwald 1993). This object was, together with other PG CVs selected from Green et al. (1982), regularly monitored by the VSOLJ members for potential activities. In 1992, Makoto Iida (VSOLJ) noticed that this object shows dwarf nova-like outbursts (Iida 1994, see *vsnet-history* archive for full details in the public reports circulated in 1992–1993). Because of the presence of long-lasting states of intermediate brightness (which later turned out to be superoutbursts), this star was originally considered to be a Z Cam-type dwarf nova, which is characterized by the presense of standstills in addition to dwarf nova-type outbursts (see e.g. Hellier 2001a, Sect. 5.4; see also Warner, van Citters 1974; Meyer, Meyer-Hofmeister 1983; Oppenheimer et al. 1998).

In early 1994, following the detection of a bright outburst by Gary Poyner, our CCD observations revealed the presence of superhumps. Combined with the visual observations electronically reported, this object was finally identified as an SU UMa-type dwarf nova with an unexpectedly short (43 d) supercycle (T_{s}). The shortest T_{s} before the discovery of ER UMa was 134 d of YZ Cnc (Szkody, Mattei 1984; van Paradijs et al. 1994). It later turned out through a discussion at the Padova CV Conference in 1994[59] that this object was independently studied at least by two groups: by the RoboScope (Honeycutt 1992) team (Robertson et al. 1995) and by Misselt, Shafter (1995). Among all groups, the VSNET team was the first to unambiguously identify the nature of this object (Kato, Kunjaya 1995). Spectroscopic (Thorstensen et al. 1997) as well as photometric (Kato et al. 1995a) identifications of the orbital periods later confirmed this identification.

Once this discovery was announced, new members of the group of ER UMa stars were immediately identified through impetuous real-time competitions through the VSNET public lists: V1159 Ori (Nogami et al. 1995a; Robertson et al. 1995) and RZ LMi (Robertson et al. 1995; Nogami et al. 1995c). It later turned out that the supercycle of V1159 Ori was independently recognized by Patterson et al. (1995b), but was only interpreted within the classical framework. The supercycle of RZ LMi is exceptionally short (19 d), which is the shortest record of supercycles up to now [see also Kato et al. (1999d) for the basic observational review, and Kato et al. (2002e); Kato et al. (2003e) for recent discussions of ER UMa stars]. Later additions to these ER UMa stars include DI UMa (Kato et al. 1996a), an RZ LMi-like system, and IX Dra (Ishioka et al. 2001b), both of which were discovered by the VSNET Collaboration.

From the standpoint of the disk-instability model, these discoveries immediately led to theoretical interpretations (Osaki 1995b; Osaki 1995d), and significantly contributed to the "unified theory of dwarf novae" (Osaki 1996b; Osaki 1996a). These theoretical calculations indicate an unexpectedly (from the canonical picture based on the GWR-driven angular momentum loss) high mass-transfer rate (ER UMa: Osaki 1995b). The extremely short T_{s} of RZ LMi requires an additional (still poorly understood) mechanism (Osaki 1995d). This theoretical effort recently led to ramifications of ideas including the effect of extremely low mass ratio ($q = M_2/M_1$) (Hellier 2001b) or the effect of irradiation (Buat-Ménard, Hameury 2002), which need to be investigated by future work. It has been speculated (Hellier 2001b) that these mechanisms are partly responsible for the manifestation of the unusual properties of still the still poorly understood WZ Sge-type dwarf novae, which will be discussed later (subsection 6.1.2).

The required high mass-transfer is still a mystery. Although there have been a number of suggestions, including the long-term effect of nova explosions (Kato, Kunjaya 1995; Schenker et al. 1998; Patterson 1998; Kolb et al. 2001), which is, in some sense, a modern extension of the "old" discussion of the "nova hibernation" scenario (Shara et al. 1986; Prialnik, Shara 1986; Livio, Shara 1987; Livio et al. 1988; Duerbeck 1992; Naylor et al. 1992; Mukai, Naylor 1995), irradiation-induced mass-transfer feedback (McCormick 1998; McCormick, Frank 1998), none of them has succeeded in explaining the required high mass-transfer rates in ER UMa. Recent observations (Kato 2001a; Fried et al. 1999) suggest that the high mass-transfer rates in ER UMa stars are less likely the result from a secular evolutionary effect, but may be more related to the activity cycle in the secondary star (Bianchini 1988; Bianchini 1990; Ak et al. 2001). The origin of the unusually high activity of ER UMa stars is still an open question.

From the observational side, there has been a systematic search, mainly conducted by the VSNET Collaboration and relevant results communicated to the public in real time, to find intermediate systems between ER UMa stars and classical SU UMa-type dwarf novae. Several interesting objects were recognized as a result: NY Ser = PG 1510+234 (SU UMa-type dwarf nova in the period gap: Nogami et al. 1998b), HS Vir = PG 1341−079 (SU UMa-type dwarf nova with very short outburst recurrence times: Kato et al. 1995d; Kato et al. 1998d; Mennickent et al. 1999; Kato et al. 2001o; Mennickent et al. 2001b), SX LMi (low amplitude SU UMa-type dwarf nova: Nogami et al. 1997b; Wagner et al. 1998; Kato 2001b), CI UMa (SU UMa-type dwarf nova with irregular supercycle behavior: Nogami, Kato 1997), V1504 Cyg (SU UMa-type dwarf nova with a short T_{s}, Nogami, Masuda 1997), V503 Cyg (unusual SU UMa-type dwarf nova with unusual outburst behavior, $T_{\mathrm{s}} \sim 89$ d: Kato et al. 2002e; Harvey et al. 1995), SS UMi (normal SU UMa-type dwarf nova with a short $T_{\mathrm{s}} \sim 84.7$ d: Kato et al. 1998b; Kato et al. 2000a), BF Ara (normal SU UMa-type dwarf nova with the shortest $T_{\mathrm{s}} \sim 84.3$ d: Kato et al. 2001p; Kato et al. 2003b), V344 Lyr (large-amplitude SU UMa-type dwarf nova with a short supercycle: Kato 1993; Kato et al. 2002j), MN Dra = Var73 Dra (SU UMa-type dwarf nova with $T_{\mathrm{s}} \sim 60$ d: Nogami et al. 2003d).

---

[59] See also Honeycutt et al. (1995); the object was originally presented in the conference as the best observational proof for the mass-transfer burst model of dwarf nova outbursts. Our detection of superhumps was presented in a later talk by Osaki (1995a), which invoked a hot discussion, against the original topics to be presented by this talk, regarding the true nature of ER UMa. At the time of this conference, the tone of the audience was even skeptical about the presence of an SU UMa-type dwarf nova with such a short supercycle.

The evolution of superhumps in ER UMa stars is also known to be unusual. Kato et al. (1996d) reported the early presence of large-amplitude superhumps, which later turned out to be part of an unexpected early phase reversal of superhumps (Kato et al. 2003e). This phenomenon resembles the so-called late superhumps (Haefner et al. 1979; Vogt 1983; van der Woerd et al. 1988; Hessman et al. 1992) seen in late stages of superoutbursts in SU UMa stars, but the striking difference is that they appear in the early stage of the superoutburst. The origin of this phase reversal is not yet understood. Gao et al. (1999) reported some peculiar features in superhumps of ER UMa. Masetti, Regos (1997) suggested a possible link between ER UMa stars X-ray transients by comparing the evolution of their superhumps.

To summarize, the discovery of ER UMa stars brought a revolutionary turning point of dwarf nova studies: the SU UMa-type dwarf novae are no longer "one-parameter systems" (see e.g. Warner 1998; Nogami 1998). Research on short-P_{orb} CVs (mainly SU UMa-type dwarf novae) is continuously broadening, and is now becoming one of "the mainstreams" of CV research.

6.1.2. *WZ Sge-Type Dwarf Novae*

WZ Sge-type dwarf novae are a peculiar subtype of SU UMa-type dwarf novae.[60] The properties of WZ Sge-type dwarf novae include: (1) long (∼ 10 yr or more) outburst recurrence time, (2) large (∼ 8 mag) outburst amplitude, (3) very few, or sometimes no, occurrence of (isolated) normal outbursts, (4) presence of "early superhumps", which are modulations having periods very close to the orbital periods, during the earliest stage of superoutbursts, and (5) frequent occurrence of post-superoutburst rebrightening (see Kato et al. 2001l and Kato 2002b for modern observational reviews).

These properties are difficult to explain, even with the recent progress of the disk-instability theory, and the WZ Sge-type dwarf novae have been continuously providing challenging problems to both theoreticians and observers. One of the main difficulties reside in their extremely long recurrence times. If one assumes the standard disk-instability model, the recurrence time is limited by the diffusion time in quiescence (Ichikawa, Osaki 1994; Osaki 1996a). In order to avoid a thermal instability resulting from this diffusion, one needs to assume an extremely low viscosity parameter in quiescence ($\alpha_C <$ 0.00005, Smak 1993; $\alpha_C < 0.003$, Osaki 1995c). Although the origin of such a low viscosity is becoming positively resolved by considering a very cold disk with a low electronic conductivity (Meyer-Hofmeister et al. 1998), there still exist a number of arguments to avoid an extremely low quiescent viscosity. For example, Lasota et al. (1995), Warner et al. (1996) assumed evaporation/truncation of the inner disk to prevent thermal instability to occur. Hameury et al. (1997) and Buat-Ménard, Hameury (2002) presented slight modifications of these ideas. The models, however, are expected to show much shorter outburst lengths than in the low-α_C model (Osaki 1995c). Some models thereby assume an enhanced mass-transfer during a superoutburst (as originating from Patterson et al. 1981 and extended by Patterson et al. 2002), there is, however, no concrete observational evidence supporting this supposed enhanced mass-transfer (Osaki, Meyer 2003). A careful analysis of the observations of the 2001 outburst of WZ Sge also supports this lack of mass-transfer (R. Ishioka et al. in preparation). Some authors ascribed the existence of a brown-dwarf secondary star to the origin of the required low α_C (Meyer, Meyer-Hofmeister 1999). This possibility, together with the expectations from the theoretical viewpoint (e.g. Paczyński 1971), led to a wide interest to search for brown dwarfs in WZ Sge-type dwarf novae (Howell et al. 1997; Politano et al. 1998; Ciardi et al. 1998; van Teeseling et al. 1999; Patterson 2001; Howell, Ciardi 2001; Mennickent et al. 2001a; Kato et al. 2002p; Mennickent, Diaz 2002; Littlefair et al. 2003). Although earlier reports tended to suggest the presence of a brown dwarf, the evidence is less clear from more recent detailed studies (Steeghs et al. 2001a; Kato et al. 2002p; Mennickent, Diaz 2002; Howell et al. 2002; Littlefair et al. 2003). Most recently, the observational lack of CVs having brown dwarf secondaries is even becoming a serious problem (Barker, Kolb 2003). In all aspects of disk-

---

[60] Historically, the concept of WZ Sge-type dwarf novae referred to large-amplitude, rarely outbursting dwarf novae (cf. Bailey 1979; Downes, Margon 1981; Patterson et al. 1981; O'Donoghue et al. 1991). This extension of this standpoint led to the concept of Tremendous Outburst Amplitude Dwarf Novae (TOADs), mainly proposed by Howell et al. (1995). WZ Sge-type dwarf novae were sometimes even referred to as extreme TOADs (cf. Howell et al. 2003). The modern viewpoint, however, is that the outburst amplitude is one of the general properties of WZ Sge-type dwarf novae, rather than the definition. WZ Sge-type dwarf novae are now better recognized as an extreme population of SU UMa-type dwarf novae (Kato et al. 2001l, see also the discussion in relation to the "TOAD" concept: Kato et al. 2003f). We therefore primarily use the term WZ Sge-type dwarf novae rather than TOADs.



Fig. 8. CCD light curve of the 2001 outburst of WZ Sge by the VSNET Collaboration.

instability problems, origin of disk viscosity, and late-stage evolution of compact binaries, WZ Sge-type dwarf novae continue to be key objects.

Since the outbursts of WZ Sge-type dwarf novae are quite rare, these objects have best illustrated the ability and the achievement of the VSNET as the real-time network. Before the VSNET was established, most (presumable) WZ Sge-type outbursts were only poorly studied (e.g. AL Com: Bertola 1964, UZ Boo: Bailey 1979, PQ And: McAdam et al. 1988; Hurst et al. 1988a; Hurst et al. 1988b; Hurst, Young 1988, GW Lib: Maza, Gonzalez 1983, other systems: Richter 1992). The establishment of the VSNET has been one of the greatest steps toward understanding the WZ Sge-type dwarf novae. In particular, without the collaboration with the VSOLJ and without the VSNET alerts, the early part of the 2001 outburst of WZ Sge would have still remained a mystery.

The earliest work on WZ Sge-type dwarf novae was on HV Vir in 1992. Although this object was originally recorded as a classical nova in 1929 (Schneller 1931; Duerbeck 1984), we started monitoring for a future potential outburst as a candidate WZ Sge-type dwarf nova (Kato et al. 2001l). This outburst was discovered by Patrick Schmeer (Schmeer et al. 1992), and was immediately relayed through the alert network. The VSNET team was the first to record periodic modulations (early superhumps) in the light curve.[61] The final result on this superoutburst was published as Kato et al. (2001l), which has been chosen as a best modern reference on new WZ Sge-type stars in Trimble, Aschwanden (2003). After ~10 yr, the object again went into superoutburst, which was observed in detail by the VSNET Collaboration (Ishioka et al. 2003).

The next object was UZ Boo in 1994 (Iida, York 1994; Szentasko et al. 1994). The outburst detection was also immediately relayed through the VSNET alert network. Due to the short visibility, only a preliminary superhump period of 0.0619 d was obtained (Kato et al. 2001l), although the post-superoutburst rebrightenings reported through the VSNET have raised a possible link between WZ Sge-type dwarf novae and soft X-ray transients (SXTs, or X-ray novae, subsection 6.4) (Kuulkers et al. 1996). This suggestion has been more substantiated by Uemura et al. (2000c), Kuulkers (2000), Kuulkers (2001); this relation is becoming one of the major contemporary topics in SXT outbursts.

In 1995 and 1996, outstanding WZ Sge-type outbursts of AL Com and EG Cnc occurred. These outburst detections were immediately relayed through the VSNET, and produced a wealth of scientific results as already introduced in subsection 4. AL Com again underwent an superoutburst in 2001. A timely outburst announcement by Steve Kerr on VSNET enabled early-time observations when the object was still rising. The detection of growing early superhumps was already reported (Ishioka et al. 2002b), which was actually the first detection of growing early superhumps before the 2001 spectacular outburst of WZ Sge itself.

---

[61] This was immediately reported to the CBAT (see *vsnet-history* archive), but was not published for an unknown reason. See Kato et al. 2001l more details.

In late 2000, another spectacular outburst of RZ Leo occurred, which had long been suspected as a candidate WZ Sge-type dwarf nova (McNaught 1985; Mattei et al. 1985; Cristiani et al. 1985; Richter 1985). The VSNET Collaboration succeeded in detecting both early superhumps and ordinary superhumps (Ishioka et al. 2000a; Ishioka et al. 2001c), giving credence to the WZ Sge-type nature of this dwarf nova.

The unexpected outburst of the prototype WZ Sge in 2001 was one of the greatest astronomical phenomena in recent years. It was detected by a Japanese amateur observer, Tomohito Ohshima, was immediately relayed to the VSNET, enabling early coverage (Ishioka et al. 2001e; Ishioka et al. 2002c; Kuulkers et al. 2002), which was one of the greatest achievements of the VSNET as an alert network (see figure 8). This outburst produced a burst of scientific results from a number of researchers, both ground-based and satellite-borne: detection of early superhumps (Kato et al. 2001b), growth of ordinary superhumps (Kato et al. 2001j; Kato et al. 2001i), spectroscopic detection of spiral patterns (Steeghs et al. 2001b; Baba et al. 2001) whose details have been published in Baba et al. (2002), "real-time" modeling of the outburst (Cannizzo 2001), Chandra observation (Wheatley et al. 2001), HST observation (Knigge et al. 2002), far ultraviolet spectroscopy (Long et al. 2003), infrared spectroscopy (Howell et al. 2003), extensive photometry (Ishioka et al. 2002b; Patterson et al. 2002; R. Ishioka et al. in preparation), and theoretical modeling of the superhump light curve (Osaki 2003).

Other WZ Sge-type dwarf novae observed and reported by the VSNET Collaboration include: UW Tri (Kato et al. 2001h), LL And (Kato 2003b; data cited in Howell, Hurst 1994, Howell, Hurst 1996), V2176 Cyg (Novák et al. 2001), CG CMa (Kato et al. 1999b), V592 Her (Kato et al. 2002p). Among them, V2176 Cyg showed a "dip" phenomenon, noted for first time since the 1995 outburst of AL Com (Nogami et al. 1997a), and V592 Her was confirmed to be a dwarf nova with an exceptionally large outburst amplitude.

The VSNET Collaboration has first systematically demonstrated that all well-observed WZ Sge-type dwarf novae show "early superhumps" during the earliest stage of their superoutbursts (which may be the best defining characteristic of WZ Sge-type dwarf novae: Kato 2002b). These early superhumps are usually double-wave (sometimes more complex) variations, which have periods extremely close to the orbital periods (e.g. Ishioka et al. 2002b). Although there is a historical (Patterson et al. 1981) and a modern version (Patterson et al. 2002) interpretation, that the phenomenon results from an enhanced mass-transfer, it is now understood as the result of some sort of resonance on the disk (2:1 resonance: Osaki, Meyer 2002 or vertical 2:1 resonance: Kato 2002b).

The identification of RZ Leo as a WZ Sge-type dwarf nova (Ishioka et al. 2001c), is a surprise from this point of view, since the superhump period (0.078529 d) of RZ Leo is anomalously long compared to the canonical picture of WZ Sge-type dwarf novae. This identification indicates that neither a brown-dwarf secondary (Meyer, Meyer-Hofmeister 1999) nor an extreme mass-ratio, enabling 2:1 resonance (Osaki, Meyer 2002), may be a necessary condition for the WZ Sge-type outburst phenomenon. This implication is presently under discussion for WZ Sge-type dwarf novae (cf. Kato 2002b; Osaki, Meyer 2002). As concerns early superhumps, Kato (1997a) detected a smooth transition from the orbital to superhump period in a more usual SU UMa-type dwarf nova, T Leo. This phenomenon may be somehow related to the evolution of early superhumps (Kato et al. 1996b).

Several large-amplitude SU UMa-type dwarf novae share some common properties with WZ Sge-type dwarf novae, particularly in the lengthening of the superhump period. The periods of the superhumps are usually not constant, but show a significant period derivative ($P_{\rm dot}$). The $P_{\rm dot}$ are usually negative in classical SU UMa-type dwarf novae (Warner 1985; Patterson et al. 1993; for a recent progress, see Kato et al. 2003h). This negative $P_{\rm dot}$ is usually considered to be a result of a decrease in the angular velocity of precession of a shrinking eccentric disk (Osaki 1985). The decrease may also be a result of inward propagation of the eccentricity wave (cf. Lubow 1991a; Lubow 1991b; Lubow 1992). A small number of SU UMa-type dwarf novae, notably many WZ Sge-type dwarf novae, are known to show positive $P_{\rm dot}$. This effect was first clearly detected in V1028 Cyg (Baba et al. 2000), although this phenomenon first appeared in solid publications on SW UMa (Semeniuk et al. 1997; Nogami et al. 1998a), V485 Cen (Olech 1997b), and on the WZ Sge-type star AL Com (Pych, Olech 1995b; Howell et al. 1996; Nogami et al. 1997a). The SU UMa-type dwarf novae newly identified by the VSNET Collaboration as having positive $P_{\rm dot}$ include: HV Vir (WZ Sge-type star, Kato et al. 2001l), WX Cet (Kato et al. 2001c), EG Cnc (WZ Sge-type star, Kato et al. 1997), and XZ Eri (Uemura et al. 2003c). The true origin of this phenomenon is not yet well understood (Baba et al. 2000; Kato et al. 2001l; Ishioka et al. 2003; Kato et al. 2003h; Kato et al. 2003b).

Post-superoutburst rebrightenings are also a renowned feature of the WZ Sge-type dwarf novae (Kato et al. 1997; Patterson et al. 1998; Ishioka et al. 2003), for which Osaki et al. (1997), and Osaki et al. (2001) presented an interpretation based on the slow viscosity decay in the early post-superoutburst state. This mechanism would require a mass reservoir in the outer disk (Osaki, Meyer 2003), whose original observational implication was proposed by Kato et al. (1998c). The existence, or non-existence, of post-superoutburst rebrightenings have been systematically studied by the VSNET Collaboration in almost all SU UMa-type dwarf nova. Recent examples include GO

Com (Kato et al. 1995c), WX Cet (Kato 1995f; Kato et al. 2001c), V1028 Cyg (Baba et al. 2000), V725 Aql (Uemura et al. 2001c), and in the unusual system (see subsection 6.1.3) EI Psc (Uemura et al. 2002e). The statistics is presented in Kato et al. (1998c).

A missing link between WZ Sge-type dwarf novae and ordinary SU UMa-type dwarf novae has been sought. CT Hya (Nogami et al. 1996; Kato et al. 1999a) was suggested to be one such system, though a more recent statistical analysis suggests a more rigid segregation between WZ Sge-type dwarf novae and ordinary SU UMa-type dwarf novae (Kato et al. 2003f). Several systems possibly related to WZ Sge-type dwarf novae have been also studied by the VSNET Collaboration: CC Scl = RX J2315.5−3049 (Ishioka et al. 2001a).

Long $P_{\rm orb}$ SU UMa-type dwarf novae with rare outbursts have also been systematically studied by the VSNET Collaboration. The objects include: EF Peg (Kato, Takata 1991; Kato 2002c) and V725 Aql (Nogami et al. 1995d; Uemura et al. 2001c). The origin of the necessary low mass-transfer rate in such long $P_{\rm orb}$ systems is still a problem.

6.1.3. Ultra-Short Period Systems and Helium Cataclysmic Variables

From the standard evolutionary scenario of compact binaries, there should be a "period minimum" (e.g. Paczyński 1971) at which the mass-losing secondary star becomes degenerate and the binary period starts to lengthen. This period is observationally determined to be ∼1.3 hr, which is about 10% longer than the theoretical predictions (Paczyński 1971; Kolb, Baraffe 1999; King et al. 2001; King et al. 2002). This discrepancy has not been yet resolved, although several attempts have been made (e.g. Renvoizé et al. 2002; Barker, Kolb 2003) to reconcile theory with observation.

There exist, however, hydrogen-rich systems with periods well below this theoretical minimum period. The "classical" object is V485 Cen (Augusteijn et al. 1993; Augusteijn et al. 1996; Olech 1997b); the faintness of this object, however, prevented a detailed observational study. In the past few years, the VSNET team found a hydrogen-rich, nearby bright system (EI Psc = 1RXS J232953.9+062814) having a short period comparable to V485 Cen (Uemura et al. 2001b; Uemura et al. 2002f; Uemura et al. 2002e). Both the radial-velocity study (Thorstensen et al. 2002b) and the superhump period analysis (Uemura et al. 2002e; Skillman et al. 2002) independently confirmed that the secondary star of this system is more massive than what is expected for this orbital period, suggesting that the mass donor is an evolved core with a thin hydrogen envelope. From this finding, Uemura et al. (2002f) suggested, following the evolutionary calculations by Podsiadlowski et al. (2003), that both EI Psc and V485 Cen can be ancestors of helium CVs (or AM CVn stars, Warner 1995a; Solheim 1995) consisting of a white dwarf and a mass-losing helium white dwarf. If this interpretation is confirmed, this object would become the first direct observational evidence that helium CVs are descendants of a certain class of CVs with hydrogen-rich appearance (cf. Webbink 1984; Tutukov, Yungelson 1996; Nelemans et al. 2001; Yungelson et al. 2002). This object, given its proximity and relatively heavy component masses, is also considered to be an excellent candidate for next generation experiments of directly detecting gravitational wave radiation (Hils, Bender 2000; Strohmayer 2002).

In clarifying the nature of EI Psc, proper motion studies played an independent important role in identifying the object as a nearby object. This finding was soon confirmed by later researchers (Biggs, Walsh 2001), and the same technique has been applied to different sorts of objects by the VSNET Collaboration (RX J2309.8+2135: Kato, Yamaoka 2002, V379 Peg: Kato et al. 2002u, CW Mon: Kato et al. 2003j). This application of astrometry soon became the global standard in studying CVs and related systems (Kato et al. 2001t, see also recent entries of a survey of proper motions in Downes et al. online CV catalog[62]).

Helium CVs have been one of the best observed targets by the VSNET Collaboration. Among them, CR Boo has been identified as the first "helium ER UMa star" with a supercycle of 46.3 d (cf. subsection 6.1.1) by Kato et al. (2000c). The VSNET team also joined a campaign to study superhumps in CR Boo (Patterson et al. 1997). Kato et al. (2000d) identified a similar supercycle in the helium CV, V803 Cen. Kato et al. (2001m) further identified standstills in V803 Cen, which were initially suggested for CR Boo in Kato et al. (2000c). Kato et al. (2001k) later detected a transition of CR Boo to a state of a short supercycle (14.7 d), which they called "the second supercycle". This phenomenon is still difficult to explain. Kato et al. (2003i) studied V803 Cen for its long-term behavior and its 2003 June superoutburst, and revealed that the object (and probably also CR Boo) shows outburst behavior similar to WZ Sge. From these studies, both CR Boo and V803 Cen have been well-established "helium counterparts" to hydrogen-rich SU UMa-type dwarf novae, in contrast to the traditional VY Scl-type (variable mass-transfer rate from the secondary, see subsection 6.3.3) explanation of high and low states in these systems (Warner 1995a; Woudt, Warner 2003b). This interpretation is perfectly in line with the dwarf nova-type interpretation (Tsugawa, Osaki 1997), although this interpretation was not originally correctly applied to observation (Patterson et al. 1997; Patterson et al. 2000b).

Outburst detections of other helium dwarf novae (e.g. KL Dra = SN 1998di) have been also announced through the VSNET, and provided necessary fundamentals[63] for detailed research (e.g. Jha et al. 1998b; Samus 2000b; Wood et al. 2002).

6.1.4. Eclipsing Dwarf Novae

Since CVs are close binary systems, high-inclination systems show eclipses. The presence of eclipses in CVs historically provided most crucial information about the geometry and fundamental physics of the accretion disk or the accretion stream (Walker 1954; Walker 1955; Kraft 1958; Greenstein, Kraft 1959; Kraft 1962; Warner, Nather 1971) and the clarification of the cause of outbursts (Osaki 1974; Smak 1976; Smak 1984; Krzeminski, Vogt 1985).

---

[62] ⟨http://icarus.stsci.edu/~downes/cvcat/⟩.

[63] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/DNe/kldra.html⟩.

In recent years, the eclipse mapping technique (Horne 1985) has been used to geometrically resolve the accretion disk by numerically modeling the eclipse light curve (and sometimes line variations) of CVs (Horne, Stiening 1985; Marsh, Horne 1987; Wood et al. 1989a; Baptista, Steiner 1991; Baptista, Steiner 1993). This and analogous methods have also been used to study the time-evolution of the accretion disk during dwarf nova outbursts (Horne, Cook 1985; Wood et al. 1989b; Rutten et al. 1992; Wolf et al. 1993; Bobinger et al. 1997), discuss the presence of a spiral pattern (which may be theoretically predicted spiral shocks: Sawada et al. 1986; Spruit et al. 1987; Matsuda et al. 1990; Matsuda et al. 2000) in dwarf novae (Baptista, Catalán 2001; Baptista, Catalán 2000; Baptista et al. 2000), for mapping of the superoutbursting disks, (Bruch et al. 1996), to spectrally resolve the accretion disk (Rutten et al. 1993; Rutten et al. 1994), and to directly obtain physical parameters of the accretion disk (Vrielmann et al. 1999; Vrielmann et al. 2002).

The VSNET Collaboration played an important role in studying eclipsing CVs, especially eclipsing dwarf novae. The initial efforts were made to follow the eclipses of the rising phase of an IP Peg outburst (Nogami 1996). The VSNET alert lists provided an initiative role of systematic studies when the northern new eclipsing dwarf nova EX Dra (= HS 1804+6753) was discovered (Nogami 1996). This action was soon extended to observe rare outbursts of an eclipsing SU UMa-type dwarf nova, DV UMa (Nogami et al. 2001b). The 1995 outburst of HT Cas was also spectacular. This is a well-known eclipsing dwarf nova (cf. Patterson 1981; Zhang et al. 1986; Wood et al. 1995; Mukai et al. 1997). In 1995, the VSNET team received a request for an optical ground-based campaign coordinated with the Hubble Space Telescope (HST). The observation by the VSNET team succeeded in correcting the eclipse ephemeris, which is readily reflected on the HST observing schedule, when the object suddenly jumped into an outburst! Thanks to this coincidence, we were able to obtain eclipse information only two days prior to the outburst maximum, which precluded the enhancement of the hot spot as expected from the mass-transfer burst (Baba et al. 1999). The eclipses during this outburst was also followed by another group (Ioannou et al. 1999), who observed this object in response to this outburst detection. The results of eclipse mapping, together with later outburst observations, have been recently reported (Baba et al. 2003b).

IR Com (= S 10932), a system very similar to HT Cas, has also been extensively studied by the VSNET team. In particular, we detected the 1996 January outburst, and succeeded in taking the earliest eclipse observations (Kato et al. 2002a), The true nature of this object had remained unclear before (Richter, Greiner 1995; Richter et al. 1997). The real-time circulation of this outburst detection and eclipse information enabled a third-party follow-up observations (Richter et al. 1996).

In most recent years, the VSNET Collaboration discovered a deeply eclipsing bright SU UMa-type dwarf nova (IY UMa = Tmz V85) in the northern hemisphere (Uemura et al. 2000d; Uemura et al. 2000b; Samus 2000a). For the first time in history, this observation yielded the simultaneous discovery of superhumps and eclipses. This system, the only bright normal SU UMa-type dwarf novae suitably situated for northern telescopes, has been proposed as the best candidate object for next generation detectors on huge telescopes (cf. Moon et al. 2001; Steeghs et al. 2003; Reynolds et al. 2003). This and the subsequent outbursts were followed by a number of teams, resulting in rich physical insights (Patterson et al. 2000a; Wu et al. 2001; Stanishev et al. 2001; Rolfe et al. 2001; Rolfe et al. 2002; Smak 2003). With strong emission lines of HeII and CIII/NIII in outburst (Wu et al. 2001; such objects are known to be quite rare: Morales-Rueda, Marsh 2002), this object is a good candidate for spatially resolving a superoutbursting disk by the emission-line eclipse mapping method (Makita, Mineshige 2002), as well as with classical Doppler tomography of the velocity field (Marsh, Horne 1988). We also succeeded in identifying the supercycle (Kato et al. 2001n).

The other outstanding object is DV UMa, which was observed during the entire stage of the 1999 December superoutburst following the outburst report by Timo Kinnunen.[64] This observation first fully covered the early evolution of eclipses in this rarely outbursting system.

In 2002 February, a collaborative effort on GY Cnc = RX J0909.8+1849 led to the discovery of the eclipsing nature of this dwarf nova (Gänsicke et al. 2000; Kato et al. 2000f; Shafter et al. 2000; Thorstensen 2000). Eclipse observations during the 2001 November outburst revealed the noticeable absence of the hot spot during the late stage of an outburst (Kato et al. 2002f). This observation suggested that GY Cnc may be the first long-$P_{\rm orb}$ object sharing common properties with HT Cas and IR Com.

Recent detailed outburst (or superoutburst) observations of eclipsing dwarf novae include: XZ Eri (Uemura et al. 2003c) and OU Vir (R. Ishioka et al. in preparation). Both stars show prominent superhumps as well as eclipses. XZ Eri is the first eclipsing SU UMa-type dwarf nova with a positive period derivative (Uemura et al. 2003c). OU Vir is another object continuously receiving world-wide attention (Vanmunster et al. 2000b; Mason et al. 2002), for which we succeeded in determining the first reliable orbital and superhump periods. V2051 Oph is another eclipsing CV, which had been thought to be a low-field polar (Warner, Cropper 1983; Wenzel 1984; Warner, O'Donoghue 1987) for which the VSNET Collaboration first provided unambiguous clarification of its SU UMa-type nature by securely detecting superhumps and supercycles (Kiyota, Kato 1998; Kato et al. 2001n). With the help by this clear identification, this object has also been receiving special attention both with ground-based and satellite-borne observations (Baptista et al. 1998; Bruch 2000; Steeghs et al. 2001c; Vrielmann et al. 2002)

---

[64] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/DNe/dvuma9912.html⟩.

The recovery and clarification of the nature of the "lost" dwarf nova V893 Sco is another noteworthy achievement by the VSNET Collaboration. This object had long been lost, when Katsumi Haseda (VSOLJ) reported an outbursting object (at a nominally different position from that of the originally reported V893 Sco) to the VSNET. After careful research of the discovery material, as a part of confirmatory process of a new variable star (subsection 5.2), this newly reported object was eventually identified with the lost V893 Sco (Kato et al. 1998a). This was only the beginning of the story; the object soon turned out to be the brightest, and presumably one of the nearest, eclipsing dwarf novae below the period gap (Thorstensen 1999; Matsumoto et al. 2000; Bruch et al. 2000). This object has been extensively studied since its recovery (Mason et al. 2001; Kato et al. 2002h).

CW Mon (Kato et al. 2003j) shows grazing eclipses during certain stages of outbursts. Together with the transient appearance of pulsed signals, the presence of a premaximum halt in the outburst light curve and relatively strong X-ray radiation, this object has been suspected to be an intermediate polar (see also subsection 6.3.1).

6.1.5. Super-Quasi-Periodic Oscillations

Quasi-periodic oscillations (QPOs) are short-period, quasi-periodic oscillations widely observed in accreting binary systems including CVs (cf. Warner 1995b). QPOs in CVs are usually subdivided into two classes. One is dwarf nova oscillations (DNOs) observed during outbursts of dwarf novae. DNOs have short periods (usually 19–29 s) and long coherence times (Robinson 1973; Szkody 1976; Patterson 1981; Hildebrand et al. 1980). The other is QPOs, which have longer (40 to several hundred seconds) and shorter coherence times (usually less than ∼10 wave numbers). We discovered a potentially new class of QPOs (super-QPOs) during the 1992 superoutburst of SW UMa (Kato et al. 1992a). These super-QPOs have long (several hundred seconds) periods and long coherent times (more than several tens of wave numbers). In some cases (Kato et al. 1992a), the amplitude can be quite large (up to 0.2 mag). The most outstanding feature of super-QPOs is that they are observed only during certain stages of SU UMa-type superoutbursts. In SW UMa (1992) and EF Peg (Kato 2002c), the super-QPOs were observed during the growing stage of superhumps. A similar, but less striking, probable appearance of super-QPOs was also recorded by the VSNET Collaboration during the early stage of an superoutburst of NSV 10934 (Kato et al. 2003d).

During the 2000 superoutburst of SW UMa, similar super-QPOs temporarily appeared during the decay phase of a superoutburst (vsnet-alert 4331).[65] These observations suggest that the appearance of super-QPOs is closely related to the growth and decay of superhumps, or related to the existence of heating/cooling waves (Yamasaki et al. (1995) proposed that some sort of QPOs can be an excitation of trapped oscillations around the discontinuity of physical parameters).

Different interpretations have also been suggested. For example, Warner, Woudt (2002) suggested that the super-QPOs may be a result of interaction between the weak magnetism of the white dwarf and some kind of wave in the inner accretion disk. Although this explanation would be compatible with the suggested presence of a weak magnetic field in SW UMa (Shafter et al. 1986; Robinson et al. 1987; Szkody et al. 1988; Rosen et al. 1994), there would be a need for a different mechanism to explain why these super-QPOs only appear only during temporary stages of superoutbursts.

In EF Peg (Kato 2002c), a rapid decrease in the periods of super-QPOs was recorded. From this finding, Kato (2002c) suggested a rapid removal of angular momentum from an orbiting blob in the accretion disk, via a reasonable viscosity in a turbulent disk. The origin of super-QPOs is still an open question, but their prominent profile is expected to provide crucial information about the origin of QPOs in CVs (Warner, Woudt 2002).

6.1.6. Low States in Dwarf Novae

Although some nova-like CVs (VY Scl-type stars, subsection 6.3.3) are best known to show "low states", during which mass-transfer from the secondary is reduced (Warner 1995b; Robinson et al. 1981), this phenomenon has not been clearly confirmed to occur in dwarf novae. Although there have been claims for "low states" (HT Cas: Zhang et al. 1986; Wood et al. 1995; Robertson, Honeycutt 1996, IR Com: Richter, Greiner 1995; Kato et al. 2002a, WW Cet: Ringwald et al. 1996; BZ UMa: Kaluzny 1986), it is not evident whether or not these phenomena directly reflect a reduced mass-transfer from the secondary, since the state change in the disk (especially the viscosity parameter) would reproduce similar phenomena. Extensive studies on selected well-observed dwarf novae (e.g. Honeycutt et al. 1998; Oppenheimer et al. 1998) have found no evidence for the long-term variation of mass-transfer.

During the extensive work by the VSNET Collaboration, we discovered that the Z Cam star RX And underwent a deep fading in 1996 September (vsnet-obs 3750).[66] This fading (and the lack of outbursts) lasted until 1997 January, which yielded the first unambiguous detection of a temporarily reduced mass-transfer in dwarf novae. This phenomenon was thoroughly studied by Kato et al. (2002i). Careful research on the historical light curve indicated that similar phenomena were sporadically observed in RX And (Kato et al. 2002i; Schreiber et al. 2002), but had been overlooked mainly because of the confusion of the true quiescent identification, and the lack of real-time circulation of information. The VSNET Collaboration succeeded in detecting such a phenomenon through real-time, regular monitoring of light curves of dwarf novae, and through a prompt reaction when an anomalous change was observed (the deep quiescence of RX And was originally confirmed by our own CCD observation). This is another aspect how effectively the VSNET alert network worked, besides other outburst-type transient events. This detection, announced world-wide, thereby led to a prompt HST observation (Sion et al. 2001), which revealed the presence of a hot white dwarf. Kato (2003c) further reported the detection of short fading episodes in RX And and SU UMa, which may be a result of temporary reduction of mass-transfer rate.

---

[65] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/DNe/swuma00.html⟩.

[66] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/obs3000/msg00750.html⟩.

6.1.7. *Other Dwarf Nova Works*

Other dwarf nova-related works, not covered by the above subsections, by the VSNET Collaboration include:

(a) Dwarf novae in the period gap: GX Cas, V419 Lyr (Nogami et al. 1998c) (see subsection 6.1.1 for NY Ser and MN Dra)
(b) Time-variation in more usual SU UMa-type dwarf novae and candidates: AW Gen (Kato 1996b), RZ Sge (Kato 1996a), V1113 Cyg (Kato et al. 1996e; Kato 2001h), CC Cnc (Kato, Nogami 1997a; Kato et al. 2002o), VZ Pyx (Kato, Nogami 1997b), CY UMa (Kato 1997b; Kato, Matsumoto 1999a), PU Per (Kato, Matsumoto 1999b), AQ Eri (Kato, Matsumoto 1999c; Kato 2001g), CT Hya (Kato et al. 1999a), V364 Peg (Kato, Matsumoto 1999d), QW Ser (Kato, Uemura 1999b; Nogami et al. 2003c), BZ UMa (Kato 1999e), CI Gem (Kato, Schmeer 1999), TY Vul (Kato, Uemura 1999d), KV Dra (Nogami et al. 2000), V844 Her (Kato, Uemura 2000b), QY Per (Kato et al. 2000e), TY Psc (Kunjaya et al. 2001; Kunjaya et al. 2003), V630 Cyg (Nogami et al. 2001a), V369 Peg (Kato, Uemura 2001c), UV Gem, FS And, AS Psc (Kato, Uemura 2001f), RX Cha (Kato et al. 2001a), YZ Cnc (Kato 2001f), IR Gem (Kato 2001i), FT Cam (Kato et al. 2001t; Kato 2002d), GZ Cnc, NSV 10934 (Kato et al. 2001r; Kato et al. 2002b), DM Dra (Kato et al. 2002n), SU UMa (Kato 2002a), DM Lyr (Nogami et al. 2003a)
(c) Time-variation in SS Cyg-type dwarf novae: V1008 Her (Kato, Uemura 1999c), V1101 Aql (Kato et al. 1999c), DK Cas (Kato, Uemura 2001a), HH Cnc = Tmz V36 (Kato, Uemura 2001e), IS Del (Kato 2001c), IZ And (Kato 2001e), DX And (Kato, Nogami 2001a), AH Eri (Kato, Nogami 2001b), CG Dra (Kato, Nogami 2001c),
(d) Standstills of Z Cam-type dwarf novae: VW Vul (Kato 1999f), AT Cnc (Nogami et al. 1999, Doppler tomography; Kato et al. 2001q), Z Cam (Kato 2001d), HL CMa (Kato 2002e), FX Cep (Kato et al. 2001f), V363 Lyr (Kato et al. 2001g), EY Cyg (Kato et al. 2002m), IW And (unusual Z Cam star: Kato et al. 2003c),
(e) Quiescent dwarf novae: UV Per (Kato 1990b), GO Com (Kato, Hirata 1990a)
(f) Classification: BF Eri (Kato 1999a; Kato, Uemura 2000a), LX And (Uemura et al. 2000e; Kato 2002f), HP And (with Subaru, Nogami et al. 2003b)
(g) Statistics and compilation: Nogami et al. (1997b); Kato et al. (2003f)

Most of the work was done with the help of the alert network and the collaboration described in subsections 2.3 and 4.1.

6.1.8. *Other Global Contributions*

As the VSNET has been mediating an enormous number of outburst alerts of dwarf novae since its very early history, these alerts, as well as long-term observations, contributed to world-wide dwarf nova studies by different teams. Since they are so numerous, we only list representative ones:

- V485 Cen: Olech (1997a); Olech (1997b)
- PV Per: Vanmunster (1997)
- TU Crt: Mennickent et al. (1998)
- KS UMa: Watanabe (1998a); Olech et al. (2003b)
- KV Dra: Vanmunster et al. (2000a)
- WY Tri: Vanmunster (2001)
- KX Aql: Tappert, Mennickent (2001)
- XY Psc: Henden et al. (2001)
- AH Her: Spogli et al. (2001); Spogli et al. (2002)
- CVs in the 2MASS survey: Hoard et al. (2002b)
- V844 Her: Thorstensen et al. (2002c)
- V2051 Oph: Vrielmann et al. (2002); Vrielmann, Offutt (2003)
- QZ Ser: Thorstensen et al. (2002a), another peculiar dwarf nova discovered by Katsumi Haseda (Had V04), and announced in collaboration with the VSNET (subsection 5.2)
- IP Peg: Baptista et al. (2002)
- RX J0944.5+0357: Mennickent et al. (2002)
- V1504 Cyg: Pavlenko, Dudka (2002)
- GZ Cnc: Tappert, Bianchini (2003)
- V1141 Aql: Olech (2003)
- QW Ser: Olech et al. (2003a)
- FS Aur: Tovmassian et al. (2003)
- EM Cyg: Spogli et al. (2003)
- short-period dwarf novae: Thorstensen, Fenton (2003)
- faint CVs survey: Mason, Howell (2003)
- SU UMa stars: Patterson et al. (2003)

6.2. *Novae and Recurrent Novae*

6.2.1. *Classical Novae and Related Objects*

From the beginning of the VSNET, novae and recurrent novae have been widely studied as one of the classical representatives of transient objects. The earliest observations include the recurrent nova V3890 Sgr (1990), whose exact identification was clarified by us

(Kato et al. 1990). The next advancement was with V838 Her (Nova Her 1991), whose eclipsing nature and exact orbital period was clarified by our observation (Kato, Hirata 1991). This is the first classical nova whose evolution of eclipses was caught from the early decline stage of the outburst (Leibowitz 1993; Szkody, Ingram 1994; Ingram et al. 1992). The "nova of the century" (Starrfield, Shore 1994) V1974 Cyg (Nova Cyg 1992) was followed with the advent of the e-mail alert list (see *vsnet-history* messages). This nova later turned out to be a permanent superhumper system (Semeniuk et al. 1994; Semeniuk et al. 1995; Retter et al. 1997; Skillman et al. 1997). The VSNET Collaboration later contributed to the international observing campaigns of the superhumps (A. Retter et al., in preparation).

The next major step was with V705 Cas (Nova Cas 1993), as introduced in subsection 1.4. After this nova, the VSNET has continuously provided public pages on individual novae,[67] which are referenced as a primary resource on recent novae. V723 Cas (Nova Cas 1995) has been one of the best studied novae in the VSNET history. The object was discovered by Minoru Yamamoto, whose report immediately triggered early follow-up observations (Hirosawa et al. 1995). Several early reports discussed against the classical nova-type classification (e.g. Munari et al. 1995a). We were the first, with the enormous amount of information collected by the VSNET, to predict that the object is a premaximum phase slow nova resembling HR Del (vsnet 223).[68] This prediction was later confirmed by a number of works (Ohsima et al. 1996; Munari et al. 1996; Iijima, Rosino 1996; Iijima, Rosino 1997; Iijima et al. 1998).

Although it was not a classical nova, V4334 Sgr (Sakurai's object) in 1996 brought a major breakthrough in "stellar evolution in real time" (Duerbeck et al. 1996; Duerbeck, Benetti 1996). This star is one of the best studied and discussed variable stars since the late 1990's (Duerbeck et al. 1997; Arkhipova, Noskova 1997; Asplund et al. 1997; Arkhipova et al. 1998a; Evans et al. 1998; Arkhipova et al. 1999b; Kipper, Klochkova 1999a; Kerber 1999; Kerber 2001; Asplund et al. 1999; Pavlenko et al. 2000; Duerbeck et al. 2000; Pavlenko, Duerbeck 2001; Shenavrin, Yudin 2001; Herwig 2001; Tatarnikov et al. 2001; Duerbeck 2002; Clayton 2002; Jeffery, Pollacco 2002; Kamath, Ashok 2002; Lawlor, MacDonald 2002; Lynch et al. 2002; Käufl et al. 2003). The VSNET not only relayed this breathtaking discovery of a final helium flash object, but also promptly provided prediscovery observations by Kesao Takamizawa (vsnet-alert 341)[69] (finally published by Takamizawa 1997), which has been frequently referenced and employed for theoretical modeling. See subsection 6.7 for its relation with R CrB-like stars (cf. Lawlor, MacDonald 2003).

In recent years, the VSNET Collaboration has been issuing scientific results, as well as early announcements and identifications, on individual novae: V2487 Oph (Nova Oph 1998, recurrent nova candidate: Nakano et al. 1998; Hachisu et al. 2002), V4444 Sgr (Nova Sgr 1999, the possibility of this being a recurrent nova has been discussed: Kato et al. 2004), V463 Sct (Nova Sct 2000, fast nova with an unusually prominent premaximum halt: Uemura et al. 2000g; Kato et al. 2002l), V445 Pup (Nova Pup 2000, unusual nova with no indication of hydrogen features: Kato et al. 2000b; Kato, Haseda 2000; Kato, Nakamura 2001; Nakano et al. 2001; M. Uemura et al. in preparation), V1548 Aql (Nova Aql 2001, slow nova initially reported as a more usual variable star: Uemura et al. 2001e; Kato, Takamizawa 2001), V1178 Sco (Nova Sco 2001, object originally confusedly reported to be a novalike object, later turned out to be a genuine nova with early stage oscillations: Haseda et al. 2001; Kanatsu et al. 2001; Kato, Fujii 2001), V2540 Oph (Nova Oph 2002, large-amplitude slow nova with strong post-outburst oscillations: Seki et al. 2002; Kato et al. 2002q). The VSNET Collaboration also joined multiwavelength campaigns on novae with satellites (e.g. V4743 Sgr: Wagner et al. 2003). The VSNET has recently been relaying discovery, independent and prediscovery detections of novae, as well as spectroscopic confirmation and early photometric observations: V2274 Cyg (Nova Cyg 2001: Sato et al. 2001), V4643 Sgr (Nova Sgr 2001: Kato et al. 2001d), V4740 Sgr (Nova Sgr 2001 No. 3: West et al. 2001); V4741 Sgr (Nova Sgr 2002: Liller et al. 2002a; Liller et al. 2002b), V4742 Sgr (Nova Sgr 2002 No. 2: Yamaoka et al. 2002d), V4743 Sgr (Nova Sgr 2002 No. 3: Haseda et al. 2002; West et al. 2002), V4744 Sgr (Nova Sgr 2002 No. 4: Yamaoka et al. 2002b), V4745 Sgr (Nova Sgr 2003: Kato et al. 2003g; Kato, Fujii 2003), Possible Nova (2002) in NGC 205 (Yamaoka et al. 2002a), Possible Nova (2003) in Scutum: (Nakano et al. 2003b).

The discovery of a very unusual eruptive object (V838 Mon, Brown et al. 2002; Wagner et al. 2002; Della Valle, Iijima 2002) was relayed through the VSNET alert system during its early stage of eruption, enabling early-stage observations (Bedient 2002a; Bedient 2002b). The identification of the object with GSC, IRAS and 2MASS objects was reported (Kato et al. 2002s). The early stage of this object was most unusual, showing an M-type spectrum at outburst maximum. On 2002 February 2, the object underwent a second major brightening, which was quickly relayed via the VSNET alert system worldwide, and the star became world-popular within a day. The object subsequently showed a prominent light echo when it faded (by the obscuration by the forming dust). There have been a number of works based on these observations (Kimeswenger et al. 2002; Munari et al. 2002; Goranskii et al. 2002; Sobotka et al. 2002; Bond et al. 2003; Crause et al. 2003; Wisniewski et al. 2003), which more or less employed VSNET observations and findings when discussing the peculiarity of this object. The origin of this eruption is still a mystery. Based on an idea proposed by Soker, Tylenda (2003), Kato (2003a) tried to explain the historical mysterious eruption of CK Vul.

---

[67] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Novae/novae.html⟩.

[68] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet/msg00223.html⟩.

[69] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-alert/msg00341.html⟩.

6.2.2. Recurrent Novae and Supersoft X-Ray Sources

The last two decades have dramatically changed our view of recurrent novae. The discovery of recurrent outbursts of V394 CrA in 1987 (Liller et al. 1987; Duerbeck 1988; Sekiguchi et al. 1989), V745 Sco in 1989 (Liller 1989; Duerbeck 1989; Sekiguchi et al. 1990b), V3890 Sgr in 1990 (Jones, Pearce 1990; Gonzalez-Riestra 1992; Harrison et al. 1993) and Nova LMC 1990 No. 2 (Sekiguchi et al. 1990a; Shore et al. 1991) resulted in a dramatic increase of our knowledge in recurrent novae (cf. Shara 1989; Anupama 1992; Sekiguchi 1995).

Another epoch of recurrent nova discoveries arrived like a flurry in the late 1990's and early 2000's. All of these recurrent nova outbursts were mediated via VSNET and followed in detail. The initial object in this series of recurrent nova outbursts was U Sco in 1999, whose early history was described in detail in subsection 4.1. This outburst of U Sco first enabled eclipse observations in real time during outburst [the eclipsing nature of U Sco was revealed only in 1990 (Schaefer 1990); the 1999 outburst was the first outburst since this discovery; see also Kato (1999l) for the retrospective detection of an eclipse during the 1987 outburst]. This observation led to the first detection of the period change in this system (Matsumoto et al. 2003b). This finding severely constrained the mass-transfer rate in quiescence (Hachisu et al. 2000a; Hachisu et al. 2000b), which makes U Sco the most promising candidate for an immediate precursor of a type-Ia supernova (Hachisu et al. 2000a).

The next object in this series was CI Aql. This object had long been suspected to be a dwarf nova based on the (apparently) small outburst amplitude of the 1917 outburst (Duerbeck 1987). Based on this potential identification, the object had been monitored by amateur variable star observers, notably by VSOLJ members and by members of the Recurrent Object Programme (see subsection 6.1). The proposed quiescent counterpart, however, was found to be an eclipsing binary which did not show CV characteristics (Mennickent, Honeycutt 1995). The same conclusion had been reached with snapshot spectroscopy (Szkody, Howell 1992), showing no indication of hydrogen emission lines. With this information, almost all observers stopped monitoring for an outburst, although there was the unexplained presence of a He II emission line (Greiner et al. 1996).

The news of a possible nova detection by Kesao Takamizawa on films taken on 2000 April 28 arrived at the VSNET on April 29. The reported position was extremely close to that of CI Aql. Minoru Yamamoto independently detected this phenomenon, and reported it to be a brightening of CI Aql. After careful examination of the identification, this possible nova is identified as a recurrent outburst of CI Aql, 83 years after the 1917 discovery (Takamizawa et al. 2000; Yamaoka et al. 2000).[70] The nova-type nature of the outburst was soon clarified with spectroscopy (Uemura, Kato 2000), confirming that CI Aql is a new recurrent nova. Being already known as an eclipsing binary, the evolution of the outburst light curve and the eclipse profile were precisely followed (Matsumoto et al. 2001; Matsumoto et al. 2003a). In particular, a dip-like sudden fading in 2000 November was noted (Matsumoto, Kato 2000). With these observational constraints, Hachisu, Kato (2001) succeeded in modeling the light curve, which was further refined and extended (Hachisu, Kato 2003b; Hachisu, Kato 2003c) to explain the unique high/low transitions in the supersoft X-ray source RX J0513.9−6951 in the LMC. Its galactic counterpart, V Sge, has been very recently identified (Hachisu, Kato 2003a). These objects are now considered to be promising candidates for precursors of type-Ia supernovae (e.g. Hachisu et al. 1999b; Hachisu, Kato 2003c; for earlier and other suggestions, see e.g. Hachisu et al. 1989; Hachisu et al. 1996; della Valle, Livio 1996; Hachisu et al. 1999a). The discoveries and modern detailed observations of U Sco and CI Aql outbursts thus provided firm observational evidence for recurrent novae and supersoft X-ray sources as immediate precursors of type-Ia supernovae. Other representative works (outside the VSNET) on CI Aql include photometry (Schmeja et al. 2000), spectroscopy (Kiss et al. 2001; Burlak, Esipov 2001), Chandra X-ray observation (Greiner, Di Stefano 2002), modeling (Lederle, Kimeswenger 2003), re-examination of the 1917 outburst (Williams 2000).

The next discovery of the series was on IM Nor (possible nova in 1920) in 2002 January. The outburst detection by William Liller (Liller 2002) was quickly relayed to the VSNET, and enabled early astrometric work to first firmly identify the quiescent counterpart and its recurrent nova nature (Duerbeck et al. 2002; Retter et al. 2002). The light curve of IM Nor was published more than 50 years after the 1920 outburst (Elliot, Liller 1972). There had been a suggestion of identification with the UHURU X-ray source 2U 1536−52, which was later confirmed to be spurious (Wyckoff, Wehinger 1979, see also Kato 2002g). Although the outburst light curve in 1920 resembled that of a slow recurrent nova T Pyx (Elliot, Liller 1972), the unusually faint quiescence inferred from Wyckoff, Wehinger (1979) had been a mystery. The exact identification with a new outburst solved this mystery, by the detection of a considerable variation in quiescence (Kato et al. 2002t). This suggestion was later confirmed by the detection of a short period variation with eclipse-like fadings (Woudt, Warner 2003a). Kato et al. (2002t) suggested, from the light curve and spectroscopic appearance, that both CI Aql and IM Nor are members of a new class of recurrent novae having intermediate properties between classical novae and fast recurrent novae.

6.3. Other Cataclysmic Variables

6.3.1. Intermediate Polars

Intermediate polars (IPs), which are a class of magnetic CVs (MCVs) having a magnetic white dwarf asynchronously rotating with the orbital motion (sometimes referred to as DQ Her stars: for recent reviews, see e.g. King, Lasota 1990; Patterson 1994; Hellier 1996;

---

[70] The object was later found to be positively recorded one day earlier (Liller 2000). An outburst in 1941–1942 was also later found in the plate archive (Schaefer 2001).

Buckley 2000; chapters 8 and 9 in Hellier 2001a). Although many "classical" IPs are novalike systems with thermally stable accretion disks, there are a number of IPs showing transient outbursts. Some of them look like dwarf novae (subsection 6.1), including well-known objects such as GK Per (Watson et al. 1985; Bianchini et al. 1986; Hutchings, Cote 1986; Norton et al. 1988; Patterson 1991), DO Dra (sometimes called YY Dra, see Kholopov et al. 1985b, Kholopov, Samus 1988 for the official nomenclature, Patterson et al. 1992; Haswell et al. 1997), EX Hya (Warner, McGraw 1981; Jablonski, Busko 1985; Heise et al. 1987; Hellier et al. 1987). There has been a long-standing discussion whether these IP outbursts originate from disk-instabilities or from mass-transfer bursts (Hellier et al. 1989; Hellier, Buckley 1993; Hellier et al. 2000). The modern understanding is that at least some of them are better understood as mass-transfer events (TV Col: Hellier, Buckley 1993), while others can be understood as disk-instability events (GK Per: Kim et al. 1992; Yi et al. 1992).

Rapid circulation of outburst alerts are extremely important for short-period systems, because these IP outbursts are usually very brief (usually less than 1 d) and require prompt follow-up observations (e.g. TV Col: Szkody, Mateo 1983; Szkody, Mateo 1984; Schwarz et al. 1988; Hellier, Buckley 1993; Retter et al. 2003, EX Hya: Bateson et al. 1986; Bateson, Jones 1987; Bond et al. 1987; Fujino et al. 1987; Hellier et al. 1989; Reinsch, Beuermann 1990; Jones 1994; Hellier et al. 2000, DO Dra: Wenzel 1983b; Wenzel 1983a; Hurst et al. 1985; McNaught et al. 1985; Hazen 1986; Dyck, Schmeer 1988; Kato, Dyck 1988). Although these outburst detections have historically been relayed via IAUCs, they were not usually rapid enough to enable early-stage observations of these outbursts. The VSNET Collaboration has played a role in detecting, and rapidly relaying these IP outbursts. Since IP outbursts tend to cluster (cf. Bateson et al. 1986, M. Uemura et al. in preparation), rapid electronic announcements of these outbursts have dramatically increased the chance of detailed follow-up observations, including simultaneous observations with satellites. The most remarkable recent example includes X-ray/optical simultaneous observations of two outbursts of DO Dra (Szkody et al. 2002b), whose optical coverage was based on observations by the VSNET Collaboration.

The IP outbursts for which the VSNET Collaboration played an important role of early notification include GK Per in 1996 (Ishida et al. 1996; Morales-Rueda et al. 1996; Nogami et al. 2002), in 1999, whose occurrence was notably predicted by a VSNET member, Tsutomu Watanabe (vsnet-future 2),[71] and real rise observed in detail by the VSNET Collaboration (vsnet-alert 2652).[72] These two outbursts were of special importance because detailed modern time-resolved CCD observations set a stringent limit on the expected occurrence of eclipses during outburst (Kim et al. 1992). These outbursts also provided an opportunity to study time-resolved spectroscopy of the QPOs (Morales-Rueda et al. 1996), as well as X-ray/optical detections of long-period QPOs (Morales-Rueda et al. 1999; Nogami et al. 2002). Multicolor observations of the 1996 outburst also provided observational constraints on the outburst models (Nogami et al. 2002). The 2002 outburst of GK Per provided an opportunity to study magnetic accretion (Bianchini et al. 2003). The long-term visual data, including the data reported to the VSNET, have been used for analysis of outburst properties (Simon 2002).

In most recent years, the VSNET Collaboration has succeeded in characterizing outburst properties of the unusual short-period intermediate polar HT Cam (Tovmassian et al. 1998). The outbursts are extremely brief (Watanabe 1998b), showing precipitous declines during the late part of outburst. From the time-difference of outburst maximum and the maximum appearance of IP pulses during the 2001 outburst, Ishioka et al. (2002a) concluded that the outburst was triggered by a dwarf nova-type disk instability phenomenon. The existence of a precipitous later decline can be explained by the truncation of the inner accretion disk. The 2001 outburst of HT Cam was also studied by another group (Kemp et al. 2002). The outburst of HT Cam thus has been one of milestones in the study of IP-dwarf nova relation.

Kato, Starkey (2002) reported short-period, nearly coherent, QPOs in V592 Cas, a nova-like star in the period gap. This object has been suggested to be a unique object in the period gap showing both properties of superhumps and occasional IP-like, nearly coherent, photometric oscillations. Several other dwarf novae have been suspected of IP-type signature, and were briefly discussed in the relevant parts of subsection 6.1.

6.3.2. *Polars*

Bright polars (MCVs with synchronously rotating white dwarfs) have been regularly monitored by the VSNET Collaboration. Many polars, notably AM Her (figure 9) and V834 Cen, occasionally show low states, whose occurrence has been notified through the VSNET alert system. Long-term CCD monitoring of faint polars have been regularly reported by Berto Monard.

Coordinated multi-wavelength observations have been conducted on several occasions. The targets include AM Her, AR UMa, ST LMi, VV Pup, AN UMa and EF Eri. Some of these collaborative studies has been already published as a solid paper (Hoard et al. 2002a).

6.3.3. *VY Scl-Type Stars*

VY Scl-type stars are novalike CVs with occasional low states, or fading episodes (cf. Warner 1995b; Robinson et al. 1981; see figure 10 for the VSNET light curve of MV Lyr). In some systems, these low states occur very infrequently (the best example being TT Ari: Bortle et al. 1980; Fuhrmann 1981; Shafter et al. 1982; Hudec

---

[71] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-future/msg00002.html⟩.

[72] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/alert2000/msg00652.html⟩.



Fig. 9. Light curve of AM Her from VSNET observations.



Fig. 10. Light curve of the VY Scl-type star MV Lyr from VSNET observations.

et al. 1984; Shafter et al. 1985; Hutchings, Cote 1985).[73] Such low states provide a unique opportunity to study the white dwarf atmosphere or to directly detect the secondary star (e.g. Jameson et al. 1982a; Jameson et al. 1982b). From the viewpoint of disk-instability model, the decreasing mass-transfer rate would produce a dwarf nova-type disk instability if there is no special mechanism to suppress the instability (Honeycutt et al. 1994; King, Cannizzo 1998). Observations, however, tend to show smooth monotonous declines (Honeycutt et al. 1994; Greiner 1998). This makes a clear contrast to the "low states" in dwarf novae (cf. 6.1.6). There must be a mechanism in VY Scl-type stars to somehow thermally stablizing the disk when the mass-transfer is reduced (cf. Leach et al. 1999). Dense observational coverage of VY Scl-type stars is therefore highly needed immediately after the start of their declines.

It is also known that VY Scl-type stars tend to show superhumps (cf. Patterson 1999). Since the mass-ratios (q) of VY Scl-type stars are not usually considered sufficiently small to enable excitation of the 3:1 resonance to produce superhumps (cf. Hirose, Osaki 1990; Wood et al. 2000; Whitehurst, King 1991; Molnar, Kobulnicky 1992; Murray 1998), there is apparently the need for an explanation of the cause of the superhumps. Murray et al. (2000) considered the variation in the mass-transfer rate in VY Scl-type stars, and presented a working hypothesis why VY Scl-type stars, with intermediate q, can show superhumps. This hypothesis also needs to be tested by more observations of VY Scl-type stars during different brightness states.

The VSNET Collaboration succeeded in early announcing a rare fading of V751 Cyg, whose VY Scl-type nature was suspected more than 20 years ago (Robinson et al. 1974) but had no comparable fading in recent years. The 1997 fading of V751 Cyg was originally reported to the VSNET by Laszlo Szentasko, and its progress was followed in detail by the VSNET Collaboration members, notably with CCD photometry at Ouda Station. This fading not only presented authentication of V751 Cyg as a genuine VY Scl-type star, but also enabled X-ray observations which led to the discovery of transient supersoft X-ray emission (Greiner et al. 1999; Greiner 2000). This observation, suggesting a possible extension of luminous supersoft X-ray sources (SSXS) toward a less-massive white dwarf, led to a revolutionary change in our view of VY Scl-type stars. Other rare low states of VY Scl-type stars announced through the VSNET Collaboration include LQ Peg (=PG 2133+115) in 1999 (the second historical fading: Watanabe 1999a; Kato, Uemura 1999a; Schmidtke et al. 2002) and BZ Cam in 1999 (the second historical fading: Watanabe 2000; Watanabe 2001; Kato, Uemura 2001b; Greiner et al. 2001). The 1999 fading of BZ Cam is notable in that transient superhumps were detected during the fading (Kato, Uemura 2001b), which may give support for the explanation by Murray et al. (2000).

---

[73] Although there was an interpretation of this object as a Z Cam-type dwarf nova permanently at standstill (Krautter et al. 1981), which was adopted by the GCVS team, it is generally considered as a VY Scl-type star with an occasional dramatic decrease of mass-transfer.

The VSNET Collaboration also succeeded in presenting the first-ever light curve of V504 Cen (Kato, Stubbings 2003), which has been suspected to be a VY Scl-type star from spectroscopy (Kilkenny, Lloyd Evans 1995), but had no solid photometric record qualifying the VY Scl-type nature.

Well-known VY Scl-type stars, such as MV Lyr, have long been best-observed targets by the VSNET Collaboration. Some of these long-term observations were employed to qualify the light curves of VY Scl-type stars (Greiner 1998). The well-known VY Scl-type star KR Aur was intensively studied by the VSNET Collaboration (Kato et al. 2002g), which led to the detection of short-term variations having power-law type temporal properties.

The other notable object is V425 Cas, whose low state in 1998 was announced by Timo Kinnunen through the VSNET, which led to our own discovery of short-term (2.65 d), large-amplitude (up to 1.5 mag) variations (Kato et al. 2001s). Such a type of variation had never been never in any class of hydrogen-rich CVs, and Kato et al. (2001s) suggested that they are dwarf nova-type instabilities in a moderately stabilized disk. This discovery was introduced as the "shortest period dwarf nova" in Astrophysics in 2002 (Trimble, Aschwanden 2003).

6.4. X-Ray Binaries and X-Ray Transients

X-ray binaries are close binary systems which consist of a compact object and a normal star. The mass accretion from the normal star onto the compact object generates strong X-ray emission. A number of X-ray binaries have been discovered as transients (cf. Liu et al. 2001). Their outbursts can be observed in all wavelengths, hence simultaneous multi-wavelength observations have played a key role to reveal the nature of X-ray binaries and X-ray transients (e.g. Orosz, Bailyn 1997; Hynes et al. 2000). Their outburst cycle is generally longer than a year, and in some cases, longer than decades. Prompt observations of an early outburst phase are hence important. The VSNET has been providing information about X-ray transients which enables prompt observations, not only for optical observers, but also for X-ray, UV, IR, and radio observers. Besides these recent studies, the earliest work by the authors include infrared quiescent observation of V404 Cyg = GS 2023+338 (Kato, Hirata 1990b), which first revealed the existence of the photometric period of 5.76 hr. Another outstanding early result was the discovery of superhumps and orbital variation in the outbursting X-ray transient GRO J0422+32 = V518 Per (Kato et al. 1992b; Kato et al. 1993a; Kato et al. 1995b). This observation has been one of the most comprehensive studies of superhumps in "classical" soft X-ray transients up to now (cf. O'Donoghue, Charles 1996).

Here we focus on the soft X-ray transients for which the VSNET collaboration conducted intense world-wide campaigns. Soft X-ray transients are also called X-ray novae, whose common characteristics were established in the mid-1990's (Tanaka, Lewin 1995; Chen et al. 1997): Their light curves are typically described with a Fast Rise and an Exponential Decay (FRED). The e-folding time is 30–40 d during the decay phase. In the FRED-type outburst, a reflare, or a secondary maximum is observed ~ 50 d after the outburst maximum. The outbursts are considered to occur due to a sudden increase of mass accretion rate in an accretion disk (Mineshige, Wheeler 1989; Huang, Wheeler 1989). Radial velocity studies of the secondary star have revealed that a dozen of soft X-ray transients contains stellar-mass black holes. Soft X-ray transients thus provide an ideal laboratory of the accretion physics onto the black hole, and they are called black hole X-ray transients.

In the framework of classical soft X-ray transients, the optical and X-ray emissions originate from the outer and the inner region of the accretion disk, respectively. Simultaneous optical and X-ray observations therefore enable us to study the evolution of the accretion disk and the mechanism of its activity. The VSNET collaboration enables us to obtain dense samples throughout outbursts. With these observations, we summarize the progress of studies of classical soft X-ray transients in subsection 6.4.3.

On the other hand, several recent transients have exhibited characteristics which are difficult to explain within the classical framework. The prompt observations by the VSNET collaboration has played an important role, in particular, for the research on the luminous fast transients (Uemura et al. 2002h) and rapid optical variations (Uemura et al. 2000c; Uemura et al. 2002d; Uemura et al. 2004a). We summarize our studies on these new classes of activity in subsections 6.4.1 and 6.4.2.

6.4.1. Fast X-ray transients, V4641 Sgr and CI Cam

It has been proposed that the outburst of classical soft X-ray transients is induced by thermal instability of the accretion disk (Mineshige, Wheeler 1989; Huang, Wheeler 1989). This model can explain the large e-folding time during their decay phase by considering a strong X-ray irradiation which stabilizes the outer disk. This standard picture cannot, however, explain two atypical outbursts, that is, the fast transients V4641 Sgr (Hjellming et al. 2000; Uemura et al. 2002h) and CI Cam (Belloni et al. 1999). Their outburst duration was only a few days, which is too short to be interpreted with the viscous decay of the classical soft X-ray transients.

On 1998 March 31, the All-Sky Monitor (ASM) of the Rossi X-Ray Timing Explorer (RXTE) detected a new X-ray transient named XTE J0421+560 (Smith et al. 1998). We were among the first to point out the presence of a supposed symbiotic star CI Cam (Bergner et al. 1995) within the error (vsnet-alert 1621).[74] This information was immediately relayed to X-ray and optical observers through the VSNET, and the proposed identification was securely confirmed with the discovery of an outbursting object at the location of CI Cam (Hjellming, Mioduszewski 1998). Contrary to classical X-ray transients, this object started a rapid fading with an *e*-folding time of ~ 0.5 d just after outburst maximum (Belloni et al. 1999). Its X-ray spectrum can be described by an absorbed power-law model with high-energy cutoff, which is atypical for X-ray transients (Belloni et al. 1999). Optical observations reported to the VSNET show that CI Cam brightened to 8.8 mag on April 3,[75] and faded by ~ 2 mag within two days. Spectroscopic observations revealed that CI Cam is a B[e]–X-ray binary (Belloni et al. 1999). The nature of the compact object is still unknown (Belloni et al. 1999; Orlandini et al. 2000; Hynes et al. 2002; Robinson et al. 2002). The VSNET Collaboration further obtained quiescent observations, which revealed the presence of weak activity (Kato, Uemura 2001d).

V4641 Sgr is a variable whose optical spectrum is that of an A-type star (Downes et al. 1995; note that the object was not correctly identified in the literature until we published a reliable chart, see subsection 2.2). In 1999 August, Tsutomu Watanabe, a member of the VSOLJ, noticed that the object entered an active state in the optical range (Watanabe 1999b). This state was characterized by the presence of large-amplitude variation having a possible periodicity of 2.5 d (Kato et al. 1999e). On September 15, the state was terminated by a short outburst reaching 8.8 mag independently detected by Rod Stubbings and the Kyoto team, following Berto Monard's detection of a brightening preceding this event (vsnet-alert 3475, 3477, 3478; Stubbings, Pearce 1999).[76] All these reports were immediately circulated through the VSNET alert network; the resultant vigil eventually enabled the historical detection of this perfectly unexpected giant outburst.

Following our notification, a corresponding X-ray outburst was found in the backlog recorded with the RXTE/ASM; the X-ray outburst was not discovered "real-time" even with X-ray all-sky monitor. The X-ray outburst, after reaching an astonishing flux of 12 Crab at the X-ray maximum, rapidly faded and returned to the pre-outburst level within only 2 hours (Smith et al. 1999). The duration of the optical outburst was also short (~ 5 d) (Uemura et al. 2002h). The VSNET collaboration team performed prompt observation of this short X-ray outburst, and revealed that the optical variation exhibited anti-correlation against the X-ray variation (Uemura et al. 2002h). Spectroscopic observations revealed that V4641 Sgr consists of a black hole and a late B-type star (Orosz et al. 2001). As reported below, V4641 Sgr experienced active phases in the next couple of years (Uemura et al. 2002d; Uemura et al. 2004a; Rupen et al. 2003), however, no comparable outburst to the 1999 September giant one has been yet observed.

While their binary components are totally different, the short outbursts of V4641 Sgr and CI Cam have several common characteristics: First, spatially resolved radio jets were associated with the outburst in both systems (Hjellming et al. 1998; Hjellming et al. 2000). Second, the peak luminosities reached the Eddington luminosity (Robinson et al. 2002; Revnivtsev et al. 2002b). Their high luminosity implies that supercritical accretion occurred. Since the matter falls with almost free-fall time scale in the supercritical accretion, the problem of the short duration of the outbursts can be reconciled. The optical–X-ray anti-correlation of V4641 Sgr may be understood with the scenario that the optically-thick supercritical accretion flow absorbed the X-ray emission and re-emitted the optical emission (Revnivtsev et al. 2002a; Uemura et al. 2002h). It is, however, still unknown how the supercritical accretion was induced.

6.4.2. *Short-term, non-thermal optical variations*

In the classical picture, the optical emission is thermal emission from the outer portion of the accretion disk where the temperature is relatively low ($\sim 10^4$ K). The observed time scale of optical variations is hence long, such as superhumps (e.g. O'Donoghue, Charles 1996) and orbital period variations (e.g. Kato et al. 1995b). The black hole binary system, GX 339-4 is, however, known to show rapid optical variations of the time scale of seconds (Motch et al. 1982; Motch et al. 1983). Such short time scale variations indicate that they originate from an inner portion of the accretion flow. They are proposed to be cyclo-synchrotron emission from the inner region (Motch et al. 1982), however, the mechanism to generate the emission and variations is poorly known. The VSNET collaboration team has recently observed optical rapid variations of two sources, that is, XTE J1118+480 = KV UMa (Uemura et al. 2000c) and V4641 Sgr (Uemura et al. 2002d; Uemura et al. 2004a).

XTE J1118+480 was discovered with the RXTE/ASM on 2000 March 29 (Remillard et al. 2000). We discovered an optical counterpart at 12.92 mag on March 30 (Uemura et al. 2000f). The optical light curve showed so many fluctuations that Uemura et al. (2000c) suggested the presence of optical rapid variations. Spruit, Kanbach 2002 detected optical short-term (a few tens of

---

[74] See ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/alert1000/msg00621.html⟩.

[75] See ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Xray/cicam.html⟩.

[76] See ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Xray/gmsgr.html⟩ for the full story.



Fig. 11. "Optical flash" from V4641 Sgr observed on 2002 July 7.

seconds) variations which correlate with the X-ray variations. These rapid optical variations are also proposed to be synchrotron emission from the inner accretion flow (Spruit, Kanbach 2002). Since XTE J1118+480 remained at a low/hard state throughout the outburst, the inner region is considered to be filled not by the standard disk, but by the advection dominated accretion flow (ADAF: Narayan, Yi 1994). In the ADAF region, the gas density is so low that the magnetic pressure is dominant (Mineshige et al. 1995). The optical rapid variations are probably generated at shock regions in such an inner region, which are formed by magnetic reconnections or collisions of blobs in the magnetically-dominated accretion flow (Uemura et al. 2002d).

After the luminous, short outburst in 1999 September, V4641 Sgr experienced a new active phase in 2002 and 2003. The VSNET collaboration succeeded in detecting rapid optical variations during the 2002 active state (figure 11), whose detailed features and interpretation are reported in Uemura et al. (2004a) in this volume.

6.4.3. Classical soft X-ray transients

Superhumps, which have been originally studied in SU UMa-type dwarf novae, are also observed in soft X-ray transients (e.g. O'Donoghue, Charles 1996). To observe the evolution of superhumps was difficult in the case of soft X-ray transients because of their rare outbursts, relatively long orbital period, and long durations of outbursts made a dense sampling throughout outbursts difficult. Our intense campaign of XTE J1118+480, however, first revealed the evolution of superhumps in soft X-ray transients (Uemura et al. 2002g). The superhump period was first 0.43% longer than the orbital period, and then decreased during the main outburst. Superhumps in SU UMa-type dwarf novae also exhibit this behavior, which can be interpreted by the contraction of the elliptical accretion disk (Osaki 1985) or the inward propagation of the eccentricity wave (Lubow 1992).

It is proposed that the reflare is induced by the growing tidal dissipation (Ichikawa et al. 1994; Truss et al. 2002). This model was developed based on the fact that superhumps appeared only after the reflare (O'Donoghue, Charles 1996). On the other hand, superhumps appeared even before the reflare in XTE J1118+480 (Uemura et al. 2002g). The VSNET collaboration obtained a dense sample around the reflare of XTE J1859+226, which is reported in Uemura et al. 2004c in this issue. We detected periodic variations, which may be superhumps, even before the reflare of XTE J1859+226. These observations are unfavorable for the above scenario for the reflare. It is also proposed that strong X-ray irradiation onto the outer accretion disk may induce the reflare (e.g. King, Ritter 1998).

6.4.4. High-Mass X-Ray Binaries

SS 433 = V1343 Aql is an active high-mass X-ray binary (HMXB) with relativistic jets, and the nature of the binary system is still mysterious in many aspects. Its optical magnitude is frequently monitored by VSNET observers, and remarkable (brightening or fading) behavior is reported via *vsnet-campaign-xray* when it occurs.

In 1995, 1998, and 2000, simultaneous multi-wavelength observations were organized by the ASCA team of RIKEN (N. Kawai et al., in preparation) in order to determine an accurate ephemeris of the eclipse and to compare light curves in the X-ray and optical wavelengths, which yield a clue to understand the emitting regions in the binary system. In these campaigns, VSNET played the role of a medium for exchanging information on the optical parts, such as calls for optical information, explanation of the background, and practical conditions and notes for observation (e.g. vsnet 103, for the 1995 campaign).[77]

6.5. *Supernovae*

Our contributions to the spectroscopic type determination of supernovae are summarized in table 1. Confirmatory observations, identifications and similar activities reported in IAUCs are also listed in table 2. We also run the mailing list *vsnet-campaign-sn*,[78] which is now widely known as the most reliable and up-to-date information of supernovae. The "Latest Supernovae" page (subsection 3.3) has been collaborating with it, which also provides USNO-A2.0 (Monet et al. 1998b) based charts made by Odd Trondal for almost every supernova brighter than 20th magnitude. The charts gives blue and red magnitudes for photometric reference stars, the latter ones have been widely used for "CR" measurements (see Appendix 2.1).

Besides objects described in the subsections 1.3 and 5.4, here we mention here remarkable examples.

SN 1997ef in UGC 4107 was discovered by Yasuo Sano, one of the most active participants of VSNET (Nakano et al. 1997). It was originally announced as "possible supernova", because its spectra was quite unusual and could not convincingly be classified as that of a supernova (Garnavich et al. 1997b). Further spectroscopic observations revealed that it was likely an explosion with massive (several $M_\odot$) ejecta. The object consequently received the designation of SN 1997ef (Hu et al. 1997, Filippenko, Martin 1997). Filippenko, Martin (1997) and Garnavich et al. (1997a) suggested that it was likely be an explosion of a stripped very massive star, a suggestion supported by theoretical modeling (Iwamoto et al. 2000). It is the first example of the so-called "type-Ic hypernova", a concept developed after the suggested association SN 1998bw with GRB 980425 (Galama et al. 1998, Iwamoto et al. 1998), SN 2002ap (see subsection 5.4), and the unambiguous identification of the supernova signature (SN 2003dh) in GRB 030329 (see subsection 6.15).

SN 1997ei was discovered by Masakatsu Aoki (Nakano, Aoki 1997) in NGC 3963. The first spectroscopy reported in IAUC indicated that it is a type-Ia supernova (Garnavich et al. 1997c). Our spectroscopy showed some peculiarity, then we reported that it could be a peculiar type-Ia supernova (Ayani, Yamaoka 1997). From later spectroscopic observations, it finally turned out to be a type-Ic supernova (Wang et al. 1998, Filippenko, Moran 1998).

The SN 1998T case taught us the importance of identification. It was produced in a pair of interacting galaxies, and some catalogues of galaxies gave discrepant designations for them. The blobby nature of the host galaxies also led to misidentifications of the supernova. The discussion on *vsnet-chat*, including the consultation of the NGC/IC project [79], led us to a correct identification of the galaxy, which was accompanied by precise astrometry (Yamaoka et al. 1998).

SN 1998bu in M 96 was discovered by Marko Villi. It was the nearest supernova since SN 1993J in M 81. The first spectroscopy reported in IAUC was a high-dispersion one, which could only determine the depth of the interstellar absorption within our Galaxy and the host galaxy (Munari et al. 1998). Our spectrum revealed that it is a type-Ia supernova, and this report was naturally posted to vsnet-alert 1785[80]. It was distributed earlier than the relevant IAUC (Ayani et al. 1998), which also included other spectroscopy (Meikle et al. 1998). In response to the type determination, the COMPTEL instrument was pointed towards SN 1998bu in order to detect the line γ-ray of ^{56}Co decay, which had been detected only from SN 1991T, the peculiar luminous type-Ia supernova. Despite SN 1998bu is as close to us as SN 1991T, the γ-ray lines was not detected (Georgii et al. 2002), which may suggest a diversity among type-Ia supernovae in the line γ-ray, as well as in the light curves (Riess et al. 1996) and in the spectra (Nugent et al. 1995).

The first spectroscopic observation of SN 1998es indicated that it is an intrinsically bright type-Ia supernova like SN 1991T (Jha et al. 1998a). Our report confirmed it, giving in addition the spectral evolution and the interstellar extinction (Ayani, Yamaoka 1998c).

SN 1999dn was a case similar to SN 1997ei. Our spectrum suggested that it was of type Ic with week He I lines (Ayani et al. 1999). The same IAUC also contained two spectroscopic observations, one of which (Turatto et al. 1999) reached the same conclusion, but the other suggested it was of type Ia (Qiu et al. 1999). Later spectroscopy revealed that SN 1999dn is an intermediate event between type Ib and type Ic (Pastorello et al. 1999).

SN 2000ch was a very subluminous supernova. It was originally announced as a variable star in field of NGC 3432 (Papenkova, Li 2000). We noticed that the object can be seen on the DSS images since 1998 (vsnet-chat 2908),[81] (Yamaoka 2000). On the other hand, spectroscopy of this object (Wagner et al. 2000) suggested that it is located within NGC 3432. From these findings, Filippenko (2000) assigned the supernova designation of this object, resembling the "type-V" SN 1961V. The comment of the discoverer (vsnet-chat 2944)[82] finally supported this classification.

SN 2001bf was yet another example similar to SN 1997ei and SN 1999dn. Despite of its low signal-to-noise ratio, our spectrum clearly showed a deep Si II absorption feature, from which we estimated that it is a type-Ia supernova (Kawakita et al. 2001a). Another group suggested that it is of type Ic (Phillips, Krisciunas 2001), but the later spectral evolution revealed that it is indeed a type-Ia supernova (Chornock et al. 2001).

---

[77] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet/msg00103.html⟩.

[78] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-campaign-sn/msg00785.html⟩.

[79] ⟨http://www.ngcic.com/⟩.

[80] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/alert1000/msg00785.html⟩.

[81] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/chat2000/msg00908.html⟩.

[82] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/chat2000/msg00944.html⟩.

SN 2002ao was a slightly different case. The first report on IAUC quoted the resemblance with type-IIb supernovae (Gal-Yam et al. 2002a, Gal-Yam et al. 2002b). We estimated that it is of type Ic (Kinugasa et al. 2002d), which is consistent with a later report (Filippenko, Chornock 2002) indicating the resemblance with SN Ic 1999cq. The rapid decline (Yamaoka et al. 2002c) of SN 2002ao was another common feature with SN 1999cq (Matheson et al. 2000).

Other IAUC issues in relation with the activity of the VSNET administrators include IAUC 7033 (SN 1998eg: Hurst et al. 1998), IAUC 8101 (SN 2003cg: Hirose, Yamaoka 2003), and IAUC 8171 (SN 2003gs: Evans et al. 2003).

6.6. *Symbiotic Variables*

Bright symbiotic variables (cf. Allen 1984; Kenyon 1986; Belczynski et al. 2000; Munari, Zwitter 2002) have been well-observed by many members of the VSNET Collaboration (see figure 12). These observations provided a number of detections of outbursts and eclipses, which were immediately relayed to more specialized researchers for detailed study.

The eclipse phenomenon in the outbursting object FN Sgr was discovered through the VSNET regular activity.[83] This work was summarized by Watanabe et al. (2000).

The outburst and possible eclipse phenomenon in V343 Ser = AS 289 was discovered by Kesao Takamizawa (= Tmz V17, vsnet-obs 8957)[84] and Minoru Wakuda. The final publication is by Takamizawa et al. (1998).

V1413 Aql = AS 338 is another object whose eclipsing symbiotic nature (cf. figure 13) was revealed by amateur astronomers (Wakuda 1988).[85] The outbursts and eclipses were regularly announced in VSNET, which have been followed by a number of researchers (e.g. Esipov et al. 2000; Kolotilov et al. 2001).

Kato (2000) reported the similarity of light curves between CH Cyg and the supersoft X-ray sources (V Sge and RX J0513.9−6951) based on VSNET observations.

The 1997 and 2000 outbursts of Z And were reported in vsnet-alert 938,[86] and vsnet-alert 5232, 5233,[87] respectively. These outbursts enabled, more or less owing to the VSNET alerts, modern observations of this classical symbiotic binary (Sokoloski, Bildsten 1999; Skopal et al. 2000; Skopal 2003; Tomov et al. 2003).

Ishioka et al. (2001d) systematically studied short-term variations of V694 Mon = MWC 560. During the entire period of observations, the object showed pronounced flickering activity. This work has been referred to as one of the most intensive photometric observations of ths peculiar symbiotic variable (Schmid et al. 2001; see also Sokoloski et al. 2001 for a recent survey work). We now have a dedicated mailing list for symbiotic stars *vsnet-symbio*[88] for informing about recent activities, particularly announcements of outbursts and eclipses, of symbiotic variables.

6.7. *R Coronae Borealis Stars and Related Objects*

R CrB stars are hydrogen-deficient carbon stars which show occasional fadings caused by dust formation (Clayton 1996). A representative light curve of R CrB from VSNET observations is shown in figure 14. Early announcements of the fadings of R CrB stars can provide the best opportunities to study the formation mechanism of dust in these stars. Before the VSNET era, these fadings had been only widely announced only when the objects had unmistakably faded (typically below ≥ 1 mag below their usual maximum); the early decline stage had been usually overlooked. The real-time communication via the VSNET public lists broke this historical limitation. The most dramatic instance was with the first-ever fading of FG Sge (also known as a final helium flash object) in 1992 late August–September (cf. history partly recorded in vsnet-history 200, 202). It was only when the object had already faded by 1 mag when IAUCs were issued (Papousek, Granslo 1992; Waagen et al. 1992; Woodward et al. 1992). This detection of a possible fading (1992 August 30) reported by Nobuhiro Makiguchi (VSOLJ, see subsection 2.3) was immediately attended by a number of world-wide observers. This fading was the first one of the series of fadings successively occurring up tp now (cf. Iijima, Strafella 1993; Iijima 1996; Iben, Livio 1993; Jurcsik 1993; Stone et al. 1993; Woodward et al. 1993; Arkhipova 1994; Simon 1994; van Genderen 1994; Kipper et al. 1995; van Genderen, Gautschy 1995; Arkhipova 1996; Arkhipova, Esipov 1996; Arkhipova et al. 1994; Arkhipova et al. 1996; Arkhipova et al. 1998b; Arkhipova et al. 1998c; Arkhipova et al. 1999a; Kipper, Klochkova 1999b; Gonzalez et al. 1998; Jurcsik, Montesinos 1999; Kipper, Klochkova 2001). The discovery of the fading in FG Sge brought a breakthrough in the understanding of late-time low-mass stellar evolution: our understanding of FG Sge had been slow and limited before this phenomenon (cf. Cowley et al. 1985; Aikawa 1985; Arkhipova et al. 1991; although the unusual nature of this object was recognized more than 30 yr ago and had long been discussed from different standpoints, including binary hypothesis and a thermal pulse in stellar evolution (e.g. Kukarkin 1960; Furtig, Wenzel 1962; Herbig, Boyarchuk 1968; Wenzel, Fürtig 1971; Paczynski 1970; Paczynski 1971; Sparks, Kutter 1973; Christy-Sackmann, Despain 1974; Langer et al. 1974; Cohen et al. 1980; Kurtz 1986; Feibelman, Bruhweiler 1990). It is now widely believed that FG Sge, V605 Aql, and V4334 Sgr (cf. subsection 6.2.1), as

---

[83] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Symbio/fnsgr.html⟩.

[84] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/obs8000/msg00957.html⟩.

[85] See ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/docs/v1413aql.html⟩ for a full story. See also Schulte-Ladbeck (1985), Munari (1992), Munari et al. (1993).

[86] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-alert/msg00938.html⟩.

[87] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/alert5000/msg00232.html⟩ and ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/alert5000/msg00233.html⟩.

[88] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-symbio/maillist.html⟩.

Table 1. Spectroscopy of supernovae.

SN name	Type	VSNET article*	Relevant IAUC	Remark
1995al	Ia	alert 266	6256 (Ayani, Kawakita 1995)	see sec. 5.4.
1997ei	Ic	alert 1399	6800 (Ayani, Yamaoka 1997)	see text.
1998an	Ia	alert 1684	6878 (Ayani, Yamaoka 1998a)	
1998aq	Ia	alert 1681	6878 (Ayani, Yamaoka 1998b)	
1998bu	Ia	alert 1785	6905 (Ayani et al. 1998)	see text.
1998es	Ia,pec	alert 2394	7059 (Ayani, Yamaoka 1998c)	see text.
1999bg	II	alert 2816	7137 (Ayani, Yamaoka 1999a)	
1999dn	Ib/c	alert 3380	7244 (Ayani et al. 1999)	see text.
1999gn	II	alert 3842	7336 (Ayani, Yamaoka 1999b)	
1999gq	II	alert 3862	7339 (Ayani, Yamaoka 1999c)	
2001bf	Ia	alert 5873	7625 (Kawakita et al. 2001a)	see text.
2001bg	Ia	alert 5871	7622 (Kawakita et al. 2001b)	
2001dp	Ia	alert 6299	7683 (Migliardi et al. 2001)	
2002ao	Ic	camp-sn 339	7810 (Gal-Yam et al. 2002a)	see text.
2002ap	Ic,pec	alert 7120	7811 (Kinugasa et al. 2002a)	see sec. 5.4.
2002bj	II(n?)	camp-sn 363	7844 (Kinugasa et al. 2002e)	luminous
2002bo	Ia	alert 7241	7848 (Kinugasa et al. 2002c)	
2002bu	IIn	alert 7259	7864 (Ayani et al. 2002)	
2002fk	Ia	alert 7516	7976 (Ayani, Yamaoka 2002)	
2003J	II	camp-sn 534	8048 (Ayani et al. 2003)	
2003K	Ia	camp-sn 534	8048 (Ayani et al. 2003)	

*alert for *vsnet-alert*, chat for *vsnet-chat*,
camp-sn2002ap for *vsnet-campaign-sn2002ap*,
camp-sn for *vsnet-campaign-sn*,

Table 2. Identification, astrometry and follow-up of supernovae.

SN name	VSNET articles*	Relevant IAUCs	Remark
1998T	chat 775, 779, 780, 786, 787	6859 (Yamaoka et al. 1998)	see text.
1999et	···	7344 (Yamaoka, Yamamoto 2000)	
2000M	alert 4320	7373 (Dimai et al. 2000)	
2000P	alert 4363, 4365, 4366, 4368, 4369	7378 (Colas et al. 2000), 7379 (corrigendum)	
2000ch	chat 2908, 2941, 2944	7415 (Papenkova, Li 2000), 7417 (Wagner et al. 2000), 7419 (Yamaoka 2000), 7421 (Filippenko 2000)	see text.
2000cm	alert 4944, chat 3035	7436 (Mueller, Rykoski 2000), 7437 (Jha et al. 2000a), 7438 (Jha et al. 2000b)	
2001dp	alert 6317	7683 (Migliardi et al. 2001)	
2002ao	alert 7190	7836 (Yamaoka et al. 2002c)	see text.
2002ap	camp-sn2002ap 154	7836 (Yamaoka, Rogers 2002)	prediscovery
2002dm	camp-sn 440	7921 (Sanders et al. 2002), 7923 (Yamaoka, Sanders 2002)	
2002ed	alert 7441, camp-sn 454	7940 (Monard, Li 2002), 7943 (Yamaoka 2002)	
2003ez	···	8141 (Papenkova, Li 2003), 8142 (Yamaoka, Ayani 2003)	

*Abbreviations same as in table 1.



Fig. 12. Light curve of CH Cyg from VSNET observations.



Fig. 13. Light curve of V1413 Aql from VSNET observations. Deep eclipses are superimposed on a gradual fading from the 1995 outburst (Elizalde et al. 1995; Munari et al. 1995b.



Fig. 14. Light curve of R CrB from VSNET observations.

well as some unusual R CrB stars (cf. V348 Sgr, e.g. Clayton 2002) comprise a sequence of final helium-flash objects (cf. Lawlor, MacDonald 2003).

The dramatic fading of ES Aql, which had been suspected to be an R CrB-type star, was first announced through the VSNET alert network; Peter Williams was the first to detect this object getting fainter than 14.0 on 2001 March 22. Based on this information, Clayton et al. (2002) succeeded in clarifying the R CrB-type nature of this object.

The other R CrB-type star recognized through the VSNET activity is V2552 Oph = Had V98 (Kato, Haseda 2003, Rao, Lambert 2003; see subsection 5.3).

We now have a dedicated mailing list for R CrB stars and related objects *vsnet-rcb*,[89] which is best employed by the researchers of this field. More recent announcements of rare R CrB-type fadings include the 2003 fading of V3795 Sgr (vsnet-rcb 585).[90] Although the attempt was not very successful, Rosenbush (2001) tried to predict future light curve of R CrB using VSNET observations.

6.8. Be Stars

Among recent phenomena in Be stars (B-type emission-line stars), the case of δ Scorpii is still fresh in our memory. The star, which had been considered as a non-variable B-type star, underwent a dramatic change in 2000 July. Sebastian Otero, a VSNET member, visually noticed a 0.1 mag brightening in δ Scorpii and issued an alert through the VSNET (vsnet-be 2).[91] The supposed transition to a Be star was subsequently confirmed by spectroscopic observations (Fabregat et al. 2000). The star further brightened to a maximum of $V \sim 1.8$ around 2000 July 30. Such a dramatic change in a bright naked-eye star is extremely rare. The only comparable precedent phenomenon was γ Cassiopeiae in 1937, which brightened to $V \sim 1.6$. This news was widely distributed through public news media, as originating from the VSNET, and became the one of the most popular astronomical phenomena in that year. This object, after reaching a temporal minimum just following the initial peak, continues to show remarkable activities up to 2003 (Otero et al. 2001; Gandet et al. 2002). On several occasions in 2002 and 2003, the star even brightened close to $V \sim 1.5$, even slightly surpassing in brightness the historical event of γ Cassiopeiae.

In the aftermath of this event, visual monitoring of bright Be stars has been conducted by a number of VSNET members, most intensively by Otero. These observations have detected of number of outbursts, e.g. in μ Centauri (cf. Baade et al. 2001) and ω Canis Majoris (cf. Otero 2002). The VSNET runs dedicated lists on the Be-star phenomenon, *vsnet-be*[92] and *vsnet-campaign-be*.[93]

6.9. Wolf–Rayet Stars

Wolf–Rayet (WR) stars are massive, luminous, and hot stars which have lost their hydrogen envelope, and are considered to be immediate precursors of some kinds of supernovae, and likely GRBs. In spite of their astrophysical importance, WR stars were less conspicuous objects in terms of optical variability. The only known categories of variability in WR stars had been occasional short-period obscuration by dust production, instabilities in the wind, or possible pulsation (see Williams et al. 1990; Williams 1995; Veen et al. 1998 for recent references).

This situation dramatically changed by the discoveries of the two most actively variable WR stars by the VSNET team (WR 104 = Had V82 = V5097 Sgr: Kato et al. 2002d, WR 106 = Had V84 = V5101 Sgr: Kato et al. 2002c). Both objects were initially reported as variable stars by Katsumi Haseda, detected during his search for novae. The variables soon turned out to be identical with known WR stars. In particular, WR 104 is a well-known binary consisting of a late-type WR star and an OB star. The most remarkable feature in this object is the presence of a “dusty pinwheel nebula” (Tuthill et al. 1999; Tuthill et al. 2002) co-rotating with the interferometric binary. The importance of the detection of large-amplitude optical variation was immediately recognized and communicated via VSNET alert network. Such a large variation in WR 104 required a non-classical interpretation (Kato et al. 2002d). With these conspicuous discoveries, the VSNET significantly broadened the scope of variability studies of WR stars, and this field is now becoming one of the contemporary topics in studying the WR-type activity. Combined with the recent advance in GRB astronomy, the importance of fundamental understanding of various phenomena in WR stars will be a matter of key importance.

6.10. Pre-main Sequence Stars

Some pre-main sequence stars show dramatic outbursts (FU Ori stars: FUORs) and smaller outbursts (EX Lup stars: EXORs), probably originating from some kind of instabilities in the circumstellar disk (e.g. Kawazoe, Mineshige 1993). Outbursts of these objects were occasionally followed by the VSNET Collaboration. The best example was the 1995 outburst of V1143 Ori (Baba et al. in preparation). Outbursts of V1118 Ori have also been occasionally reported (e.g. Hayakawa et al. 1998).

Among pre-main sequence stars, Herbig Ae/Be stars have been one of the best observed objects by VSNET members. The recent discovery of a likely Herbig Ae/Be star with large-amplitude variation, MisV1147, has been widely studied through the VSNET (M. Uemura et al. in preparation). Other objects of this class, well observed and timely informed through the VSNET, include UX Ori, RR Tau, CQ Tau, AB Aur, RZ Psc and WW Vul (see figure 15). Remarkable variations in these objects have been relayed through a dedicated list of pre-main sequence

---

[89] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-rcb/maillist.html⟩.

[90] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-rcb/msg00585.html⟩.

[91] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-be/msg00002.html⟩.

[92] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-be/maillist.html⟩.

[93] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-campaign-be/maillist.html⟩.



Fig. 15. Complex light curve of the Herbig Ae/Be star RR Tau from VSNET observations.

variables, *vsnet-orion*,[94] and utilized for detailed follow-up studies.

6.11. *Peculiar Variable Stars*

Some peculiar (or unique) variable stars were also studied by the VSNET Collaboration. The most striking objects include V651 Mon (the central star of the planetary nebula NGC 2346). This binary system containing a B-type subdwarf underwent a totally unexpected fading in 1981–1985 (for a summary of this event, see Costero et al. 1986). Although no similar phenomenon was recorded in the century-long past photographic records, the object underwent another unexpected fading episode in 1996–1997. This phenomenon was detected by Danie Overbeek, and immediately reported to the VSNET (vsnet-alert 548).[95] Following this announcement, Kato et al. (2001e) succeeded in recording the phenomenon in detail. There was a sharply defined transient clearing (brightening) even during this fading, which was ascribed to a sharply defined, small (several times $\sim 10^{11}$ cm) lucent structure within the obscuring body. See figure 16 for the recent light curve.

6.12. *Pulsating Variables*

Although pulsating variables were only occasionally selected as intensive targets of the VSNET Collaboration, the VSNET public data archive (cf. figure 17), as well as the VSOLJ database, have been frequently used in period analysis, and correlation studies with other observational modalities. These references to the VSNET/VSOLJ data have been so numerous that we only list most recent ones: historical archive for μ Cephei (Skiff 2001), evolution of R Hya (Zijlstra, Bedding 2002), dust formation in L$_2$ Pup (Bedding et al. 2002), possible chaotic behavior in R Cyg (Kiss, Szatmary 2002), period determination of V648 Oph (Kazarovets, Pastukhova 2002), non-variability of EK And (Yoshida et al. 2003), and period variation in T UMi (Szatmary et al. 2003).

Numerous new variable stars reported in *vsnet-newvar* (subsection 5.2) have been studied in detail, and have been reported in a numerous number of papers published in Information Bulletin on Variable Stars (IBVS).

ρ Cas, originally classified as a yellow semiregular variable, is now considered to be a low-temperature counterpart (yellow hypergiants: de Jager 1998; Israelian et al. 1999; de Jager et al. 2001; Stothers, Chin 2001) of the extremely luminous hot hypergiants (luminous blue variables: LBVs). This object occasionally undergoes temporary optical fadings caused by huge mass-loss events, usually once in decade(s) (figure 18). The most spectacular recent event occurred in 2000 (Lobel et al. 2003). Independent detections, including that by one of the authors (TK), of this phenomenon were circulated through the VSNET (vsnet-alert 5186, 5187),[96] which enabled first dense optical coverage of this kind of phenomenon. Upon recognition of the astrophysical significance of this event, we set up a dedicated mailing list *vsnet-campaign-rhocas* in 2000 August.[97] These observations have provided the primary resource on this rare event, and are referenced on Lobel's dedicated page on ρ Cassiopeiae.[98] In 2003 there was a small signature of a line variation similar to the precursor event in 2000. This news was widely announced through *vsnet-campaign-rhocas* and the object has been intensively observed.

6.13. *Eclipsing Binaries*

Although classical (other than CV-type or symbiotic type) binaries are usually not intensive targets for the VSNET Collaboration, there have been several calls for observations, particularly for long-period

---

[94] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-orion/maillist.html⟩.

[95] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-alert/msg00548.html⟩.

[96] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/alert5000/msg00186.html⟩ and ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/alert5000/msg00187.html⟩.

[97] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-campaign-rhocas/maillist.html⟩.

[98] ⟨http://cfa-www.harvard.edu/~alobel/⟩.



Fig. 16. Light curve of V651 Mon from VSNET observations.



Fig. 17. Light curve of the RV Tau-type star R Sct from VSNET observations.



Fig. 18. Light curve of ρ Cas from VSNET observations.

eclipsing binaries (e.g. OW Gem, EE Cep). Visual and CCD/photoelectric observations, as well as comparison star sequence information, have been exchanged on VSNET lists. The discovery of eclipses in the bright naked-eye star δ Velorum was one of the breaking news mediated through the VSNET (see also Otero et al. 2000). A dedicated list for δ Velorum observations has been set up, *vsnet-campaign-deltavel*.[99]

The VSNET runs dedicated lists on eclipsing binaries, *vsnet-ecl*[100] which are now recognized as one of world-wide networks exchanging information in eclipsing binaries. Kazuo Nagai (VSOLJ) has been summarizing the times of minimum of eclipsing binaries reported to *vsnet-ecl*.

6.14. Blazars

Blazars (BL Lac objects and optically violently variable quasars) are also one of the targets of long-term and intensive observing campaign. We run a dedicated list *vsnet-campaign-blazar*[101] for exchanging information on blazar activities and upcoming collaboration with other world-wide blazar groups. From the very establishment of the Whole Earth Blazar Telescope (WEBT: Villata et al. 2002b),[102] we have been continuously in collaboration with this group.

In addition to continuous visual monitoring campaigns on strongly active blazars, such as OJ 287, Markarian 421, 3C 279, quick electronic circulation and prompt feedback via the VSNET discussion group led to the notable discovery of unexpectedly large, short-term intranight variation of BL Lac in 1997.[103] This intensive observation was initiated by a report of a bright state of BL Lac in IAUC (Mattox et al. 1999), which initially revealed "outbursts" every five to ten days. Following this stage, the object entered a more active phase in mid-July to early August. During this stage, real-time comparisons of visual observations through the VSNET discussion group revealed substantial discrepancies depending on observers' longitudes. These discrepancies, which were much larger than what had been recognized as blazar microvariability, were initially considered as a result of an inhomogeneous magnitude system. This possibility was soon disproved by the real-time distribution of the modern photoelectric comparison star magnitudes. By the end of 1997 July, the incredible short-term variation (0.8 mag in four hours) in BL Lac had become a doubtless phenomenon.

In response to these visual observations, the Kyoto University team obtained long time-resolved CCD photometry at Ouda Station, revealing the unprecedented complexity and fast variation in the light curve. A similar conclusion was reached with time-resolved CCD photometry by Tonny Vanmunster, which was also rapidly communicated via VSNET. The variation has a power-law temporal properties, analogous to those of AGN variability (although the variation in BL Lac was much more violent and rapid), which can be tracked down to five minutes. This is one of the shortest time scales ever recorded in blazar optical variation (Matsumoto et al. 1999). The early results by the VSNET team were presented at the 23rd IAU General Assembly held in Kyoto held in 1997 August.

Since then, several studies have been performed in collaboration with the WEBT (Ravasio et al. 2002; Villata et al. 2002a). Long-term light curves of selected blazars from VSNET observations have been widely used for correlation with multiwavelength data.

6.15. Gamma-Ray Bursts

The VSNET collaboration team has tried to search optical afterglows of GRBs and reported their detection or upper-limits of magnitudes. Earlier announcements of Beppo Satellite per Astronomia X (BeppoSAX) GRB detections were reported to *vsnet-alert*. Since 2000 June, we set up a dedicated list for GRB announcements (*vsnet-grb*).[104] This mailing list is currently employed to distribute satellite-borne alerts, as a secondary distribution node of the GRB Coordinates Network (GCN).[105] A public mailing list *vsnet-grb-info*,[106] originally extended from *vsnet-grb*, currently provides GCN Circular information as well as our own observations by the VSNET Collaboration. Table 3 summarizes our own GRB afterglow observations (not all observations are listed), which is an extension of table 1 in Yamaoka et al. (2003b). See the corresponding GCN Circulars for more details.

Here we summarize our significant detections and their importance in GRB astronomy:

GRB010222: Our observation of the afterglow of GRB010222 covered a period around the jet break (Uemura et al. 2001a). This is the first detection of the GRB afterglow in Japan.[107] This observation encouraged observers having small telescopes in Japan to try observations of GRB afterglows.

GRB020331: We succeeded in observing an early afterglow with $R_\mathrm{c} = 17.9$ (17.5–18.7; 1-σ limits) 64 min after the burst (Kato et al. 2002r). This is the earliest observation for this afterglow. Our observation revealed that the light curve can be described with a single power-law from the early phase (~ 1 hr) of the afterglow.

GRB021004: Owing to the prompt identification by the HETE-2 satellite (Shirasaki 2002), we first revealed the continuous behavior of the GRB afterglow around 1 hr af-

---

[99] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-campaign-deltavel/maillist.html⟩.

[100] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-ecl/maillist.html⟩.

[101] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-campaign-blazar/maillist.html⟩.

[102] ⟨http://www.to.astro.it/blazars/webt/⟩.

[103] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/BLLAC/bllac.html⟩.

[104] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-grb/maillist.html⟩.

[105] ⟨http://gcn.gsfc.nasa.gov/⟩.

[106] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-grb-info/maillist.html⟩.

[107] The first detection with a Japanese telescope was the infrared afterglow of GRB000926 observed with the Subaru telescope in Hawaii (Kobayashi et al. 2000).

Table 3. GRB Afterglows observed by the Kyoto team.

GRB name	Alert type	Epoch (d)*	Telescope (cm)	Magnitude†	GCN Circular
001025A	IPN	1.7	60	>18.0	866 (Uemura et al. 2000a)
001025B	IPN	1.6	60	>19.0	unpublished
001212	IPN	2.0	25	>17.5	902 (Ishioka et al. 2000b)
010214	SAX	0.29	30	>18.0	948 (Uemura et al. 2001d)
010220	SAX	1.5	30	>17.5	unpublished
010222	SAX	0.23	30	17.8	984 (Uemura et al. 2001a)
		0.55	30	19.3	
011030	SAX	0.28	25	>16.0	unpublished
011130	HETE	0.47	25	OF	unpublished
020124	HETE	0.079	25,30	OF	unpublished
020331	HETE	0.033	25,30	17.9	1363 (Kato et al. 2002r)
020812	HETE	0.018	60	>21.2	1515 (Ohashi et al. 2002)
		0.038	25,30	>19.1	1521 (Uemura et al. 2002a)
020813	HETE	0.36	60	∼19	unpublished
020819	HETE	0.14	60	>15.0	unpublished
020903	HETE	0.18	25	OF or >17.5	1537 (Uemura et al. 2002b)
021004	HETE	0.036	25,30	16.3	1566 (Uemura et al. 2002c)
030227	HETE	0.046	25	>18.0	1899 (Uemura et al. 2003a)
030329	HETE	0.053	25,30	12.6	1989 (Uemura 2003), 1994 (Uemura et al. 2003f), 2147 (Yamaoka et al. 2003a)
030528	HETE	0.004	25	>16.0	2252 (Uemura et al. 2003b)
030823	HETE	1.24	60	>20.5	2370 (Imada et al. 2003)

*Day after the burst detection.

†Unfiltered CCD or R_c magnitude. OF = optical counterpart out of the field of view.

ter the burst (Uemura et al. 2002c; Uemura et al. 2003d). An early afterglow was also observed in GRB990123 (Sari, Piran 1999a), however, our observation covered a period corresponding to an observing gap of the early afterglow light curve of GRB990123. In the light curve of GRB021004, the initial fading phase was terminated by a short plateau phase that lasted for about 2 hours from 0.024 to 0.10 d after the burst (Uemura et al. 2003d). The object then entered an ordinary power-law fading phase. We propose that the plateau phase is evidence that the maximum of the synchrotron emission from a forward shock region appears around ~ 0.1 d after the burst, as expected from theoretical calculations (Sari, Piran 1999b; Kobayashi, Zhang 2003). The initial fading phase can be interpreted as part of the optical flash, which was recorded in GRB990123. If this is the case, the color of the afterglow would have dramatically changed from blue to red around the maximum, while no color information is available in this early phase (Sari et al. 1998). Detections the color change in the early phase will be an important future step for GRB astronomy. On the other hand, the feature around 0.1 d after the burst can be one of a series of bumps observed 1 d after the burst (Nakar et al. 2003; Lazzati et al. 2002; Fox et al. 2003; Schaefer et al. 2003; Li, Chevalier 2003).

GRB030329: The world-famous "monster GRB" GRB030329 (figure 19) occurred closest to us ($z = 0.168$), which gave a chance for us to study the detailed structure of the GRB afterglow (Vanderspek et al. 2003; Greiner et al. 2003; Caldwell et al. 2003). We detected a bright, 12th-mag optical afterglow 76 min after the burst (Peterson, Price 2003; Torii 2003; Uemura et al. 2003f). With the international collaboration through the VSNET, we obtained an ~ 11 hr continuous light curve of this afterglow (Uemura et al. 2003e). Our observation revealed that the afterglow experienced repetitive modulations even in the early phase of our observation. In conjunction with public data reported to the GCN Circulars, the light curve was rich in unexpectedly complicated structures throughout ~ 6 days after the burst. It is a surprise that the amplitude of modulations was almost constant with time. This feature of modulations is difficult to be understood with density variations of the interstellar medium (Nakar et al. 2003). The energy in the shock region must have changed with time, while the mechanism to generate additional energy is an open issue (Uemura et al. 2004b). The VSNET team also contributed to a another collaborative work on the earliest stage afterglow (Torii et al. 2003).

7. Conclusion

Variable Star Network (VSNET) is a global professional-amateur network of researchers in variable stars and related objects, particularly in transient objects, such as cataclysmic variables, black hole binaries,



Fig. 19. Fading afterglow of GRB030329. The images were taken by the VSNET Collaboration. The individual images (from left top to right bottom) correspond to 1.27, 3.37, 8.18 and 28.65 hr after the gamma-ray burst.

supernovae and gamma-ray bursts. The VSNET has been playing a pioneering role in establishing the field of *transient object astronomy*, by effectively incorporating modern advance in observational astronomy and global electronic network, as well as collaborative progress in theoretical astronomy and astronomical computing. The VSNET is now one of the best-featured global networks in this field of astronomy. We review on the historical progress, design concept, associated technology. We also review on the breathtaking scientific achievements, as well as regular variable star works, particularly focusing on dwarf novae (discovery of ER UMa stars, works in WZ Sge-type dwarf novae, more usual SU UMa-type dwarf novae, eclipsing dwarf novae), black hole X-ray transients (discoveries of an unexpected violent outburst of V4641 Sgr, rapid optical variations from the same object), and recent achievements in gamma-ray bursts.

We are grateful to Seiji Masuda and Katsura Matsumoto, who greatly contributed to the activities of the VSNET administrator group. We are grateful to many VSNET members who have been continuously supporting our activity. We are grateful to Emile Schweitzer (AFOEV), Keiichi Saijo and Makoto Watanabe (VSOLJ) for kindly allowing us to use AFOEV and VSOLJ public database for drawing light curves. We are also grateful to Dave Monet for making USNO A1 CD-ROMs readily available for us. This work is partly supported by a grant-in-aid [13640239, 15037205 (TK), 14740131 (HY)] from the Japanese Ministry of Education, Culture, Sports, Science and Technology. Part of this work is supported by a Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists (MU, RI). This research has made use of the astronomical catalogs at Astronomical Data Centers operated by National Astronomical Observatory, Japan, and NASA. This research has also made use of the Digitized Sky Survey produced by STScI, the ESO Skycat tool, the VizieR catalogue access tool, and the Electronic Edition of the GCVS.

Appendix 1. Poster Presented at the 23rd IAU General Assembly

Development of a Mailing List on Variable Stars, VSNET

(Daisaku Nogami, Taichi Kato,
Hajime Baba, Chatief Kunjaya)

Abstract

As the computer environment has been developing drastically these years, the style of astronomical study has been changed. The key words of these changes are thought to be "real-time" and "interactivity". Suspecting that they have potential to made an great effect on the study of transient objects (cataclysmic variables, X-ray binary, super novae, and so on), we set up the mailing list, VSNET, in 1994. Our policy on VSNET since the start is that VSNET is world-widely opened to any kinds of researchers including professionals and amateurs, observers and theorists. Subscribers have increased as time goes and now are over 400 from over 40 countries. Although VSNET started as a mailing list, it at present consists of five submaling lists, vsnet: vsnet-alert, vsnet-obs, vsnet-chat, and vsnet-chart, and each of these lists works independently for different purposes. Using the different characteristics of these sub-lists to advantage, various types of study have been proposed and carried out. VSNET will be developed further with cooperation of subscribers. If you have any comments or questions, please feel free to contact with VSNET administrators (vsnet-adm@kusastro.kyoto-u.ac.jp).

Introduction

The computer environment is making remarkable progress with development of the infrastructure and the machine power. The World-Wide Web (WWW, Berners-Lee et al. 1992) is explosively coming into wide as well, redrawing the common sense on the information distribution.

These have an great affect on the astronomical study, for example, 1) the complicated rapid control of telescopes (adaptive optics, etc.) becomes possible, 2) the scale of simulations are getting much larger, 3) editors of various journals encourage the electronic submission, 4) the electronic publication of papers on WWW or by CD-ROM is discussed and partly realized , 5) IAU circulars are distributed by e-mail, 6) preprints are usually distributed by e-mail or on web pages (e.g. http://www.lanl.gov, Ginsparg 1996), 7) the search of papers has become quite easy by the foundation of the Astrophysics Data System (NASA, Accomazzi et al. 1995), and so on.

The key words of this revolution are thought to be "realtime" and "interactivity".

Is there a possibility for these characteristics to be quite useful in the study of transient objects? We had an eye on the mailing list since the system makes it possible to share much information among much people quite quickly and can be used interactively by much people. Then, in July 1994, we started a mailing list, VSNET (vsnet@kusastro.kyoto-u.ac.jp), mainly on cataclysmic variables, supernovae, X-ray binaries, and so on, with a policy that we aim to make a contribute to the astronomical community by providing a mechanism to make information shared and a room to discuss all aspects of the astronomy concerning those transient objects among subscribers from all over the world irrespective of the status (an amateur or a professional), the style of astronomical study (an observer or a theorist), and other properties.

History of VSNET

The first member of VSNET is several tens of amateurs belongs to amateur associations (The American Association of Variable Stars (AAVSO, see http://www.aavso.org/), Association Française des Observateurs d'Etoiles Variables (AFOEV), Variable Star Observer League in Japan (VSOLJ), and so on) and a few tens of professionals. Subscribers, then, have increased by mouse-to-mouse advertising and our invitation to authors of papers in astronomical journals. Now the number of subscribers from about 50 countries exceeds 400, which means that major part of researchers in this field has already subscribed.

At first, all e-mails other than administrative ones for subscription and unsubscription were distributed to all subscribers. However, accepting a request mainly from theorists that they do not need daily data but want to know final results deduced from observations, we modified the system of VSNET to divide VSNET to three sub-mailing lists in October 1994, 1) vsnet-obs (vsnet-obs@kusastro.kyoto-u.ac.jp) for reporting daily observations, 2) vsnet-alert (vsnet-alert@kusastro.kyoto-u.ac.jp) for alert on the discovery of supernovae, novae, rare outbursts, discovery of new variable star, dramatic change of known variable stars, and so on, and 3) vsnet (vsnet@kusastro.kyoto-u.ac.jp) for the general information not suited for the former two lists, for example, complied data, finding charts, preprints, calls for observations for international co-observation campaigns, and so on.

In order to refer to an old log correctly, serial numbers in the subject were added as [vsnet-obs 1997] since January 1995.

In June 1995, the VSNET Web pages (http://www.kusastro.kyoto-u.ac.jp/vsnet/) were opened. You can see all articles ever posted to VSNET on WWW. You can see light curves drawn from observations distributed via vsnet-obs in the recommended format, too. The most important results among vsnet-alert logs and public information like conference announcement are gathered on the top page almost daily up-dated. Many useful tools developed and provided by various groups are available on the pages as well. At the same time we started VSNET anonymous ftp service (ftp.kusastro.kyoto-u.ac.jp/vsnet/) where you can get objects almost same as on WWW.

In January 1997, we made an addition of a new sub-mailing-list of vsnet-chat (vsnet-chat@kusastro.kyoto-u.ac.jp) in order to discuss various subjects. Though VSNET was used nearly only for distribution of information, this modification widened the usage of VSNET.

Having the background as mentioned here, VSNET at present consists of four sub-mailing-lists (vsnet, vsnet-alert, vsnet-obs, and vsnet-chat) as well as VSNET Web pages and anonymous ftp service. However, one of the sub-lists, vsnet, is now closed, since commercial informations never related to astronomy, so-called SPAM, were posted again and again. This type of problem, although common to all mailing lists, is hard to completely solve.

The number of e-mail posted to vsnet-obs, vsnet-alert, and vsnet-chat is 20-40, a few, ∼5 per day, although changing from day to day.

Tables show several e-mails posted to VSNET andf igures show light curves available on WWW.

VSNET: Now and Future

The major usages of VSNET are 1) to forecast the behavior of variable stars from daily observations available via vsnet-obs and plot a strategy of the observation, 2) to check the status of a variable star in optical on vsnet-obs at the time of the observations in UV or X-ray, 3) to publish new results on vsnet-alert (discoveries of supernovae, determination of superhump period in SU UMa-type dwarf novae, and so on), 4) to call the follow-up observations of transient objects detected in UV or X-ray, 5) to call the co-operation for international co-observation campaigns, 6) to notify conferences newly held, 7) to discuss various subjects ranging from the first step of the observation to theoretical interpretations of new interesting phenomena, and so on. For reference, tables list a part of subjects of e-mails posted to VSNET. These usages would be born in the active environment under the complex of "real-time", "interactivity", and the VSNET policy.

In addition, new original applications of VSNET are appreciated all time, and, actually, will be proposed. We, VSNET administrators, would be much glad if studies completed on use of VSNET contribute to the progress of astrophysics. If you have any questions, requests, and suggestions, please do not hesitate to communicate with VSNET administrators (vsnet-adm@kusastro.kyoto-u.ac.jp).

Part of this work was supported by a Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists (D.N.).

References

Accomazzi, A., Eichhorn, G., Grant, C. S., Murray, S. S., and Kurtz, M. J. 1995, Vistas in Astronomy 39, 63

Berners-Lee, T. J., et al. 1992, in "Electronic Networking: Research, Applications and Policy" (Meckler Publishing, Westport), Vol. 2, No.1, p52

Ginsparg, P. 1996, http://xxx.lanl.gov/blurb/pg96unesco.html

Appendix 2. VSNET Data Reporting System

As described in subsection 2.1, the VSNET data handling, database managing, and data analysis tools are mostly common to the ones developed for the VSOLJ database project (Kato 1990a), and later adapted for wider range of observations. The programs were originally written for microcomputers running on MS-DOS$^{\mathrm{TM}}$, and later ported to Linux. The following information is mainly an excerpt from Kato (1990a), partly rewritten for the recent changes for the VSNET management.

A.2.1. *Electronic Format of Observations*

The VSNET reporting format is an UT extension of the VSOLJ electronic reporting format, whose design was established in 1987. The standard format VSNET observations are composed of lines separated by new-line characters. Since a full description of the format used in these data or files may not be necessary to all readers, the minimal requirements to interpret the data which are made available through the VSNET World-Wide Web service.

Each line contains the following items.

(a) Name of the object
(b) Time of the observation in decimals of UT
(c) Observed magnitude
(c') Magnitude system (CCD/photoelectric) or film and filter (photographic)
(d) Observer's code

These items were designed to express original observations as exactly as possible (mainly in terms of significant digits). Comparison stars, charts used, or any other text information can be written as a comment following the item (d); they are usually used for future reference or examination, and not directly used at present in the VSNET regular database management.

Each item is separated by one or more space characters (ASCII code 20 in hexadecimal) and does not contain spaces within the item.

(a) Name of the Object

This is an identifier for the object. If the name is unique enough to discriminate the object from other celestial objects, any expression is basically allowed; the use of the names listed in the General Catalogue of Variable Stars (GCVS), Beyer names, numbers in the New catalogue of Suspected Variable Stars (NSV), and Durchmusterung numbers (BD, CD, CPD), however, is strongly recommended. When there is no specific relevant catalog, the use of GSC and USNO identifier can be used.

For GCVS and Beyer names, three-letter IAU code of constellation (in upper-case letters) precedes the name of the star in the constellation. Greek letters are written in the standard English expressions.

Examples: ANDR for R And,
ORIalpha for Betelgeuse (α Ori)

Translation of reported names into standard expressions, when necessary, is performed either automatically or manually using the *ALIAS* database. This process particularly becomes necessary when new GCVS names are released in the form of regular name-list updates, or when a new GCVS designation is given for a new nova. All the name translation rules are centralized in the *ALIAS* database and reflected on the entire VSNET system, eliminating additional efforts to modify individual observation reports or expressions in VSNET circulars. This is an great advantage of the VSNET database managing process, and a newly released name-list update (containing some hundreds of newly designated variable stars) can be usually reflected within a day of the release. Observers are allowed to continue using the old expressions as long as the expressions can be uniquely and automatically translated into the new expressions.

(b) Time of the Observation

In order to facilitate detecting errors by eye, times are expressed in the following decimal format using UTC.

Example: 20030701.123 (2003 July 1.123 UTC).

No heliocentric or barycentric corrections are introduced at this stage. The conversion to TAI or TD (in any expression including Julian Date), or helio-(or bary-)centric corrections are left to data analysis software and users. These measures are partly because widely used software packages at the observer's end are known to frequently contain problems. The other reason is that leap seconds do not allow to distribute fixed tables for conversion beforehand.

(c) Observed Magnitude

Decimal points are explicitly used to show the significant digits. Upper limit observation (non-detection) is expressed by a prefix '<'. Expression may be followed by one of ':' or '?' to show uncertainty.

(c') Magnitude System, Film and Filter

If the observation is visually performed, this item is not necessary. Otherwise, the code of the magnitude system (or the film and the filter if the observation is done photographically) follows the item (c) without placing a space. Well-defined standard photoelectric systems (e.g. Johnson-Cousins U, B, V, $R_{\rm c}$, $I_{\rm c}$, $R_{\rm j}$, $I_{\rm j}$) systems are used as in usual sense. Other VSNET-specific codes include "C" (unspecified unfiltered CCD magnitude), "CR" (unfiltered CCD magnitude calibrated on $R_{\rm c}$-band) and "p" (unspecified photographic magnitude). The complete list of the codes representing the films and filters at present is available at the VSNET website.[108]

Example: <15.5Rc (fainter than 15.5 in $R_{\rm c}$-band)

(d) Observer's code

The code is usually a three-letter code, as canonically used by the American Association of Variable Star Observers (AAVSO)[109] and the VSOLJ. The code may be immediately followed by a period mark ('.') and the organization code. When there is no fixed affiliation for the observer, the VSNET manager group issues a code, which scheme also enables archiving historical or literature observations.

With this format, each line of observation has all information in itself; any operation of moving lines in a data file (such as sorting) does not affect the properties of the data. The above design of electronic data has greatly facilitated overall aspects of the following electronic data management, and this pioneering concept has been taken over by a number of world-wide variable star organizations.

Appendix 3. Design of the VSNET Database

Most of programs are written in the language C, and were originally compiled with Turbo-C (Borland International). Many of machine-independent source codes (handling text-based data) can be compiled by ANSI C (ANSI X3.159) compilers (such as gcc) without much correction. In the actual VSNET database management, these codes are either compiled on Windows$^{\rm TM}$/DOS personal computers, or Linux workstations. Most of the text-based works are currently done on Linux workstations.

Besides storing original data as text files (as reported in vsnet-obs), they are also incorporated into an observational database in rewritable media allowing random access. Several interesting graphic programs and data analysis programs run in this mode. The maximum number of observations handled at one time depends only on capacity of the media and the addressing capacity of the operating system. For example, we now handle dynamic and random access to the entire VSNET observations, consisting of more than 1.2 million visual observations (~130 MB) of randomly accessible data, and we have registered more than 1.3 million CCD observations submitted to the VSNET Collaboration. We have also confirmed that a combination test of the VSNET, VSOLJ and AFOEV public data (about 3 million observation) has yielded satisfactory efficiency of handling a huge volume of randomly accessible data.

In addition to the observational database, there also exist other system databases including *VARTYPE* (database on individual variable star types), and *ALIAS* (name resolver database). A programmer on this system can inquire these databases for the type of variability or the standard expression of the name for the object specified by a given identifier.

A.3.1. Structure of the Database

Because the number of the data or the objects is very large, and both random and sequential accesses are necessary for easy operation on the light curves, we adopted the combination of B+tree and bidirectional linear list. This basic structure of the database was originally established in 1989 to fully incorporate the VSOLJ database.

Complete data of observations (other than remarks) are stored in a file equivalent to an index file of an relational database, and there is no need for reading an additional data file. The size of a storage block is 16 kilobytes, but this can be modified upon different compilation. The main memory is dynamically allocated to simulate a virtual memory in order to minimize access to the storage media. With this feature, the main database module only requires an order of a few hundreds of megabytes of actual memory. The kernel of the database module is written transparently to the upper modules, so upper modules have only to pass the key or the virtual address to the kernel to inquire the next or the previous data. The modules are written to enable simultaneous handling of multiple databases (e.g. observational database and *VARTYPE*) without interference. The database functions are prepared as a form of C-language based application interface (API), but we skipped the details of individual APIs because they are too technical to be presented in this paper.

---

[108] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/etc/format.html⟩.

[109] ⟨http://www.aavso.org⟩.

A.3.2. Basic Command Operations

Basic user operations to the database can be performed by command-line tasks on MS-DOS$^{\mathrm{TM}}$ or Linux. The most frequently used and basic commands include:

inittree	creates a new database.
mrg	merges a text file to a database.
vdel	deletes data specified by a text file from a database.
vmrg	merges an observational data file in the standard format to the observational database, and lists potential errors by referring to the previously registered data and *VARTYPE* database.
vcut	lists data of a given object for a given chronological period.
vren	replaces an identifier for an object by another.
vlist	lists all data in a database.
liststar	lists all stars in a database.
seltype	selects data from a text file by specifying types of variability, by referring to the *VARTYPE* database.
chknew	checks observation data file if they contain newly reported objects by comparison with the existing observational database..
chkstd	checks observation data file for grammar, and lists discordant data and potential errors by referring to the previously registered data and *VARTYPE* database.
looktype	returns the variable star type by referring to the *VARTYPE* database.
settype	sets or modifies the variable star type in the *VARTYPE* database.
calias	converts a variable star name to the standard name by referring to *ALIAS* database.
alias	sets or modifies the entry in the *ALIAS* database (this command has a different name on Linux because of the collision with the shell built-in command).

Most basic operations on the VSNET data are done with these basic commands, and are frequently used as a combination regulated by a shell script.

A.3.3. Data Analysis Application Programs

Most of these programs use a GUI. These packages are architecture-specific, and presently run in limited environments (e.g. NEC PC 9801 machines), although efforts have been taken to port these applications to Windows$^{\mathrm{TM}}$ operating system, or to write an equivalent wrapper GUI applications written in Java$^{\mathrm{TM}}$. Figure 20 shows a sample image by **grp** interactive light curve viewer program.

grp	displays light curves, and enables interactive zooming, data selection, data editing.
autogrp	displays and prints light curves automatically. An equivalent package written in Java$^{\mathrm{TM}}$ is presently used to produce VSNET online light curves.
circ	automatically produces VSNET CV Circulars.

A.3.4. Data Processing

Besides drawing light curves, there also exist several kinds of scientific data-analysis programs. Some require databases and others require only data files in the text format. A few samples are shown here. Many of the general-purpose programs (mostly with source codes) are available from the VSNET website "Tools and Programs" section.[110]

Heliocentric and Barycentric Corrections

We have implemented both conventional heliocentric corrections for the observed JDs, by using the well-known Newcomb's expansion of the planets. We have also implemented barycentric corrections by numerically integrating the DE 200 ephemeris produced by NASA. The original source code of the DE 200 ephemeris was imported from a software package written by Mr. Novas.

Period Analysis Tools

We have our own implementations of Discrete Fourier Transform (DFT), Phase Dispersion Minimization (PDM: Stellingwerf 1978), Discrete Wavelet Transform (DWT: Foster 1996) and other period analysis tools. The PDM package was ported to Windows$^{\mathrm{TM}}$ by Andreas Wijaya, and has been conveniently used by many users.

CV Circulars

CV Circulars were originally prepared monthly and issued by the VSOLJ, from the observations reported by the end of the next month. In the modern VSNET service, these circulars are issued almost daily to fully incorporate the daily changes in the rapidly varying CVs. The corresponding list is *vsnet-cvcirc*.[111]

CV Circulars contain information about all reported outbursts and standstills of dwarf novae in the form of nightly averages, and nightly averages of other peculiar objects (CVs other than dwarf novae, X-ray binaries, symbiotic variables, eruptive variable such as R CrB stars, S Dor stars, FU Ori stars, and supernovae, active galactic nuclei and other objects of special interest).

Unpredictable variations of brightness of these objects make it extremely difficult to check observations as automatically done for pulsating variables. Together with faintness of these objects (which implies existence of large number of negative observations), cross checks between different observers are indispensable.

The present program **circ**, referring to a list of more than 1000 objects containing their properties (type of variability, normal range of variation, ephemerides of eclipses etc.), checks all observations using databases. The list is updated by the editor whenever new information becomes available in order to maintain circulars up-to-date.

---

[110] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/etc/prog.html⟩.

[111] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-cvcirc/maillist.html⟩.



Fig. 20. Sample image by **grp** interactive light curve viewer program. The light curve is BL Lac from VSNET observations. White and red marks represent positive and upper limit observations, respectively. The program enables interactive GUI-based operation on photometric database constructed on the database engine described in Appendix 3.1.

The results are then listed as a file in the prototypical form of CV Circulars, containing the reports of potentially discordant data (or rapid intrinsic changes) with special marks. The present version can handle more than tens of thousands of observations per month, and produces a circular in a minute.

The same program, slightly modified to produce long-term averages, is used to produce VSNET Mira Circulars *vsnet-miracirc*.[112]

Appendix 4. Chart-Plotting and Star Identification Program

When identifying variable stars and locating the CCD field of view, chart-plotting and star identification programs are very useful, particularly when the network resource was unavailable. For this purpose, one of the authors (TK) developed in 1990 a chart-plotting computer program, when machine-readable Guide Star Catalog (GSC) 1.0 (Lasker et al. 1990) was released. The program was designed to run on a stand-alone personal computer running on MS-DOS$^{\mathrm{TM}}$ non-extended (less than 640 KB) memory.

Since the distribution form of GSC 1.0 was composed of huge ASCII tables on two CD-ROMs (1.2 GB), and since the objects were randomly arranged within individual files covering 2–3 degrees square, it was an absolute requirement to compress these data and make them quickly and randomly accessible. In this software, we subdivided the entire GSC data into 0.5 degree bins in declination, and sorted the objects within the same bin according to right ascension. In order to compress the data, we used binary files which can be directly mapped in C structure. In order to avoid redundancy in right ascensions and enable quick random access, we used a separate jump table which records the file positions of given coordinate meshes. Individual entries in the 0.5-deg bins only contain residuals to the mesh coordinates. In original compression of GSC 1.0, each catalog entry corresponded to 5 bytes. By combining the information of coordinate mesh in the jump table and the individual entries, one can obtain the fully decoded coordinates. These functions (including sequential reading functions of a given box) were implemented as transparent APIs, many of which were designed to take a pointer to the display function as a call-back function. With this compression, the entire GSC 1.0 (disregarding the object names, plate numbers) can be compressed into 100 MB. This program was one of the earliest chart-plotting software packages that used GSC as source catalogs and implement encoded compressed catalogs, and the design was taken over in various successive third-party applications. The program is also able to display GCVS and NSV variable star catalogs, and IRAS PSC objects, which were compressed in a similar manner employed in GSC compression.

---

[112] ⟨http://www.kusastro.kyoto-u.ac.jp/vsnet/Mail/vsnet-miracirc/maillist.html⟩.

The program was later updated to accommodate GSC 1.1 (Lasker et al. 1996) and USNO A1.0 (Monet et al. 1998a) catalogs in 1998, and other catalogs (with variable-length object labels). With the increase of capacity of storage media, the present version uses 12 bytes for one GSC 1.1 full entry and 8 bytes for one USNO A1.0 entry. With this compression method, the entire USNO A1.0 data (more than 4.8×10^8 objects) can be stored in 4 GB storage and can be quickly and randomly accessed. The present version is equipped with a function to handle the 2MASS point source catalog in the same manner. The program also handles a name resolver by using the same database engine described in Appendix 3.1. The entire program is presently ported to Windows$^{\mathrm{TM}}$ and Linux using *xlib* graphic library (see figure 21 for an example). With this program and APIs, one can very quickly identify new and known variable stars either interactively or in a batch. Online VSNET charts with Hipparcos and Tycho magnitudes (subsection 2.2) have been prepared with this software operated in a batch mode. The object databases have been regularly updated, especially when a name lists for new variable stars is released. The source codes are available upon request to the author.

References

Aikawa, T. 1985, Ap&SS, 116, 401

Ak, T., Ozkan, M. T., & Mattei, J. A. 2001, A&A, 369, 882

Allen, D. A. 1984, Publ. Astron. Soc. Australia, 5, 369

Anupama, G. C. 1992, PASP, 104, 308

Anupama, G. C., & Dewangan, G. C. 2000, AJ, 119, 1359

Arkhipova, V. P. 1994, Pis'ma AZh, 20, 919

Arkhipova, V. P. 1996, Pis'ma AZh, 22, 828

Arkhipova, V. P., & Esipov, V. F. 1996, Pis'ma AZh, 22, 264

Arkhipova, V. P., et al. 1998a, Pis'ma AZh, 24, 297

Arkhipova, V. P., Esipov, V. F., & Sokol, G. V. 1998b, Pis'ma AZh, 24, 431

Arkhipova, V. P., Esipov, V. F., Sokol, G. V., & Shugarov, S. Y. 1999a, Astron. Letters, 25, 739

Arkhipova, V. P., & Noskova, R. I. 1997, Pis'ma AZh, 23, 709

Arkhipova, V. P., Noskova, R. I., Esipov, V. F., & Sokol, G. V. 1999b, Astron. Letters, 25, 615

Arkhipova, V. P., Zaitseva, G. V., Ikonnikova, N. P., Noskova, R. I., & Shugarov, S. Y. 1994, Pis'ma AZh, 20, 916

Arkhipova, V. P., Zaitseva, G. V., Ikonnikova, N. P., Noskova, R. I., & Shugarov, S. Y. 1996, Pis'ma AZh, 22, 836

Arkhipova, V. P., Zaitseva, G. V., Ikonnikova, N. P., Noskova, R. I., Sokol, G. V., & Shugarov, S. Y. 1998c, Pis'ma AZh, 24, 427

Arkhipova, V. P., Zaitseva, G. V., & Noskova, R. I. 1991, Astron. Tsirk., 1547, 9

Asplund, M., Gustafsson, B., Lambert, D. L., & Kameswara Rao, N. 1997, A&A, 321, L17

Asplund, M., Lambert, D. L., Kipper, T., Pollacco, D., & Shetrone, M. D. 1999, A&A, 343, 507

Augusteijn, T., & Stehle, R. 1995, in Cataclysmic Variables, ed. A. Bianchini, M. della Valle, & M. Orio (Dordrecht: Kluwer Academic Publishers), 103

Augusteijn, T., van der Hooft, F., de Jong, J. A., & van Paradijs, J. 1996, A&A, 311, 889

Augusteijn, T., van Kerkwijk, M. H., & van Paradijs, J. 1993, A&A, 267, L55

Ayani, K., Furusho, R., Kawakita, H., Fujii, M., & Yamaoka, H. 1999, IAU Circ., 7244

Ayani, K., Hashimoto, T., & Yamaoka, H. 2003, IAU Circ., 8048

Ayani, K., Kawabata, T., & Yamaoka, H. 2002, IAU Circ., 7864

Ayani, K., & Kawakita, H. 1995, IAU Circ., 6256

Ayani, K., Nakatani, H., & Yamaoka, H. 1998, IAU Circ., 6905

Ayani, K., & Yamaoka, H. 1997, IAU Circ., 6800

Ayani, K., & Yamaoka, H. 1998a, IAU Circ., 6878

Ayani, K., & Yamaoka, H. 1998b, IAU Circ., 6878

Ayani, K., & Yamaoka, H. 1998c, IAU Circ., 7059

Ayani, K., & Yamaoka, H. 1999a, IAU Circ., 7137

Ayani, K., & Yamaoka, H. 1999b, IAU Circ., 7336

Ayani, K., & Yamaoka, H. 1999c, IAU Circ., 7339

Ayani, K., & Yamaoka, H. 2002, IAU Circ., 7976

Baade, D., Rivinius, T., Stefl, S., Otero, S., & Liu, X. 2001, IAU Circ., 7658

Baba, D., Nagata, T., Iwata, I., Kato, T., & Yamaoka, H. 2003a, IAU Circ., 8112

Baba, H., Kato, T., Nogami, D., & Hirata, R. 1999, in Disk Instabilities in Close Binary Systems, ed. S. Mineshige, & J. C. Wheeler (Tokyo: Universal Academy Press), 123

Baba, H., Kato, T., Nogami, D., Hirata, R., Matsumoto, K., & Sadakane, K. 2000, PASJ, 52, 429

Baba, H., et al. 2003b, PASJ, submitted

Baba, H., et al. 2002, PASJ, 54, L7

Baba, H., Sadakane, K., Norimoto, Y., Nogami, D., Matsumoto, K., Makita, M., & Kato, T. 2001, IAU Circ., 7678

Bailey, J. 1979, MNRAS, 189, 41P

Baptista, R., & Catalán, M. S. 2000, ApJL, 539, L55

Baptista, R., & Catalán, M. S. 2001, MNRAS, 324, 599

Baptista, R., Catalán, M. S., Horne, K., & Zilli, D. 1998, MNRAS, 300, 233

Baptista, R., Harlaftis, E. T., & Steeghs, D. 2000, MNRAS, 314, 727

Baptista, R., Haswell, C. A., & Thomas, G. 2002, MNRAS, 334, 198

Baptista, R., & Steiner, J. E. 1991, A&A, 249, 284

Baptista, R., & Steiner, J. E. 1993, A&A, 277, 331

Barker, J., & Kolb, U. 2003, MNRAS, 340, 623

Barlow, M. J., et al. 1981, MNRAS, 195, 61

Baron, E., Hauschildt, P. H., Branch, D., Wagner, R. M., Austin, S. J., Filippenko, A. V., & Matheson, T. 1993, ApJL, 416, L21

Bateson, F., & Jones, A. F. 1987, IAU Circ., 4383

Bateson, F. M., Goltz, W., Jones, A. F., & McIntosh, R. 1986, Publ. Variable Stars Sect. R. Astron. Soc. New Zealand, 14, 52

Bedding, T. R., Zijlstra, A. A., Jones, A., Marang, F., Matsuura, M., Retter, A., Whitelock, P. A., & Yamamura, I. 2002, MNRAS, 337, 70

Bedient, J. 2002a, IAU Circ., 7785

Bedient, J. 2002b, IAU Circ., 7790

Belczynski, K., Mikolajewska, J., Munari, U., Ivison, R. J., & Friedjung, M. 2000, A&AS, 146, 407

Belloni, T., et al. 1999, ApJ, 527, 345

Bergner, Y. K., Miroshnichenko, A. S., Yudin, R. V., Kuratov, K. S., Mukanov, D. B., & Shejkina, T. A. 1995, A&AS, 112, 221

Bertola, F. 1964, Ann. d'Astrophys., 27, 298



Fig. 21. Sample chart of V4743 Sgr (Nova Sgr 2002 No. 3). The nova progenitor is marked.

Bianchini, A. 1988, Inf. Bull. Variable Stars, 3136

Bianchini, A. 1990, AJ, 99, 1941

Bianchini, A., Canterna, R., Desidera, S., & Garcia, C. 2003, PASP, 115, 474

Bianchini, A., Sabbadin, F., Favero, G. C., & Dalmeri, I. 1986, A&A, 160, 367

Biggs, J., & Walsh, S. 2001, IAU Circ., 7749

Bobinger, A., Horne, K., Mantel, K., & Wolf, S. 1997, A&A, 327, 1023

Bond, H. E., et al. 2003, Nature, 422, 405

Bond, I., Freeth, R. V., Marino, B. F., & Walker, W. S. G. 1987, Inf. Bull. Variable Stars, 3037

Bortle, J. E., et al. 1980, IAU Circ., 3541

Brown, N. J., Waagen, E. O., Scovil, C., Nelson, P., Oksanen, A., Solonen, J., & Price, A. 2002, IAU Circ., 7785

Bruch, A. 2000, A&A, 359, 998

Bruch, A., Beele, D., & Baptista, R. 1996, A&A, 306, 151

Bruch, A., Steiner, J. E., & Gneiding, C. D. 2000, PASP, 112, 237

Buat-Ménard, V., & Hameury, J.-M. 2002, A&A, 386, 891

Buckley, D. A. H. 2000, New Astron. Rev., 44, 63

Burlak, M. A., & Esipov, V. F. 2001, Astron. Letters, 27, 574

Caldwell, N., Garnavich, P., Holland, S., Matheson, T., & Stanek, K. Z. 2003, GCN, 2053

Cannizzo, J. K. 2001, ApJL, 561, L175

Cannizzo, J. K., Gehrels, N., & Mattei, J. A. 2002, ApJ, 579, 760

Chen, W., Shrader, C. R., & Livio, M. 1997, ApJ, 491, 312

Chornock, R., Filippenko, A. V., & Li, W. D. 2001, IAU Circ., 7701

Christy-Sackmann, I. J., & Despain, K. H. 1974, ApJ, 189, 523

Ciardi, D. R., Howell, S. B., Hauschildt, P. H., & Allard, F. 1998, ApJ, 504, 450

Clayton, G. C. 1996, PASP, 108, 225

Clayton, G. C. 2002, Ap&SS, 279, 167

Clayton, G. C., Hammond, D., Lawless, J., Kilkenny, D., Evans, T. L., Mattei, J., & Landolt, A. U. 2002, PASP, 114, 846

Clocchiatti, A., Wheeler, J. C., Barker, E. S., Filippenko, A. V., Matheson, T., & Liebert, J. W. 1995, ApJ, 446, 167

Cohen, R. D., Marcy, G. W., & Harlan, E. A. 1980, AJ, 85, 867

Colas, F., Chassagne, R., & Yamaoka, H. 2000, IAU Circ., 7378

Costero, R., Tapia, M., Méndez, R. H., Echevarría, J., Roth, M., Quintero, A., & Barral, J. F. 1986, Rev. Mexicana Astron. Astrof., 13, 149

Cowley, C. R., Jaschek, M., & Acker, A. 1985, A&A, 149, 224

Crause, L. A., Lawson, W. A., Kilkenny, D., van Wyk, F., Marang, F., & Jones, A. F. 2003, MNRAS, 341, 785

Cristiani, H., Duerbeck, H. W., & Seitter, W. C. 1985, IAU Circ., 4027

de Jager, C. 1998, Astron. Astrophys. Rev., 8, 145

de Jager, C., Lobel, A., Nieuwenhuijzen, H., & Stothers, R. 2001, MNRAS, 327, 452

Della Valle, M., & Iijima, T. 2002, IAU Circ., 7786

della Valle, M., & Livio, M. 1996, ApJ, 473, 240

Dimai, M., Migliardi, M., Yamaoka, H., & Li, W. 2000, IAU Circ., 7373

Downes, R., Hoard, D. W., Szkody, P., & Wachter, S. 1995, AJ, 110, 1824

Downes, R., Webbink, R. F., & Shara, M. M. 1997, PASP, 109, 345

Downes, R. A., & Margon, B. 1981, MNRAS, 197, 35P

Downes, R. A., Webbink, R. F., Shara, M. M., Ritter, H., Kolb, U., & Duerbeck, H. W. 2001, PASP, 113, 764
Drissen, L., Shara, M. M., Dopita, M., & Wickramasinghe, D. T. 1994, AJ, 107, 2172
Duerbeck, H. W. 1984, Inf. Bull. Variable Stars, 2502
Duerbeck, H. W. 1987, Space Sci. Rev., 45, 1
Duerbeck, H. W. 1988, A&A, 197, 148
Duerbeck, H. W. 1989, Messenger, 58, 34
Duerbeck, H. W. 1992, MNRAS, 258, 629
Duerbeck, H. W. 2002, Ap&SS, 279, 5
Duerbeck, H. W., Baptista, R., Dutra, C. M., & Sterken, C. 2002, IAU Circ., 7799
Duerbeck, H. W., & Benetti, S. 1996, ApJL, 468, L111
Duerbeck, H. W., Benetti, S., Gautschy, A., van Genderen, A. M., Kemper, C., Liller, W., & Thomas, T. 1997, AJ, 114, 1657
Duerbeck, H. W., et al. 2000, AJ, 119, 2360
Duerbeck, H. W., Pollacco, D., Verbunt, F., Geertsema, G., van den Berg, M., & Green, D. W. E. 1996, IAU Circ., 6328
Dyck, G., & Schmeer, P. 1988, IAU Circ., 4594
Elizalde, F., Jablonski, F., Cieslinski, D., & Baptista, R. 1995, IAU Circ., 6165
Elliot, J. L., & Liller, W. 1972, ApJL, 175, L69
Esipov, V. F., Kolotilov, E. A., Mikolajewska, J., Munari, U., Tatarnikova, A. A., Tatarnikov, A. M., Tomov, T., & Yudin, B. F. 2000, Astron. Letters, 26, 162
Evans, A., Eyres, S. P. S., Naylor, T., & Salama, A. 1998, A&A, 335, 292
Evans, A., Krautter, J., Vanzi, L., & Starrfield, S. 2001, A&A, 378, 132
Evans, R., et al. 2003, IAU Circ., 8171
Fabregat, J., Reig, P., & Otero, S. 2000, IAU Circ., 7461
Faulkner, J. 1971, ApJL, 170, L99
Feibelman, W. A., & Bruhweiler, F. C. 1990, AJ, 100, 1248
Filippenko, A. V. 2000, IAU Circ., 7421
Filippenko, A. V., & Chornock, R. 2002, IAU Circ., 7825
Filippenko, A. V., & Martin, E. L. 1997, IAU Circ., 6783
Filippenko, A. V., Matheson, T., & Ho, L. C. 1993a, ApJL, 415, L103
Filippenko, A. V., et al. 1993b, IAU Circ., 5760
Filippenko, A. V., & Moran, E. C. 1998, IAU Circ., 6809
Foster, G. 1996, AJ, 112, 1709
Fox, D. W., et al. 2003, Nature, 422, 284
Fried, R. E., Kemp, J., Patterson, J., Skillman, D. R., Retter, A., Leibowitz, E., & Pavlenko, E. 1999, PASP, 111, 1275
Fuhrmann, B. 1981, Inf. Bull. Variable Stars, 1905
Fujino, S., Fujino, K., Iida, M., & Kato, T. 1987, VSOLJ Variable Star Bull., 3, 9
Furtig, W., & Wenzel, W. 1962, Inf. Bull. Variable Stars, 12
Gal-Yam, A., Shemeer, O., & Dann, J. 2002a, IAU Circ., 7810
Gal-Yam, A., Shemeer, O., & Dann, J. 2002b, IAU Circ., 7819
Galama, T. J., et al. 1998, Nature, 395, 670
Gandet, T. L., Otero, S., Fraser, B., & West, J. D. 2002, Inf. Bull. Variable Stars, 5352
Gao, W., Li, Z., Wu, X., Zhang, Z., & Li, Y. 1999, ApJL, 527, L55
Garnavich, P., Jha, S., Kirshner, R., Challis, P., Balam, D., & Berlind, P. 1997a, IAU Circ., 6798
Garnavich, P., Jha, S., Kirshner, R., Challis, P., & Berlind, P. 1997b, IAU Circ., 6778
Garnavich, P., Jha, S., Kirshner, R., Challis, P., & Macri, L. 1997c, IAU Circ., 6796
Georgii, R., et al. 2002, A&A, 394, 517
Gonzalez, G., Lambert, D. L., Wallerstein, G., Rao, N. K., Smith, V. V., & McCarthy, J. K. 1998, ApJS, 114, 133
Gonzalez-Riestra, R. 1992, A&A, 265, 71
Goranskii, V. P., Kusakin, A. V., Metlova, N. V., Shugarov, S. Y., Barsukova, E. A., & Borisov, N. V. 2002, Astron. Letters, 28, 691
Green, D. W. E. 2003, IAU Circ., 8115
Green, R. F., Ferguson, D. H., Liebert, J., & Schmidt, M. 1982, PASP, 94, 560
Green, R. F., Schmidt, M., & Liebert, J. 1986, ApJS, 61, 305
Greenstein, J. L., & Kraft, R. P. 1959, ApJ, 130, 99
Greiner, J. 1998, A&A, 336, 626
Greiner, J. 2000, New Astron. Rev., 44, 149
Greiner, J., Alcala, J. M., & Wenzel, W. 1996, Inf. Bull. Variable Stars, 4338
Greiner, J., & Di Stefano, R. 2002, ApJL, 578, L59
Greiner, J., et al. 2003, GCN, 2020
Greiner, J., et al. 2001, A&A, 376, 1031
Greiner, J., Tovmassian, G. H., di Stefano, R., Prestwich, A., González-Riestra, R., Szentasko, L., & Chavarría, C. 1999, A&A, 343, 183
Gänsicke, B. T., Fried, R. E., Hagen, H.-J., Beuermann, K., Engels, D., Hessman, F. V., Nogami, D., & Reinsch, K. 2000, A&A, 356, L79
Hachisu, I., & Kato, M. 2001, ApJL, 553, L161
Hachisu, I., & Kato, M. 2003a, ApJ, in press (astro-ph/0308065)
Hachisu, I., & Kato, M. 2003b, ApJ, 588, 1003
Hachisu, I., & Kato, M. 2003c, ApJ, 590, 445
Hachisu, I., Kato, M., Kato, T., & Matsumoto, K. 2000a, ApJL, 528, L97
Hachisu, I., Kato, M., Kato, T., & Matsumoto, K. 2002, in ASP Conf. Ser. 261, The Physics of Cataclysmic Variables and Related Objects, ed. B. T. Gänsicke, K. Beuermann, & K. Reinsch (San Francisco: ASP), 629
Hachisu, I., Kato, M., Kato, T., Matsumoto, K., & Nomoto, K. 2000b, ApJL, 534, L189
Hachisu, I., Kato, M., & Nomoto, K. 1996, ApJL, 470, L97
Hachisu, I., Kato, M., & Nomoto, K. 1999a, ApJ, 522, 487
Hachisu, I., Kato, M., Nomoto, K., & Umeda, H. 1999b, ApJ, 519, 314
Hachisu, I., Kato, M., & Saio, H. 1989, ApJL, 342, L19
Haefner, R., Schoembs, R., & Vogt, N. 1979, A&A, 77, 7
Hameury, J.-M., Lasota, J.-P., & Huré, J. M. 1997, MNRAS, 287, 937
Harrison, T. E., Johnson, J. J., & Spyromilio, J. 1993, AJ, 105, 320
Harvey, D., Skillman, D. R., Patterson, J., & Ringwald, F. A. 1995, PASP, 107, 551
Haseda, K. 2000, IAU Circ., 7382
Haseda, K., Kadota, K., Yamaoka, H., Takamizawa, K., & Kato, T. 2001, IAU Circ., 7647
Haseda, K., West, D., Yamaoka, H., & Masi, G. 2002, IAU Circ., 7975
Haswell, C. A., Patterson, J., Thorstensen, J. R., Hellier, C., & Skillman, D. R. 1997, ApJ, 476, 847
Hayakawa, T., Ueda, T., Uemura, M., Ohashi, H., Kato, T., & Matsumoto, K. 1998, Inf. Bull. Variable Stars, 4615
Hazen, M. L. 1986, Inf. Bull. Variable Stars, 2880
Heise, J., Mewe, R., Kruszewski, A., & Chlebowski, T. 1987, A&A, 183, 73
Hellier, C. 1996, in IAU Colloq. 158, Cataclysmic Variables and Related Objects, ed. A. Evans, & J. H. Wood (Dordrecht: Kluwer Academic Publishers), 143

Hellier, C. 2001a, Cataclysmic Variable Stars: how and why they vary. (Berlin: Springer-Verlag)
Hellier, C. 2001b, PASP, 113, 469
Hellier, C., & Buckley, D. A. H. 1993, MNRAS, 265, 766
Hellier, C., Kemp, J., Naylor, T., Bateson, F. M., Jones, A., Overbeek, D., Stubbings, R., & Mukai, K. 2000, MNRAS, 313, 703
Hellier, C., Mason, K. O., Rosen, S. R., & Cordova, F. A. 1987, MNRAS, 228, 463
Hellier, C., Mason, K. O., Smale, A. P., Corbet, R. H. D., O'Donoghue, D., Barrett, P. E., & Warner, B. 1989, MNRAS, 238, 1107
Henden, A. A., Munari, U., & Sumner, B. 2001, Inf. Bull. Variable Stars, 5140
Herbig, G. H., & Boyarchuk, A. A. 1968, ApJ, 153, 397
Herwig, F. 2001, ApJL, 554, L71
Hessman, F. V., Mantel, K.-H., Barwig, H., & Schoembs, R. 1992, A&A, 263, 147
Hildebrand, R. H., Spillar, E. J., Middleditch, J., Patterson, J. J., & Stiening, R. F. 1980, ApJL, 238, L145
Hils, D., & Bender, P. L. 2000, ApJ, 537, 334
Hirosawa, K., Yamamoto, M., Nakano, S., Kojima, T., Iida, M., Sugie, A., Takahashi, S., & Williams, G. V. 1995, IAU Circ., 6213
Hirose, M., & Osaki, Y. 1990, PASJ, 42, 135
Hirose, Y., & Yamaoka, H. 2003, IAU Circ., 8101
Hjellming, R. M., & Mioduszewski, A. J. 1998, IAU Circ., 6857
Hjellming, R. M., Mioduszewski, A. J., Ueda, Y., Ishida, M., Inoue, H., Dotani, T., Lewin, W. H. G., & Greiner, J. 1998, IAU Circ., 6872
Hjellming, R. M., et al. 2000, ApJ, 544, 977
Hoard, D. W., Szkody, P., Ishioka, R., Ferrario, L., Gänsicke, B. T., Schmidt, G. D., Kato, T., & Uemura, M. 2002a, AJ, 124, 2238
Hoard, D. W., Wachter, S., Clark, L. L., & Bowers, T. P. 2002b, ApJ, 565, 511
Honeycutt, R. K. 1992, PASP, 104, 435
Honeycutt, R. K., Cannizzo, J. K., & Robertson, J. W. 1994, ApJ, 425, 835
Honeycutt, R. K., Robertson, J. W., & Turner, G. W. 1995, in Cataclysmic Variables, ed. A. Bianchini, M. della Valle, & M. Orio (Dordrecht: Kluwer Academic Publishers), 75
Honeycutt, R. K., Robertson, J. W., Turner, G. W., & Mattei, J. A. 1998, PASP, 110, 676
Horne, K. 1985, MNRAS, 213, 129
Horne, K., & Cook, M. C. 1985, MNRAS, 214, 307
Horne, K., & Stiening, R. F. 1985, MNRAS, 216, 933
Howell, S., & Szkody, P. 1988, PASP, 100, 224
Howell, S. B., Adamson, A., & Steeghs, D. 2003, A&A, 399, 219
Howell, S. B., & Ciardi, D. R. 2001, ApJL, 550, L57
Howell, S. B., DeYoung, J. A., Mattei, J. A., Foster, G., Szkody, P., Cannizzo, J. K., Walker, G., & Fierce, E. 1996, AJ, 111, 2367
Howell, S. B., Dobrzycka, D., Szkody, P., & Kreidl, T. J. 1991, PASP, 103, 300
Howell, S. B., Gänsicke, B. T., Szkody, P., & Sion, E. M. 2002, ApJ, 575, 419
Howell, S. B., & Hurst, G. M. 1994, Inf. Bull. Variable Stars, 4043
Howell, S. B., & Hurst, G. M. 1996, J. British Astron. Assoc., 106, 29
Howell, S. B., Rappaport, S., & Politano, M. 1997, MNRAS, 287, 929
Howell, S. B., & Skidmore, W. 2000, New Astron. Rev., 44, 33
Howell, S. B., & Szkody, P. 1990, ApJ, 356, 623
Howell, S. B., & Szkody, P. 1995, in Cataclysmic Variables, ed. A. Bianchini, M. della Valle, & M. Orio (Dordrecht: Kluwer Academic Publishers), 335
Howell, S. B., Szkody, P., & Cannizzo, J. K. 1995, ApJ, 439, 337
Howell, S. B., Szkody, P., Kreidl, T. J., Mason, K. O., & Puchnarewicz, E. M. 1990, PASP, 102, 758
Hu, J. Y., Qiu, Y. L., Qiao, Q. Y., & Wei, J. Y. 1997, IAU Circ., 6783
Hu, J. Y., Wang, L. F., Jiang, X. J., Hanzl, D., Kato, M., Karlovsky, V., & Prugniel, P. 1993, IAU Circ., 5783
Huang, M., & Wheeler, J. C. 1989, ApJ, 343, 229
Hudec, R., Huth, H., & Fuhrmann, B. 1984, Observatory, 104, 1
Hurst, G. M., et al. 1998, IAU Circ., 7033
Hurst, G. M., Lubbock, S., & Buczynski, D. 1985, IAU Circ., 4130
Hurst, G. M., McAdam, D., Mobberley, M., & James, N. 1988a, IAU Circ., 4570
Hurst, G. M., & Young, A. 1988, IAU Circ., 4579
Hurst, G. M., Young, A., Manning, B., Mobberley, M., Oates, M., Boattini, A., & Scovil, C. 1988b, IAU Circ., 4577
Hutchings, J. B., & Cote, T. J. 1985, PASP, 97, 847
Hutchings, J. B., & Cote, T. J. 1986, PASP, 98, 104
Hynes, R. I., et al. 2002, A&A, in press (astro-ph/0207026)
Hynes, R. I., Mauche, C. W., Haswell, C. A., Shrader, C. R., Cui, W., & Chaty, S. 2000, ApJL, 539, L37
Iben, I. J., & Livio, M. 1993, ApJL, 406, L15
Ichikawa, S., Mineshige, S., & Kato, T. 1994, ApJ, 435, 748
Ichikawa, S., & Osaki, Y. 1994, in Theory of Accretion Disks-2, ed. W. J. Duschl, J. Frank, F. Meyer, E. Meyer-Hofmeister, & W. M. Tscharnuter (Dordrecht: Kluwer Academic Publishers), 169
Iida, M. 1994, VSOLJ Variable Star Bull., 19, 2
Iida, M., Kato, T., Nogami, D., & Baba, H. 1995a, Inf. Bull. Variable Stars, 4279
Iida, M., Nogami, D., & Kato, T. 1995b, Inf. Bull. Variable Stars, 4208
Iida, M., & York, D. 1994, IAU Circ., 6056
Iijima, T. 1996, MNRAS, 283, 141
Iijima, T. 2002, A&A, 387, 1013
Iijima, T., & Rosino, L. 1996, IAU Circ., 6365
Iijima, T., & Rosino, L. 1997, IAU Circ., 6703
Iijima, T., Rosino, L., & della Valle, M. 1998, A&A, 338, 1006
Iijima, T., & Strafella, F. 1993, Inf. Bull. Variable Stars, 3959
Ikeda, Y., Kawabata, K. S., & Akitaya, H. 2000, A&A, 355, 256
Imada, A., Kato, T., & Yamaoka, H. 2003, GCN, 2370
Ingram, D., Garnavich, P., Green, P., & Szkody, P. 1992, PASP, 104, 402
Ioannou, Z., Naylor, T., Welsh, W. F., Catalán, M. S., Worraker, W. J., & James, N. D. 1999, MNRAS, 310, 398
Ishida, M., Yamashita, A., Ozawa, H., Nagase, F., & Inoue, H. 1996, IAU Circ., 6340
Ishioka, R., Kato, T., Matsumoto, K., Uemura, M., Iwamatsu, H., & Stubbings, R. 2001a, Inf. Bull. Variable Stars, 5023
Ishioka, R., et al. 2002a, PASJ, 54, 581
Ishioka, R., Kato, T., Uemura, M., Iwamatsu, H., Matsumoto, K., Martin, B. E., Billings, G. W., & Novak, R. 2001b, PASJ, 53, L51
Ishioka, R., et al. 2001c, PASJ, 53, 905
Ishioka, R., et al. 2003, PASJ, 55, 683

Ishioka, R., Uemura, M., & Kato, T. 2001d, Inf. Bull. Variable Stars, 5028
Ishioka, R., Uemura, M., Kato, T., Iwamatsu, H., Matsumoto, K., Billings, G., Masi, G., & Kiyota, S. 2000a, IAU Circ., 7552
Ishioka, R., Uemura, M., Matsumoto, K., Iwamatsu, H., Kato, T., & Yamaoka, H. 2000b, GCN, 902, 1
Ishioka, R., et al. 2001e, IAU Circ., 7669
Ishioka, R., et al. 2002b, A&A, 381, L41
Ishioka, R., Uemura, M., T., Kato, & Team, The VSNET Collaboration 2002c, in ASP Conf. Ser. 261, The Physics of Cataclysmic Variables and Related Objects, ed. B. T. Gänsicke, K. Beuermann, & K. Reinsch (San Francisco: ASP), 491
Israelian, G., Lobel, A., & Schmidt, M. R. 1999, ApJL, 523, L145
Iwamoto, K., et al. 1998, Nature, 395, 672
Iwamoto, K., et al. 2000, ApJ, 534, 660
Jablonski, F., & Busko, I. C. 1985, MNRAS, 214, 219
Jameson, R. F., King, A. R., & Sherrington, M. R. 1982a, MNRAS, 200, 455
Jameson, R. F., Sherrington, M. R., King, A. R., & Frank, J. 1982b, Nature, 300, 152
Jeffery, C. S., & Pollacco, D. 2002, Ap&SS, 279, 15
Jha, S., Challis, P., & Kirshner, R. 2000a, IAU Circ., 7437
Jha, S., Challis, P., Kirshner, R., Yamaoka, H., & Ewalt, F. M. 2000b, IAU Circ., 7438
Jha, S., Garnavich, P., Challis, P., Kirshner, P., & Calkins, M. 1998a, IAU Circ., 7054
Jha, S., Garnavich, P., Challis, P., Kirshner, R., & Berlind, P. 1998b, IAU Circ., 6983
Jones, A. 1994, IAU Circ., 6120
Jones, A., & Pearce, A. 1990, IAU Circ., 5004
Jurcsik, J. 1993, Acta Astron., 43, 353
Jurcsik, J., & Montesinos, B. 1999, New Astron. Rev., 43, 415
Kahabka, P., Hartmann, H. W., Parmar, A. N., & Negueruela, I. 1999, A&A, 347, L43
Kaluzny, J. 1986, IAU Circ., 4287
Kamath, U. S., & Ashok, N. M. 2002, Ap&SS, 279, 65
Kanatsu, K., Kato, T., & Liller, W. 2001, IAU Circ., 7650
Kato, D., Nagata, T., Kato, T., Uemura, M., & Yamaoka, H. 2003a, IAU Circ., 8056
Kato, T. 1990a, in Light Curves of Variable Stars by VSOLJ 1. Mira Type Variable Stars for JD 2445950–2447050, ed. M. Watanabe, K. Hirosawa, & K. Saijo (Tokyo: Variable Star Observers League in Japan), 2
Kato, T. 1990b, Inf. Bull. Variable Stars, 3522
Kato, T. 1991a, Inf. Bull. Variable Stars, 3671
Kato, T. 1991b, IAU Circ., 5379
Kato, T. 1993, PASJ, 45, L67
Kato, T. 1994, Inf. Bull. Variable Stars, 4136
Kato, T. 1995a, Inf. Bull. Variable Stars, 4152
Kato, T. 1995b, Inf. Bull. Variable Stars, 4236
Kato, T. 1995c, Inf. Bull. Variable Stars, 4239
Kato, T. 1995d, Inf. Bull. Variable Stars, 4242
Kato, T. 1995e, Inf. Bull. Variable Stars, 4243
Kato, T. 1995f, Inf. Bull. Variable Stars, 4256
Kato, T. 1996a, Inf. Bull. Variable Stars, 4369
Kato, T. 1996b, PASJ, 48, 777
Kato, T. 1997a, PASJ, 49, 583
Kato, T. 1997b, VSOLJ Variable Star Bull., 25, 2
Kato, T. 1999a, Inf. Bull. Variable Stars, 4745
Kato, T. 1999b, Inf. Bull. Variable Stars, 4761
Kato, T. 1999c, Inf. Bull. Variable Stars, 4762
Kato, T. 1999d, Inf. Bull. Variable Stars, 4764
Kato, T. 1999e, Inf. Bull. Variable Stars, 4768
Kato, T. 1999f, Inf. Bull. Variable Stars, 4769
Kato, T. 1999g, Inf. Bull. Variable Stars, 4788
Kato, T. 1999h, Inf. Bull. Variable Stars, 4789
Kato, T. 1999i, Inf. Bull. Variable Stars, 4790
Kato, T. 1999j, Inf. Bull. Variable Stars, 4791
Kato, T. 1999k, Inf. Bull. Variable Stars, 4799
Kato, T. 1999l, VSOLJ Variable Star Bull., 33, 3
Kato, T. 2000, Inf. Bull. Variable Stars, 4918
Kato, T. 2001a, PASJ, 53, L17
Kato, T. 2001b, Inf. Bull. Variable Stars, 5071
Kato, T. 2001c, Inf. Bull. Variable Stars, 5092
Kato, T. 2001d, Inf. Bull. Variable Stars, 5093
Kato, T. 2001e, Inf. Bull. Variable Stars, 5102
Kato, T. 2001f, Inf. Bull. Variable Stars, 5104
Kato, T. 2001g, Inf. Bull. Variable Stars, 5107
Kato, T. 2001h, Inf. Bull. Variable Stars, 5110
Kato, T. 2001i, Inf. Bull. Variable Stars, 5122
Kato, T. 2001j, Inf. Bull. Variable Stars, 5137
Kato, T. 2002a, A&A, 384, 206
Kato, T. 2002b, PASJ, 54, L11
Kato, T. 2002c, PASJ, 54, 87
Kato, T. 2002d, Inf. Bull. Variable Stars, 5316
Kato, T. 2002e, Inf. Bull. Variable Stars, 5243
Kato, T. 2002f, Inf. Bull. Variable Stars, 5346
Kato, T. 2002g, IAU Circ., 7791
Kato, T. 2003a, A&A, 399, 695
Kato, T. 2003b, PASJ, submitted
Kato, T. 2003c, PASJ, in press (astro-ph/0308086)
Kato, T. 2003d, VSOLJ Variable Star Bull., in press
Kato, T. 2003e, VSOLJ Variable Star Bull., in press
Kato, T. 2003f, VSOLJ Variable Star Bull., in press
Kato, T., Baba, H., & Nogami, D. 2002a, PASJ, 54, 79
Kato, T., Bolt, G., Nelson, P., Monard, B., Stubbings, R., Pearce, A., Yamaoka, H., & Richards, T. 2003b, MNRAS, 341, 901
Kato, T., Djamaluddin, T., Hirata, R., & Saito, M 1990, IAU Circ., 5007
Kato, T., et al. 2002b, A&A, 396, 929
Kato, T., & Dyck, G. 1988, IAU Circ., 4575
Kato, T., & Fujii, M. 2001, Inf. Bull. Variable Stars, 5150
Kato, T., & Fujii, M. 2003, IAU Circ., 8132
Kato, T., Garradd, G., Stubbings, R., Pearce, A., & Nelson, P. 2001a, Inf. Bull. Variable Stars, 5117
Kato, T., Hanson, G., Poyner, G., Muyllaert, E., Reszelski, M., & Dubovsky, P. A. 2000a, Inf. Bull. Variable Stars, 4932
Kato, T., & Hanzl, D. 1993, IAU Circ., 5747
Kato, T., & Haseda, K. 2000, IAU Circ., 7553
Kato, T., & Haseda, K. 2003, Inf. Bull. Variable Stars, 5417
Kato, T., Haseda, K., Takamizawa, K., Kazarovets, E. V., & Samus, N. N. 1998a, Inf. Bull. Variable Stars, 4585
Kato, T., Haseda, K., Takamizawa, K., & Yamaoka, H. 2002c, A&A, 393, L69
Kato, T., Haseda, K., Yamaoka, H., & Takamizawa, K. 2002d, PASJ, 54, L51
Kato, T., & Hirata, R. 1990a, Inf. Bull. Variable Stars, 3489
Kato, T., & Hirata, R. 1990b, Inf. Bull. Variable Stars, 3529
Kato, T., & Hirata, R. 1991, IAU Circ., 5262
Kato, T., Hirata, R., & Mineshige, S. 1992a, PASJ, 44, L215
Kato, T., Ishioka, R., & Uemura, M. 2002e, PASJ, 54, 1029
Kato, T., Ishioka, R., & Uemura, M. 2002f, PASJ, 54, 1023
Kato, T., Ishioka, R., & Uemura, M. 2002g, PASJ, 54, 1033

Kato, T., Ishioka, R., & Uemura, M. 2003c, Inf. Bull. Variable Stars, 5376

Kato, T., et al. 2001b, IAU Circ., 7672

Kato, T., Kanatsu, K., Takamizawa, K., Takao, A., & Stubbings, R. 2000b, IAU Circ., 7552

Kato, T., Kiyota, S., Novák, R., & Matsumoto, K. 1999a, Inf. Bull. Variable Stars, 4794

Kato, T., & Kunjaya, C. 1995, PASJ, 47, 163

Kato, T., Kunjaya, C., Okyudo, M., & Takahashi, A. 1994, PASJ, 46, L199

Kato, T., Lipkin, Y., Retter, A., & Leibowitz, E. 1998b, Inf. Bull. Variable Stars, 4602

Kato, T., Masuda, S., & Nogami, D. 1995a, Inf. Bull. Variable Stars, 4233

Kato, T., & Matsumoto, K. 1999a, Inf. Bull. Variable Stars, 4763

Kato, T., & Matsumoto, K. 1999b, Inf. Bull. Variable Stars, 4765

Kato, T., & Matsumoto, K. 1999c, Inf. Bull. Variable Stars, 4767

Kato, T., & Matsumoto, K. 1999d, Inf. Bull. Variable Stars, 4776

Kato, T., Matsumoto, K., Nogami, D., Morikawa, K., & Kiyota, S. 2001c, PASJ, 53, 893

Kato, T., Matsumoto, K., & Stubbings, R. 1999b, Inf. Bull. Variable Stars, 4760

Kato, T., Matsumoto, K., & Uemura, M. 2002h, Inf. Bull. Variable Stars, 5262

Kato, T., Minehige, S., & Hirata, R. 1992b, IAU Circ., 5676

Kato, T., Mineshige, S., & Hirata, R. 1993a, IAU Circ., 5704

Kato, T., Mineshige, S., & Hirata, R. 1995b, PASJ, 47, 31

Kato, T., & Nakamura, Y. 2001, IAU Circ., 7559

Kato, T., et al. 2003d, MNRAS, in press (astro-ph/0310037)

Kato, T., et al. 2001d, IAU Circ., 7590

Kato, T., & Nogami, D. 1995, Inf. Bull. Variable Stars, 4260

Kato, T., & Nogami, D. 1997a, PASJ, 49, 341

Kato, T., & Nogami, D. 1997b, PASJ, 49, 481

Kato, T., & Nogami, D. 2001a, Inf. Bull. Variable Stars, 5098

Kato, T., & Nogami, D. 2001b, Inf. Bull. Variable Stars, 5105

Kato, T., & Nogami, D. 2001c, Inf. Bull. Variable Stars, 5124

Kato, T., Nogami, D., & Baba, H. 1995c, Inf. Bull. Variable Stars, 4228

Kato, T., Nogami, D., & Baba, H. 1996a, PASJ, 48, L93

Kato, T., Nogami, D., & Baba, H. 1999c, Inf. Bull. Variable Stars, 4766

Kato, T., Nogami, D., & Baba, H. 2001e, PASJ, 53, 901

Kato, T., Nogami, D., & Baba, H. 2001f, Inf. Bull. Variable Stars, 5103

Kato, T., Nogami, D., Baba, H., Hanson, G., & Poyner, G. 2000c, MNRAS, 315, 140

Kato, T., Nogami, D., Baba, H., & Masuda, S. 2001g, Inf. Bull. Variable Stars, 5118

Kato, T., Nogami, D., Baba, H., Masuda, S., Matsumoto, K., & Kunjaya, C. 1999d, in Disk Instabilities in Close Binary Systems, ed. S. Mineshige, & J. C. Wheeler (Tokyo: Universal Academy Press), 45

Kato, T., Nogami, D., Baba, H., & Matsumoto, K. 1998c, in ASP Conf. Ser. 137, Wild Stars in the Old West, ed. S. Howell, E. Kuulkers, & C. Woodward (San Francisco: ASP), 9

Kato, T., Nogami, D., Baba, H., Matsumoto, K., Arimoto, J., Tanabe, K., & Ishikawa, K. 1996b, PASJ, 48, L21

Kato, T., Nogami, D., Baba, H., Matsumoto, K., Arimoto, J., Tanabe, K., & Ishikawa, K. 1996c, in IAU Colloq. 158, Cataclysmic Variables and Related Objects, ed. A. Evans, & J. H. Wood (Dordrecht: Kluwer Academic Publishers), 77

Kato, T., Nogami, D., Lockley, J. J., & Somers, M. 2001h, Inf. Bull. Variable Stars, 5116

Kato, T., Nogami, D., & Masuda, S. 1996d, PASJ, 48, L5

Kato, T., Nogami, D., & Masuda, S. 2002i, PASJ, 54, 575

Kato, T., Nogami, D., & Masuda, S. 2003e, PASJ, 55, L7

Kato, T., Nogami, D., Masuda, S., & Baba, H. 1998d, PASP, 110, 1400

Kato, T., Nogami, D., Masuda, S., & Hirata, R. 1995d, Inf. Bull. Variable Stars, 4193

Kato, T., Nogami, D., Masuda, S., & Hirata, R. 1996e, PASJ, 48, 45

Kato, T., Nogami, D., Matsumoto, K., & Baba, H. 1997, ftp://vsnet.kusastro.kyoto-u.ac.jp/pub/vsnet/preprints/EG_Cnc/

Kato, T., Nogami, D., Moilanen, M., & Yamaoka, H. 2003f, PASJ, in press (astro-ph/0307064)

Kato, T., et al. 2001i, IAU Circ., 7680

Kato, T., et al. 2001j, IAU Circ., 7678

Kato, T., Okada, I., Jones, A. F., Overbeek, D., & Africa, S. 1993b, IAU Circ., 5871

Kato, T., Pojmanski, G., Nishimura, H., Haseda, K., Shida, R. Y., & Kiyota, S. 2003g, IAU Circ., 8126

Kato, T., Poyner, G., & Kinnunen, T. 2002j, MNRAS, 330, 53

Kato, T., et al. 2001k, Inf. Bull. Variable Stars, 5120

Kato, T., et al. 2003h, MNRAS, 339, 861

Kato, T., & Schmeer, P. 1999, Inf. Bull. Variable Stars, 4757

Kato, T., Sekine, Y., & Hirata, R. 2001l, PASJ, 53, 1191

Kato, T., & Starkey, D. R. 2002, Inf. Bull. Variable Stars, 5358

Kato, T., & Stubbings, R. 2003, Inf. Bull. Variable Stars, 5426

Kato, T., Stubbings, R., Monard, B., Butterworth, N. D., Bolt, G., & Richards, T. 2003i, PASJ, in press (astro-ph/0307308)

Kato, T., Stubbings, R., Monard, B., & Pearce, A. 2000d, Inf. Bull. Variable Stars, 4915

Kato, T., Stubbings, R., Monard, B., Pearce, A., & Nelson, P. 2001m, Inf. Bull. Variable Stars, 5091

Kato, T., Stubbings, R., Nelson, P., Pearce, A., Garradd, G., & Kiyota, S. 2001n, Inf. Bull. Variable Stars, 5159

Kato, T., et al. 2002k, A&A, 395, 541

Kato, T., Stubbings, R., Pearce, A., Dubovsky, P. A., Kiyota, S., Itoh, H., & Simonsen, M. 2001o, Inf. Bull. Variable Stars, 5109

Kato, T., Stubbings, R., Pearce, A., Nelson, P., & Monard, B. 2001p, Inf. Bull. Variable Stars, 5119

Kato, T., et al. 2001q, Inf. Bull. Variable Stars, 5099

Kato, T., & Takamizawa, K. 2001, Inf. Bull. Variable Stars, 5100

Kato, T., & Takata, T. 1991, IAU Circ., 5371

Kato, T., & Uemura, M. 1999a, Inf. Bull. Variable Stars, 4786

Kato, T., & Uemura, M. 1999d, Inf. Bull. Variable Stars, 4787

Kato, T., & Uemura, M. 1999b, Inf. Bull. Variable Stars, 4802

Kato, T., & Uemura, M. 1999c, Inf. Bull. Variable Stars, 4803

Kato, T., & Uemura, M. 2000b, Inf. Bull. Variable Stars, 4882

Kato, T., & Uemura, M. 2000b, Inf. Bull. Variable Stars, 4902

Kato, T., & Uemura, M. 2001a, Inf. Bull. Variable Stars, 5070

Kato, T., & Uemura, M. 2001b, Inf. Bull. Variable Stars, 5077

Kato, T., & Uemura, M. 2001c, Inf. Bull. Variable Stars, 5078

Kato, T., & Uemura, M. 2001d, Inf. Bull. Variable Stars, 5081

Kato, T., & Uemura, M. 2001e, Inf. Bull. Variable Stars, 5097

Kato, T., & Uemura, M. 2001f, Inf. Bull. Variable Stars, 5158

Kato, T., Uemura, M., Buczynski, D., & Schmeer, P. 2001r, Inf. Bull. Variable Stars, 5123

Kato, T., Uemura, M., Haseda, K., Yamaoka, H., Takamizawa, K., & Fujii, M. 2002l, PASJ, 54, 1009

Kato, T., Uemura, M., & Ishioka, R. 2002m, Inf. Bull. Variable Stars, 5268

Kato, T., Uemura, M., Ishioka, R., & Kinnunen, T. 2001s, PASJ, 53, 1185

Kato, T., Uemura, M., Ishioka, R., Matsumoto, K., & Tanabe, K. 2002n, Inf. Bull. Variable Stars, 5284

Kato, T., Uemura, M., Ishioka, R., & Pietz, J. 2002o, PASJ, 54, 1017

Kato, T., et al. 2003j, PASJ, 55, 489

Kato, T., Uemura, M., Matsumoto, K., Kinnunen, T., Garradd, G., Masi, G., & Yamaoka, H. 2002p, PASJ, 54, 999

Kato, T., et al. 2000e, IAU Circ., 7343

Kato, T., et al. 2000f, Inf. Bull. Variable Stars, 4873

Kato, T., Uemura, M., Stubbings, R., Watanabe, T., & Monard, B. 1999e, Inf. Bull. Variable Stars, 4777

Kato, T., Uemura, M., & Yamaoka, H. 2001t, Inf. Bull. Variable Stars, 5082

Kato, T., & Yamaoka, H. 2002, Inf. Bull. Variable Stars, 5334

Kato, T., Yamaoka, H., & Ishioka, R. 2002q, Inf. Bull. Variable Stars, 5309

Kato, T., Yamaoka, H., Ishioka, R., & Uemura, M. 2002r, GCN, 1363

Kato, T., Yamaoka, H., & Kiyota, S. 2002s, IAU Circ., 7786

Kato, T., Yamaoka, H., & Kiyota, S. 2004, PASJ, in press (astro-ph/0309056)

Kato, T., Yamaoka, H., Liller, W., & Monard, B. 2002t, A&A, 391, L7

Kato, T., Yamaoka, H., Torii, K., Ishioka, R., & Uemura, M. 2002u, Inf. Bull. Variable Stars, 5342

Kawabata, K. S., et al. 2002, ApJL, 580, L39

Kawakita, H., Kinugasa, K., & Yamaoka, H. 2001a, IAU Circ., 7625

Kawakita, H., Kinugasa, K., & Yamaoka, H. 2001b, IAU Circ., 7622

Kawazoe, E., & Mineshige, S. 1993, PASJ, 45, 715

Kazarovets, E. V., Kireeva, N. N., Samus, N. N., & Durlevich, O. V. 2003, Inf. Bull. Variable Stars, 5422

Kazarovets, E. V., & Pastukhova, E. N. 2002, Inf. Bull. Variable Stars, 5225

Kazarovets, E. V., Samus, N. N., & Durlevich, O. V. 2000, Inf. Bull. Variable Stars, 4870

Kazarovets, E. V., Samus, N. N., & Durlevich, O. V. 2001, Inf. Bull. Variable Stars, 5135

Kazarovets, V., Samus, N. N., & Durlevich, O. V. 1998, Inf. Bull. Variable Stars, 4655

Kemp, J., Patterson, J., Thorstensen, J. R., Fried, R. E., Skillman, D. R., & Billings, G. 2002, PASP, 114, 623

Kenyon, S. J. 1986, The symbiotic stars (Cambridge and New York: Cambridge University Press)

Kerber, F. 1999, New Astron. Rev., 43, 425

Kerber, F. 2001, Ap&SS, 275, 91

Kholopov, P. N., & Samus, N. N. 1988, Inf. Bull. Variable Stars, 3154

Kholopov, P. N., et al. 1985a, General Catalogue of Variable Stars, fourth edition (Moscow: Nauka Publishing House)

Kholopov, P. N., Samus, N. N., Kazarovets, E. V., & Perova, N. B. 1985b, Inf. Bull. Variable Stars, 2681

Kilkenny, D., & Lloyd Evans, T. 1995, Observatory, 109, 85

Kim, S.-W., Wheeler, J. C., & Mineshige, S. 1992, ApJ, 384, 269

Kimeswenger, S., Lederle, C., Schmeja, S., & Armsdorfer, B. 2002, MNRAS, 336, 43P

King, A. R., & Cannizzo, J. K. 1998, ApJ, 499, 348

King, A. R., & Lasota, J. P. 1990, MNRAS, 247, 214

King, A. R., & Ritter, H. 1998, MNRAS, 293, 42P

King, A. R., Schenker, K., & Hameury, J. M. 2002, MNRAS, 335, 513

King, A. R., Schenker, K., Kolb, U., & Davies, M. B. 2001, MNRAS, 321, 327

King, D., et al. 1993, IAU Circ., 5755

Kinugasa, K., Kawakita, H., Ayani, K., Kawabata, T., & Yamaoka, H. 2002a, IAU Circ., 7811

Kinugasa, K., et al. 2002b, ApJL, 577, L97

Kinugasa, K., Kawakita, H., Ayani, K., & Yamaoka, H. 2002c, IAU Circ., 7848

Kinugasa, K., Kawakita, H., & Yamaoka, H. 2002d, IAU Circ., 7815

Kinugasa, K., Kawakita, H., Yamaoka, H., & Naito, H. 2002e, IAU Circ., 7844

Kipper, T., Kipper, M., & Klochkova, V. G. 1995, A&A, 297, L33

Kipper, T., & Klochkova, V. G. 1999a, Ap&SS, 265, 449

Kipper, T., & Klochkova, V. G. 2001, Baltic Astron., 10, 393

Kipper, T., & Klochkova, V. V. 1999b, Inf. Bull. Variable Stars, 4661

Kiss, L. L., & Szatmary, K. 2002, A&A, 390, 585

Kiss, L. L., Thomson, J. R., Ogloza, W., Fűrész, G., & Sziládi, K. 2001, A&A, 366, 858

Kiyota, S. 1999, Inf. Bull. Variable Stars, 4736

Kiyota, S., & Kato, T. 1998, Inf. Bull. Variable Stars, 4644

Knigge, C., Hynes, R. I., Steeghs, D., Long, K. S., Araujo-Betancor, S., & Marsh, T. R. 2002, ApJL, 580, L151

Kobayashi, N., Goto, M., Terada, H., & Tomono, D. 2000, GCN, 821

Kobayashi, S., & Zhang, B. 2003, ApJ, 582, L75

Kolb, U., & Baraffe, I. 1999, MNRAS, 309, 1034

Kolb, U., Rappaport, S., Schenker, K., & Howell, S. 2001, ApJ, 563, 958

Kolotilov, E. A., Mikolajewska, J., Marrese, P. M., Munari, U., Shugarov, S. Y., & Yudin, B. F. 2001, Astron. Letters, 27, 51

Kraft, R. P. 1958, PASP, 70, 598

Kraft, R. P. 1962, ApJ, 135, 408

Krautter, J., Vogt, N., Klare, G., Wolf, B., Wargau, W., Drechsel, H., & Rahe, J. 1981, A&A, 98, 27

Krzeminski, W., & Vogt, N. 1985, A&A, 144, 124

Kube, J., Gänsicke, B. T., Euchner, F., & Hoffmann, B. 2003, A&A, 404, 1159

Kukarkin, B. V. 1960, Astron. Tsirk., 209, 21

Käufl, H. U., Koller, J., & Kerber, F. 2003, A&A, 406, 981

Kunjaya, C., Kato, T., & Hirata, R. 1994, Inf. Bull. Variable Stars, 4023

Kunjaya, C., Kinugasa, K., Ishioka, R., Kato, T., Iwamatsu, H., & Uemura, M. 2001, Inf. Bull. Variable Stars, 5128

Kunjaya, C., Kinugasa, K., Ishioka, R., Kato, T., Iwamatsu, H., & Uemura, M. 2003, in ASP Conf. Ser. 289, The Proceeding of the IAU 8th Asian-Pacific Regional Meeting, Volume II, ed. S. Ikeuchi, J. Hearnshaw, & T. Hanawa (San Francisco: ASP), 419

Kurtz, D. W. 1986, Ap&SS, 125, 311

Kuulkers, E. 2000, New Astron. Rev., 44, 27

Kuulkers, E. 2001, Astron. Nachr., 322, 9

Kuulkers, E., Howell, S. B., & van Paradijs, J. 1996, ApJL, 462, L87

Kuulkers, E., Knigge, C., Steeghs, D., Wheatley, P. J., & Long, K. S. 2002, in The Physics of Cataclysmic Variables and Related Objects, ed. B. T. Gänsicke, K. Beuermann, & K. Reinsch (San Francisco: ASP), 443

Langer, G. E., Kraft, R. P., & Anderson, K. S. 1974, ApJ, 189, 509

Lasker, B. M., Russell, J. L., Jenkner, H., Sturch, C. R., McLean, B. J., & Shara, M. M. 1996, VizieR Online Data Catalog, I/220

Lasker, B. M., Sturch, C. R., McLean, B. J., Russell, J. L., Jenkner, H., & Shara, M. M. 1990, AJ, 99, 2019

Lasota, J. P., Hameury, J. M., & Huré, J. M. 1995, A&A, 302, L29

Lawlor, T. M., & MacDonald, J. 2002, Ap&SS, 279, 123

Lawlor, T. M., & MacDonald, J. 2003, ApJ, 583, 913

Lazzati, D., Rossi, E., Covino, S., Ghisellini, G., & Malesani, D. 2002, A&A, 396, L5

Leach, R., Hessman, F. V., King, A. R., Stehle, R., & Mattei, J. 1999, MNRAS, 305, 225

Lederle, C., & Kimeswenger, S. 2003, A&A, 397, 951

Leibowitz, E. M. 1993, ApJL, 411, L29

Lépine, S., Shara, M. M., Livio, M., & Zurek, D. 1999, ApJL, 522, L121

Li, Z., & Chevalier, R. A. 2003, ApJ, 589, L69

Liller, W. 1989, IAU Circ., 4820

Liller, W. 1996, Inf. Bull. Variable Stars, 4299

Liller, W. 2000, IAU Circ., 7417

Liller, W. 2002, IAU Circ., 7789

Liller, W., Cacella, P., Kanatsu, K., Kato, T., Pearce, A., Shida, R. Y., & Linnolt, M. 2002a, IAU Circ., 7878

Liller, W., Herald, D., McNaught, R. H., Pearce, A., & Seargent, D. A. J. 1987, IAU Circ., 4428

Liller, W., Kato, T., & Haseda, K. 2002b, IAU Circ., 7879

Littlefair, S. P., Dhillon, V. S., & Martin, E. L. 2003, MNRAS, 340, 264

Liu, Q. Z., van Paradijs, J., & van den Heuvel, E. P. J. 2001, A&A, 368, 1021

Livio, M., Shankar, A., & Truran, J. W. 1988, ApJ, 325, 282

Livio, M., & Shara, M. M. 1987, ApJ, 319, 819

Lobel, A., et al. 2003, ApJ, 583, 923

Long, K. S., Froning, C. S., Gänsicke, B., Knigge, C., Sion, E. M., & Szkody, P. 2003, ApJ, 591, 1172

Lubow, S. H. 1991a, ApJ, 381, 259

Lubow, S. H. 1991b, ApJ, 381, 268

Lubow, S. H. 1992, ApJ, 401, 317

Lynch, D. K., et al. 2002, Ap&SS, 279, 57

McAdam, D., Hurst, G. M., Manning, B., Lubbock, S., & Young, A. 1988, IAU Circ., 4620

McCormick, P. 1998, in ASP Conf. Ser. 137, Wild Stars in the Old West, ed. S. Howell, E. Kuulkers, & C. Woodward (San Francisco: ASP), 415

McCormick, P., & Frank, J. 1998, in ASP Conf. Ser. 137, Wild Stars in the Old West, ed. S. Howell, E. Kuulkers, & C. Woodward (San Francisco: ASP), 493

McNaught, R. H. 1985, IAU Circ., 4036

McNaught, R. H., Hurst, G. M., Taylor, M., & Green, D. W. E. 1985, IAU Circ., 4143

Makita, M., & Mineshige, S. 2002, PASJ, 54, 463

Marsh, T. R., & Horne, K. 1987, Ap&SS, 130, 85

Marsh, T. R., & Horne, K. 1988, MNRAS, 235, 269

Masetti, N., & Regos, E. 1997, New Astron., 2, 429

Mason, E., & Howell, S. 2003, A&A, 403, 699

Mason, E., Howell, S. B., Szkody, P., Harrison, T. E., Holtzman, J. A., & Hoard, D. W. 2002, A&A, 396, 633

Mason, E., Skidmore, W., Howell, S. B., & Mennickent, R. E. 2001, ApJ, 563, 351

Matheson, T., Filippenko, A. V., Chornock, R., Leonard, D. C., & Li, W. 2000, AJ, 119, 2303

Matsuda, T., Makita, M., Fujiwara, H., Nagae, T., Haraguchi, K., Hayashi, E., & Boffin, H. M. J. 2000, Ap&SS, 274, 259

Matsuda, T., Sekino, N., Shima, E., Sawada, K., & Spruit, H. 1990, A&A, 235, 211

Matsumoto, K., Ishioka, R., Uemura, M., Kato, T., & Kawabata, T. 2003a, MNRAS, 339, 1006

Matsumoto, K., & Kato, T. 2000, IAU Circ., 7528

Matsumoto, K., Kato, T., & Hachisu, I. 2003b, PASJ, 55, 297

Matsumoto, K., Kato, T., Nogami, D., Kawaguchi, T., Kinnunen, T., & Poyner, G. 1999, PASJ, 51, 253

Matsumoto, K., Mennickent, R. E., & Kato, T. 2000, A&A, 363, 1029

Matsumoto, K., Nogami, D., Kato, T., & Baba, H. 1998, PASJ, 50, 405

Matsumoto, K., & Schmeer, P. 1996, IAU Circ., 6517

Matsumoto, K., et al. 2001, A&A, 378, 487

Mattei, J., Ducoty, R., Stanton, R., & Scovil, C. 1985, IAU Circ., 4026

Mattei, J. A. 1980, JRASC, 74, 53

Mattei, J. A., Szentasko, L., Schmeer, P., York, D., & Cragg, T. 1995, IAU Circ., 6155

Mattox, J. R., Tosti, G., & Balonek, T. 1999, IAU Circ., 7189

Maza, J., & Gonzalez, L. E. 1983, IAU Circ., 3856

Meikle, P., Hernandez, M., & Fassia, A. 1998, IAU Circ., 6905

Mennickent, R. E., Diaz, M., Skidmore, W., & Sterken, C. 2001a, A&A, 376, 448

Mennickent, R. E., & Diaz, M. P. 2002, MNRAS, 336, 767

Mennickent, R. E., & Honeycutt, R. K. 1995, Inf. Bull. Variable Stars, 4232

Mennickent, R. E., Matsumoto, K., & Arenas, J. 1999, A&A, 348, 466

Mennickent, R. E., Nogami, D., Kato, T., & Worraker, W. 1996, A&A, 315, 493

Mennickent, R. E., Patterson, J., O'Donoghue, D., Unda, E., Harvey, D., Vanmuster, T., & Bolt, G. 1998, Ap&SS, 262, 1

Mennickent, R. E., Sterken, C., Arenas, J., Diaz, M., & Matsumoto, K. 2001b, J. Astron. Data, 7, 1

Mennickent, R. E., Tovmassian, G., Zharikov, S. V., Tappert, C., Greiner, J., Gänsicke, B. T., & Fried, R. E. 2002, A&A, 383, 933

Meyer, F., & Meyer-Hofmeister, E. 1983, A&A, 121, 29

Meyer, F., & Meyer-Hofmeister, E. 1999, A&A, 341, L23

Meyer-Hofmeister, E., Meyer, F., & Liu, B. F. 1998, A&A, 339, 507

Migliardi, M., Dal Farra, E., Kato, T., & Sano, Y. 2001, IAU Circ., 7683

Mineshige, S., Kusnose, M., & Matsumoto, R. 1995, ApJL, 445, L43

Mineshige, S., & Wheeler, J. C. 1989, ApJ, 343, 241

Misselt, K. A., & Shafter, A. W. 1995, AJ, 109, 1757

Molnar, L. A., & Kobulnicky, H. A. 1992, ApJ, 392, 678

Monard, B., & Li, W. 2002, IAU Circ., 7940

Monet, D., Canzian, B., Harris, H., Reid, N., Rhodes, A., & Sell, S. 1998a, VizieR Online Data Catalog, I/243

Monet, D. B. A., et al. 1998b, VizieR Online Data Catalog, I/252

Moon, D., Pirger, B. E., & Eikenberry, S. S. 2001, PASP, 113, 646

Morales-Rueda, L., & Marsh, T. R. 2002, MNRAS, 332, 814

Morales-Rueda, L., Still, M. D., & Roche, P. 1996, MNRAS, 283, 58P

Morales-Rueda, L., Still, M. D., & Roche, P. 1999, MNRAS, 306, 753

Motch, C., Ilovaisky, S. A., & Chevalier, C. 1982, A&A, 109, L1

Motch, C., Ricketts, M. J., Page, C. G., Ilovaisky, S. A., & Chevalier, C. 1983, A&A, 119, 171

Mueller, J., & Rykoski, K. 2000, IAU Circ., 7436

Mukai, K., & Naylor, T. 1995, in Cataclysmic Variables, ed. A. Bianchini, M. della Valle, & M. Orio (Dordrecht: Kluwer Academic Publishers), 517

Mukai, K., Wood, J. H., Naylor, T., Schlegel, E. M., & Swank, J. H. 1997, ApJ, 475, 812

Munari, U. 1992, A&A, 257, 163

Munari, U., Barbon, R., Tomasella, L., & Rejkuba, M. 1998, IAU Circ., 6902

Munari, U., et al. 1996, A&A, 315, 166

Munari, U., et al. 2002, A&A, 389, L51

Munari, U., Lepardo, A., Passuello, R., & Sostero, G. 1995a, IAU Circ., 6259

Munari, U., Tomov, T., Yudin, B., Sostero, G., & Passuello, R. 1995b, IAU Circ., 6151

Munari, U., Yudin, B. F., Sostero, G., & Lepardo, A. 1993, IAU Circ., 5884

Munari, U., & Zwitter, T. 2002, A&A, 383, 188

Munari, U., et al. 1999, A&A, 347, L39

Murray, J. R. 1998, MNRAS, 297, 323

Murray, J. R., Warner, B., & Wickramasinghe, D. T. 2000, MNRAS, 315, 707

Nagata, T., et al. 2003, PASJ, submitted

Nakano, S., Abe, H., Masi, G., Uemura, M., Kato, T., & Kadota, K. 2001, IAU Circ., 7554

Nakano, S., & Aoki, M. 1997, IAU Circ., 6795

Nakano, S., Kushida, R., & Kushida, Y. 1995, IAU Circ., 6134

Nakano, S., Nishimura, H., Kushida, R., & Kushida, Y. 2003a, IAU Circ., 8113

Nakano, S., Sano, Y., Kushida, R., & Kushida, Y. 1997, IAU Circ., 6778

Nakano, S., Sato, H., Nishimura, H., Nakamura, T., Yamaoka, H., & Pearce, A. 2003b, IAU Circ., 8190

Nakano, S., Takamizawa, K., Kushida, Y., & Kato, T. 1998, IAU Circ., 6941

Nakar, E., Piran, T., & Granot, J. 2003, New Astron., 8, 495

Narayan, R., & Yi, I. 1994, ApJ, 428, L13

Naylor, T., Charles, P. A., Mukai, K., & Evans, A. 1992, MNRAS, 258, 449

Nelemans, G., Portegies Zwart, S. F., Verbunt, F., & Yungelson, L. R. 2001, A&A, 368, 939

Nogami, D. 1996, in Physics of Accretion Disks, ed. S. Kato, S. Inagaki, S. Mineshige, & J. Fukue (Amsterdam: Gordon and Breach Science Publishers), 111

Nogami, D. 1998, in ASP Conf. Ser. 137, Wild Stars in the Old West, ed. S. Howell, E. Kuulkers, & C. Woodward (San Francisco: ASP), 495

Nogami, D., Baba, H., Kato, T., & Novák, R. 1998a, PASJ, 50, 297

Nogami, D., Baba, H., Matsumoto, K., & Kato, T. 2003a, PASJ, 55, 483

Nogami, D., Buczynski, D., Baba, H., & Kato, T. 2001a, Inf. Bull. Variable Stars, 5157

Nogami, D., Engels, D., Gänsicke, B. T., Pavlenko, E. P., Novák, R., & Reinsch, K. 2000, A&A, 364, 701

Nogami, D., Ishioka, R., Makita, M., Kato, T., Yamaoka, H., Uemura, M., & Mineshige, S. 2003b, Inf. Bull. Variable Stars, 5416

Nogami, D., & Kato, T. 1995a, IAU Circ., 6164

Nogami, D., & Kato, T. 1995b, Inf. Bull. Variable Stars, 4227

Nogami, D., & Kato, T. 1997, PASJ, 49, 109

Nogami, D., Kato, T., & Baba, H. 2002, PASJ, 54, 987

Nogami, D., Kato, T., Baba, H., & Masuda, S. 1998b, PASJ, 50, L1

Nogami, D., Kato, T., Baba, H., Matsumoto, K., Arimoto, J., Tanabe, K., & Ishikawa, K. 1997a, ApJ, 490, 840

Nogami, D., Kato, T., Baba, H., Novák, R., Lockley, J. J., & Somers, M. 2001b, MNRAS, 322, 79

Nogami, D., Kato, T., & Hirata, R. 1996, PASJ, 48, 607

Nogami, D., Kato, T., & Masuda, S. 1998c, PASJ, 50, 411

Nogami, D., Kato, T., Masuda, S., & Hirata, R. 1995a, Inf. Bull. Variable Stars, 4155

Nogami, D., Kato, T., Masuda, S., & Hirata, R. 1995b, Inf. Bull. Variable Stars, 4163

Nogami, D., Kato, T., Masuda, S., Hirata, R., Matsumoto, K., Tanabe, K., & Yokoo, T. 1995c, PASJ, 47, 897

Nogami, D., Kunjaya, C., Kato, T., & Hirata, R. 1994, Inf. Bull. Variable Stars, 4059

Nogami, D., & Masuda, S. 1997, Inf. Bull. Variable Stars, 4532

Nogami, D., Masuda, S., & Kato, T. 1995d, Inf. Bull. Variable Stars, 4218

Nogami, D., Masuda, S., & Kato, T. 1997b, PASP, 109, 1114

Nogami, D., Masuda, S., Kato, T., & Hirata, R. 1999, PASJ, 51, 115

Nogami, D., et al. 2003c, PASJ, submitted

Nogami, D., et al. 2003d, A&A, 404, 1067

Nomoto, K., Suzuki, T., Shigeyama, T., Kumagai, S., Yamaoka, H., & Saio, H. 1993, Nature, 364, 507

Norton, A. J., Watson, M. G., & King, A. R. 1988, MNRAS, 231, 783

Novák, R., Vanmunster, T., Jensen, L. T., & Nogami, D. 2001, Inf. Bull. Variable Stars, 5108

Nugent, P., Phillips, M., Baron, E., Branch, D., & Hauschildt, P. 1995, ApJL, 455, L147

O'Donoghue, D., & Charles, P. A. 1996, MNRAS, 282, 191

O'Donoghue, D., Chen, A., Marang, F., Mittaz, J. P. D., Winkler, H., & Warner, B. 1991, MNRAS, 250, 363

Ohashi, H., Kato, T., Senda, T., Mii, H., Mizuno, K., Yasaka, Y., & Yamaoka, H. 2002, GCN, 1515

Ohsima, O., Akazawa, H., & Ohkura, N. 1996, Inf. Bull. Variable Stars, 4295

Ohtani, H., et al. 1992, Memoirs of the Faculty of Science, Kyoto University, Series A of Physics, Astrophysics, Geophysics and Chemistry, 38, 167

Okyudo, M., Kato, T., Ishida, T., Tokimasa, N., & Yamaoka, H. 1993, PASJ, 45, L63

Olech, A. 1997a, IAU Circ., 6666

Olech, A. 1997b, Acta Astron., 47, 281

Olech, A. 2003, Acta Astron., 53, 85

Olech, A., Kędzierski, P., Złoczewski, K., Mularczyk, K., & Wiśniewski, M. 2003a, A&A, in press (astro-ph/0307497)

Olech, A., Schwarzenberg-Czerny, A., Kędzierski, P., Złoczewski, K., Mularczyk, K., & Wiśniewski, M. 2003b, Acta Astron., 53, 175

Oppenheimer, B. D., Kenyon, S. J., & Mattei, J. A. 1998, AJ, 115, 1175

Orlandini, M., et al. 2000, A&A, 356, 1630

Orosz, J. A., & Bailyn, C. D. 1997, ApJ, 477, 876

Orosz, J. A., et al. 2001, ApJ, 555, 489

Osaki, Y. 1974, PASJ, 26, 429

Osaki, Y. 1985, A&A, 144, 369

Osaki, Y. 1989, PASJ, 41, 1005

Osaki, Y. 1995a, in Cataclysmic Variables, ed. A. Bianchini, M. della Valle, & M. Orio (Dordrecht: Kluwer Academic Publishers), 307

Osaki, Y. 1995b, PASJ, 47, L11

Osaki, Y. 1995c, PASJ, 47, 47

Osaki, Y. 1995d, PASJ, 47, L25

Osaki, Y. 1996a, PASP, 108, 39

Osaki, Y. 1996b, in IAU Colloq. 158, Cataclysmic Variables and Related Objects, ed. A. Evans, & J. H. Wood (Dordrecht: Kluwer Academic Publishers), 127

Osaki, Y. 2003, PASJ, in press (astro-ph/0307518)

Osaki, Y., & Meyer, F. 2002, A&A, 383, 574

Osaki, Y., & Meyer, F. 2003, A&A, 401, 325

Osaki, Y., Meyer, F., & Meyer-Hofmeister, E. 2001, A&A, 370, 488

Osaki, Y., Shimizu, S., & Tsugawa, M. 1997, PASJ, 49, L19

Otero, S. 2002, Be Star Newsletter, 35, 27

Otero, S., Fraser, B., & Lloyd, C. 2001, Inf. Bull. Variable Stars, 5026

Otero, S. A., Fieseler, P. D., & Lloyd, C. 2000, Inf. Bull. Variable Stars, 4999

Paczynski, B. 1967, Acta Astron., 17, 287

Paczynski, B. 1970, Acta Astron., 20, 47

Paczynski, B. 1971, Acta Astron., 21, 417

Paczyński, B. 1971, ARA&A, 9, 183

Paczynski, B. 1981, Acta Astron., 31, 1

Paczynski, B., & Sienkiewicz, R. 1981, ApJL, 248, L27

Paczynski, B., & Sienkiewicz, R. 1983, ApJ, 268, 825

Papenkova, M., & Li, W. 2003, IAU Circ., 8141

Papenkova, M., & Li, W. D. 2000, IAU Circ., 7415

Papousek, J., & Granslo, B. H. 1992, IAU Circ., 5604

Pastorello, A., Turatto, M., Rizzi, L., Cappellaro, E., Benetti, S., & Patet, F. 1999, IAU Circ., 7245

Patterson, J. 1981, ApJS, 45, 517

Patterson, J. 1984, ApJS, 54, 443

Patterson, J. 1991, PASP, 103, 1149

Patterson, J. 1994, PASP, 106, 209

Patterson, J. 1998, PASP, 110, 1132

Patterson, J. 1999, in Disk Instabilities in Close Binary Systems, ed. S. Mineshige, & J. C. Wheeler (Tokyo: Universal Academy Press), 61

Patterson, J. 2001, PASP, 113, 736

Patterson, J., Augusteijn, T., Harvey, D. A., Skillman, D. R., Abbott, T. M. C., & Thorstensen, J. 1996, PASP, 108, 748

Patterson, J., Bond, H. E., Grauer, A. D., Shafter, A. W., & Mattei, J. A. 1993, PASP, 105, 69

Patterson, J., Harvey, D., & DeYoung, J. A. 1995a, IAU Circ., 6157

Patterson, J., Jablonski, F., Koen, C., O'Donoghue, D., & Skillman, D. R. 1995b, PASP, 107, 1183

Patterson, J., Kemp, J., Jensen, L., Vanmunster, T., Skillman, D. R., Martin, B., Fried, R., & Thorstensen, J. R. 2000a, PASP, 112, 1567

Patterson, J., et al. 1997, PASP, 109, 1100

Patterson, J., et al. 1998, PASP, 110, 1290

Patterson, J., & McGraw, J. T. 1978, IAU Circ., 3311

Patterson, J., McGraw, J. T., Coleman, L., & Africano, J. L. 1981, ApJ, 248, 1067

Patterson, J., et al. 2002, PASP, 114, 721

Patterson, J., Schwartz, D. A., Pye, J. P., Blair, W. P., Williams, G. A., & Caillault, J.-P. 1992, ApJ, 392, 233

Patterson, J., et al. 2003, PASP, in press (astro-ph/0309100)

Patterson, J., Walker, S., Kemp, J., O'Donoghue, D. B., Bos, H., & Stubbings, R. 2000b, PASP, 112, 625

Pavlenko, E. P., & Dudka, O. I. 2002, Astrophysics, 45, 1

Pavlenko, Y. V., & Duerbeck, H. W. 2001, A&A, 367, 933

Pavlenko, Y. V., Yakovina, L. A., & Duerbeck, H. W. 2000, A&A, 354, 229

Pease, R. 1993, Nature, 362, 585

Pepin, M. B. 1995, S&T, 90, 92

Peterson, B. A., & Price, P. A. 2003, GCN, 1985

Phillips, M., & Krisciunas, K. 2001, IAU Circ., 7636

Podsiadlowski, P., Hsu, J. J. L., Joss, P. C., & Ross, R. R. 1993, Nature, 364, 509

Podsiadlowski, Ph., Han, Z., & Rappaport, S. 2003, MNRAS, 340, 1214

Pojmanski, G. 2002, Acta Astron., 52, 397

Politano, M., Howell, S. B., & Rappaport, S. 1998, in ASP Conf. Ser. 137, Wild Stars in the Old West, ed. S. Howell, E. Kuulkers, & C. Woodward (San Francisco: ASP), 207

Poyner, G. 1996, J. British Astron. Assoc., 106, 155

Prialnik, D., & Shara, M. M. 1986, ApJ, 311, 172

Pringle, J. E., et al. 1987, MNRAS, 225, 73

Pych, W., & Olech, A. 1995a, IAU Circ., 6162

Pych, W., & Olech, A. 1995b, Acta Astron., 45, 385

Qiu, Y. I., Huang, L., Yao, B., & Li, H. 1999, IAU Circ., 7244

Rao, N. K., & Lambert, D. L. 2003, PASP, in press (astro-ph/0308447)

Rappaport, S., Joss, P. C., & Verbunt, F. 1983, ApJ, 275, 713

Rappaport, S., Joss, P. C., & Webbink, R. F. 1982, ApJ, 254, 616

Ravasio, M., et al. 2002, A&A, 383, 763

Reinsch, K., & Beuermann, K. 1990, A&A, 240, 360

Remillard, R. A., Morgan, E., Smith, D., & Smith, E. 2000, IAU Circ., 7389

Renvoizé, V., Baraffe, I., Kolb, U., & Ritter, H. 2002, A&A, 389, 485

Retter, A., Hellier, C., Augusteijn, T., Naylor, T., Bedding, T. R., Bembrick, C., McCormick, J., & Velthuis, F. 2003, MNRAS, 340, 679

Retter, A., Leibowitz, E. M., & Ofek, E. O. 1997, MNRAS, 286, 745

Retter, A., O'Toole, S. J., & Starrfield, S. 2002, IAU Circ., 7818

Revnivtsev, M., Gilfanov, M., Churazov, E., & Sunyaev, R. 2002a, A&A, 391, 1013

Revnivtsev, M., Sunyaev, R., Gilfanov, M., & Churazov, E. 2002b, A&A, 385, 904

Reynolds, A. P., de Bruijne, J. H. J., Perryman, M. A. C., Peacock, A., & Bridge, C. M. 2003, A&A, 400, 1209

Richter, G. A. 1985, Inf. Bull. Variable Stars, 2714

Richter, G. A. 1992, in ASP Conf. Ser. 29, Vina del Mar Workshop on Cataclysmic Variable Stars, ed. N. Vogt (San Francisco: ASP), 12

Richter, G. A., & Greiner, J. 1995, in Cataclysmic Variables, ed. A. Bianchini, M. Della Valle, & M. Orio (Dordrecht: Kluwer Academic Publishers), 177

Richter, G. A., Kroll, P., Greiner, J., & Bianchini, A. 1996, IAU Circ., 6295

Richter, G. A., Kroll, P., Greiner, J., Wenzel, W., Luthardt, R., & Schwartz, R. 1997, A&A, 325, 994

Riess, A. G., Press, W. H., & Kirshner, R. P. 1996, ApJ, 473, 88

Ringwald, F. A. 1993, PASP, 105, 805

Ringwald, F. A., Thorstensen, J. R., Honeycutt, R. K., & Smith, R. C. 1996, AJ, 111, 2077

Ripero, J., et al. 1993, IAU Circ., 5731

Ritter, H. 1976, MNRAS, 175, 279

Ritter, H. 1980, Messenger, 21, 16

Ritter, H. 1985, A&A, 145, 227

Ritter, H., & Kolb, U. 1998, A&AS, 129, 83

Ritter, H., & Kolb, U. 2003, A&A, 404, 301

Robertson, J. W., & Honeycutt, R. K. 1996, AJ, 112, 2248

Robertson, J. W., Honeycutt, R. K., & Turner, G. W. 1995, PASP, 107, 443

Robinson, E. L. 1973, ApJ, 183, 193

Robinson, E. L., Barker, E. S., Cochran, A. L., Cochran, W. D., & Nather, R. E. 1981, ApJ, 251, 611

Robinson, E. L., Ivans, I. I., & Welsh, W. F. 2002, ApJ, 565, 1169

Robinson, E. L., Nather, R. E., & Kiplinger, A. 1974, PASP, 86, 401

Robinson, E. L., Shafter, A. W., Hill, J. A., Wood, M. A., & Mattei, J. A. 1987, ApJ, 313, 772

Rolfe, D. J., Abbott, T. M. C., & Haswell, C. A. 2002, MNRAS, 334, 699

Rolfe, D. J., Haswell, C. A., & Patterson, J. 2001, MNRAS, 324, 529

Rosen, S. R., Clayton, K. L., Osborne, J. P., & McGale, P. A. 1994, MNRAS, 269, 913

Rosenbush, A. E. 2001, Inf. Bull. Variable Stars, 5025

Rosino, L., & Iijima, T. 1988, A&A, 201, 89

Rupen, M. P., Mioduszewski, A. J., & Dhawan, V. 2003, IAU Circ., 8180

Rutten, R. G. M., Dhillon, V. S., Horne, K., & Kuulkers, E. 1994, A&A, 283, 441

Rutten, R. G. M., Dhillon, V. S., Horne, K., Kuulkers, E., & van Paradijs, J. 1993, Nature, 362, 518

Rutten, R. G. M., Kuulkers, E., Vogt, N., & van Paradijs, J. 1992, A&A, 265, 159

Sadakane, K., et al. 1996, PASJ, 48, 51

Samus, N. N. 2000a, IAU Circ., 7353

Samus, N. N. 2000b, IAU Circ., 7382

Samus', N. N., et al. 2003, Astron. Letters, 29, 468

Samus', N. N., Goranskii, V. P., Durlevich, O. V., Zharova, A. V., Kazarovets, E. V., Pastukhova, E. N., Hazen, M. L., & Tsvetkova, T. M. 2002, Astron. Letters, 28, 174

Samus, N. N., Kazarovets, E. V., & Durlevich, O. V. 2000, Baltic Astron., 9, 583

Sanders, E., Turatto, M., Riello, M., Altavilla, G., Benetti, S., Pastorello, A., & Cappellaro, E. 2002, IAU Circ., 7921

Sari, R., & Piran, T. 1999a, ApJ, 517, L109

Sari, R., & Piran, T. 1999b, ApJ, 520, 641

Sari, R., Piran, T., & Narayan, R. 1998, ApJ, 497, L17

Sato, H., Nakamura, Y., Fukushima, H., Yamaoka, H., & Kadota, K. 2001, IAU Circ., 7666

Sawada, K., Matsuda, T., & Hachisu, I. 1986, MNRAS, 219, 75

Schaefer, B. E. 1990, ApJL, 355, L39

Schaefer, B. E. 2001, IAU Circ., 7750

Schaefer, B.E., et al. 2003, ApJ, 588, 387

Schenker, K., Kolb, U., & Ritter, H. 1998, MNRAS, 297, 633

Schmeer, P., Hurst, G. M., Kilmartin, P. M., & Gilmore, A. C. 1992, IAU Circ., 5502

Schmeer, P., Waagen, E., Poyner, G., Vanmunster, T., Hanson, G., & Burrows, T. 1996, IAU Circ., 6513

Schmeer, P., Waagen, E., Shaw, L., & Mattiazzo, M. 1999, IAU Circ., 7113

Schmeja, S., Armsdorfer, B., & Kimeswenger, S. 2000, Inf. Bull. Variable Stars, 4957

Schmid, H. M., Kaufer, A., Camenzind, M., Rivinius, T., Stahl, O., Szeifert, T., Tubbesing, S., & Wolf, B. 2001, A&A, 377, 206

Schmidtke, P. C., Ciudin, G. A., Indlekofer, U. R., Johnson, D. R., Fried, R. E., & Honeycutt, R. K. 2002, in ASP Conf. Ser. 261, The Physics of Cataclysmic Variables and Related Objects, ed. B. T. Gänsicke, K. Beuermann, & K. Reinsch (San Francisco: ASP), 539

Schneller, H. 1931, Astron. Nachr., 243, 335

Schreiber, M. R., Gänsicke, B. T., & Mattei, J. A. 2002, A&A, 384, L6

Schulte-Ladbeck, R. 1985, Messenger, 39, 3

Schwarz, H. E., van Amerongen, S., Heemskerk, M. H. M., & van Paradijs, J. 1988, A&A, 202, L16

Schwarzenberg-Czerny, A., Ward, M., Pringle, J. E., Verbunt, F., Hanes, D. A., Jones, D. H. P., & Wade, R. A. 1985, MNRAS, 212, 645

Seki, T., Sato, H., Kato, T., Nishimura, H., Muraoka, Y., Garcia, J., Linnolt, M., & Pearce, A. 2002, IAU Circ., 7809

Sekiguchi, K. 1995, Ap&SS, 230, 75

Sekiguchi, K., Caldwell, J. A. R., Stobie, R. S., & Buckley, D. A. H. 1990a, MNRAS, 245, 28

Sekiguchi, K., et al. 1989, MNRAS, 236, 611

Sekiguchi, K., Feast, M. W., Whitelock, P. A., Overbeek, M. D., Wargau, W., & Jones, J. S. 1988, MNRAS, 234, 281

Sekiguchi, K., et al. 1990b, MNRAS, 246, 78

Semeniuk, I., de Young, J. A., Pych, W., Olech, A., Ruszkowski, M., & Schmidt, R. E. 1995, Acta Astron., 45, 365

Semeniuk, I., Olech, A., Kwast, T., & Nalezyty, M. 1997, Acta Astron., 47, 201

Semeniuk, I., Pych, W., Olech, A., & Ruszkowski, M. 1994, Acta Astron., 44, 277

Shafter, A., Nolthenius, R., Patterson, J., Bond, H. E., Grauer, A. D., Szkody, P., & Kemper, E. 1982, IAU Circ., 3748

Shafter, A. W., Clark, L. L., Holland, J., & Williams, S. J. 2000, PASP, 112, 1467

Shafter, A. W., Szkody, P., Liebert, J., Penning, W. R., Bond, H. E., & Grauer, A. D. 1985, ApJ, 290, 707

Shafter, A. W., Szkody, P., & Thorstensen, J. R. 1986, ApJ, 308, 765

Shara, M. M. 1989, PASP, 101, 5

Shara, M. M., Livio, M., Moffat, A. F. J., & Orio, M. 1986, ApJ, 311, 163

Shenavrin, V. I., & Yudin, B. F. 2001, Astron. Rep., 45, 631

Shirasaki, Y. et al. 2002, GCN, 1565

Shore, S. N., Sonneborn, G., Starrfield, S. G., Hamuy, M., Williams, R. E., Cassatella, A., & Drechsel, H. 1991, ApJ, 370, 193

Shore, S. N., Starrfield, S., Gonzalez-Riestra, R., Hauschildt, P. H., & Sonneborn, G. 1994, Nature, 369, 539

Simon, V. 1994, Inf. Bull. Variable Stars, 4026

Simon, V. 2002, A&A, 382, 910

Sion, E. M., Szkody, P., Gaensicke, B., Cheng, F. H., LaDous, C., & Hassall, B. 2001, ApJ, 555, 834

Skiff, B. A. 2001, Inf. Bull. Variable Stars, 5015

Skillman, D. R., Harvey, D., Patterson, J., & Vanmunster, T. 1997, PASP, 109, 114

Skillman, D. R., et al. 2002, PASP, 114, 630

Skopal, A. 2003, A&A, 401, L17
Skopal, A., Chochol, D., Pribulla, T., & Vanko, M. 2000, Inf. Bull. Variable Stars, 5005
Smak, J. 1976, Acta Astron., 26, 277
Smak, J. 1984, Acta Astron., 34, 93
Smak, J. 1993, Acta Astron., 43, 101
Smak, J. 2003, Acta Astron., 53, 167
Smith, D., Remillard, R., Swank, J., Takeshima, T., & Smith, E. 1998, IAU Circ., 6855
Smith, D. A., Levine, A. M., & Morgan, E. H. 1999, IAU Circ., 7253
Sobotka, P., Smelcer, L., Pejcha, O., Kral, L., Kolasa, M., Hornoch, K., & Lomoz, F. 2002, Inf. Bull. Variable Stars, 5336
Soker, N., & Tylenda, R. 2003, ApJL, 582, L105
Sokoloski, J. L., & Bildsten, L. 1999, ApJ, 517, 919
Sokoloski, J. L., Bildsten, L., & Ho, W. C. G. 2001, MNRAS, 326, 553
Solheim, J.-E. 1995, Baltic Astron., 4, 363
Soria, R., & Kong, A. K. H. 2002, ApJL, 572, L33
Sparks, W. M., & Kutter, G. S. 1973, ApJL, 183, L121
Spogli, C., Fiorucci, M., & Raimondo, G. 2003, Inf. Bull. Variable Stars, 5365
Spogli, C., Fiorucci, M., Tosti, G., Ciprini, S., Nucciarelli, G., Macchia, E., & Monacelli, G. 2002, Inf. Bull. Variable Stars, 5276
Spogli, C., Fiorucci, M., Tosti, G., & Raimondo, G. 2001, Inf. Bull. Variable Stars, 5147
Spruit, H. C., & Kanbach, G. 2002, A&A, 391, 225
Spruit, H. C., Matsuda, T., Inoue, M., & Sawada, K. 1987, MNRAS, 229, 517
Spruit, H. C., & Ritter, H. 1983, A&A, 124, 267
Stanishev, V., Kraicheva, Z., Boffin, H. M. J., & Genkov, V. 2001, A&A, 367, 273
Starrfield, S., & Shore, S. N. 1994, S&T, 87, 20
Steeghs, D., Marsh, T., Knigge, C., Maxted, P. F. L., Kuulkers, E., & Skidmore, W. 2001a, ApJL, 562, L145
Steeghs, D., Marsh, T., Kuulkers, E., & Skidmore, W. 2001b, IAU Circ., 7675
Steeghs, D., O'Brien, K., Horne, K., Gomer, R., & Oke, J. B. 2001c, MNRAS, 323, 484
Steeghs, D., Perryman, M. A. C., Reynolds, A., de Bruijne, J. H. J., Marsh, T., Dhillon, V. S., & Peacock, A. 2003, MNRAS, 339, 810
Stellingwerf, R. F. 1978, ApJ, 224, 953
Stone, R. P. S., Kraft, R. P., & Prosser, C. F. 1993, PASP, 105, 755
Stothers, R. B., & Chin, C. 2001, ApJ, 560, 934
Strohmayer, T. E. 2002, ApJ, 581, 577
Stubbings, R., & Pearce, A. 1999, IAU Circ., 7253
Swartz, D. A., Clocchiatti, A., Benjamin, R., Lester, D. F., & Wheeler, J. C. 1993, Nature, 365, 232
Szatmary, K., Kiss, L. L., & Bebesi, Zs. 2003, A&A, 398, 277
Szentasko, L., Takenaka, Y., Pietz, J., Royer, R., Kato, T., & Schmeer, P. 1994, IAU Circ., 6057
Szkody, P. 1976, ApJ, 207, 190
Szkody, P., Desai, V., Burdullis, T., Hoard, D. W., Fried, R., Garnavich, P., & Gänsicke, B. 2000, ApJ, 540, 983
Szkody, P., Gänsicke, B. T., Sion, E. M., & Howell, S. B. 2002a, ApJ, 574, 950
Szkody, P., & Howell, S. B. 1992, ApJS, 78, 537
Szkody, P., & Howell, S. B. 1995, in Cataclysmic Variables, ed. A. Bianchini, M. della Valle, & M. Orio (Dordrecht: Kluwer Academic Publishers), 339
Szkody, P., Howell, S. B., Mateo, M., & Kreidl, T. J. 1989, PASP, 101, 899
Szkody, P., & Ingram, D. 1994, ApJ, 420, 830
Szkody, P., & Mateo, M. 1983, PASP, 95, 596
Szkody, P., & Mateo, M. 1984, ApJ, 280, 729
Szkody, P., & Mattei, J. A. 1984, PASP, 96, 988
Szkody, P., et al. 2002b, AJ, 123, 413
Szkody, P., Osborne, J., & Hassall, B. J. M. 1988, ApJ, 328, 243
Szkody, P., Silber, A., Sion, E., Downes, R. A., Howell, S. B., Costa, E., & Moreno, H. 1996, AJ, 111, 2379
Taam, R. E., Flannery, B. P., & Faulkner, J. 1980, ApJ, 239, 1017
Takamizawa, K. 1997, VSOLJ Variable Star Bull., 25, 4
Takamizawa, K., Kato, T., Yamamoto, M., Takahashi, S., Hirosawa, H., & H., Maeharam 2000, IAU Circ., 7409
Takamizawa, K., Wakuda, M., & Kato, T. 1998, VSOLJ Variable Star Bull., 30-31, 1
Takao, A., Monard, L. A. G., Tabur, V., & Kato, T. 2003, IAU Circ., 8166
Tanaka, Y., & Lewin, W. H. G. 1995, in X-Ray Binaries, ed. W. H. G. Lewin, J. van Paradijs, & van den Heuvel. E. P. J. (Cambridge: Cambridge University Press), 126
Taniguchi, Y., Murayama, T., Sato, Y., Yadoumaru, Y., Ohyama, Y., Kosugi, G., Yoshida, M., & Kurakami, T. 1993, PASJ, 45, L43
Tappert, C., & Bianchini, A. 2003, A&A, 401, 1101
Tappert, C., & Mennickent, R. E. 2001, Inf. Bull. Variable Stars, 5101
Tatarnikov, A. M., Shenavrin, V. I., Whitelock, P. A., Feast, M. W., & Yudin, B. F. 2001, Astron. Letters, 27, 534
Thorstensen, J. R. 1999, Inf. Bull. Variable Stars, 4749
Thorstensen, J. R. 2000, PASP, 112, 1269
Thorstensen, J. R., & Fenton, W. H. 2003, PASP, 115, 37
Thorstensen, J. R., Fenton, W. H., Patterson, J. O., Kemp, J., Halpern, J., & Baraffe, I. 2002a, PASP, 114, 1117
Thorstensen, J. R., Fenton, W. H., Patterson, J. O., Kemp, J., Krajci, T., & Baraffe, I. 2002b, ApJL, 567, L49
Thorstensen, J. R., Patterson, J. O., Kemp, J., & Vennes, S. 2002c, PASP, 114, 1108
Thorstensen, J. R., Taylor, C. J., Becker, C. M., & Remillard, R. A. 1997, PASP, 109, 477
Tomov, N. A., Taranova, O. G., & Tomova, M. T. 2003, A&A, 401, 669
Torii, K. 2003, GCN, 1986
Torii, K., et al. 2003, ApJL, in press (astro-ph/0309563)
Tovmassian, G., et al. 2003, PASP, 115, 725
Tovmassian, G. H., et al. 1998, A&A, 335, 227
Trammell, S. R., Hines, D. C., & Wheeler, J. C. 1993a, ApJL, 414, L21
Trammell, S. R., Hines, D. C., Wheeler, J. C., Jannuzzi, B., Smith, J. A., Okyudo, M., Ishida, T., & Kato, T. 1993b, IAU Circ., 5780
Trimble, V., & Aschwanden, M. J. 2003, PASP, 115, 514
Truss, M. R., Wynn, G. A., Murray, J. R., & King, A. R. 2002, MNRAS, 337, 1329
Tsugawa, M., & Osaki, Y. 1997, PASJ, 49, 75
Turatto, M., Rizzi, L., Salvo, M., Cappellaro, E., Benetti, S., & Patet, F. 1999, IAU Circ., 7244
Tuthill, P. G., Monnier, J. D., & Danchi, W. C. 1999, Nature, 398, 487
Tuthill, P. G., Monnier, J. D., & Danchi, W. C. 2002, in ASP Conf. Ser. 260, Interacting Winds from Massive Stars, ed. A. F. J. Moffat, & N. St-Louis (San Francisco: ASP), 321

Tutukov, A., & Yungelson, L. 1996, MNRAS, 280, 1035

Tutukov, A. V., & Yungelson, L. R. 1979, Acta Astron., 29, 665

Uemura, M. 2003, GCN, 1989

Uemura, M., Ishioka, R., Iwamatsu, H., Kato, T., & Yamaoka, H. 2001a, GCN, 984

Uemura, M., Ishioka, R., Kato, T., Schmeer, P., Yamaoka, H., Starkey, D., Vanmunster, T., & Pietz, J. 2001b, IAU Circ., 7747

Uemura, M., Ishioka, R., Kato, T., & Yamaoka, H. 2002a, GCN, 1521

Uemura, M., Ishioka, R., Kato, T., & Yamaoka, H. 2002b, GCN, 1537

Uemura, M., Ishioka, R., Kato, T., & Yamaoka, H. 2002c, GCN, 1566

Uemura, M., Ishioka, R., Kato, T., & Yamaoka, H. 2003a, GCN, 1899

Uemura, M., Ishioka, R., Kato, T., & Yamaoka, H. 2003b, GCN, 2252

Uemura, M., & Kato, T. 2000, IAU Circ., 7409

Uemura, M., et al. 2003c, PASJ, submitted

Uemura, M., Kato, T., Ishioka, R., Iwamatsu, H., & Yamaoka, H. 2000a, GCN, 866

Uemura, M., et al. 2002d, PASJ, 54, L79

Uemura, M., et al. 2004a, PASJ, in press (astro-ph/0308154)

Uemura, M., Kato, T., Ishioka, R., & Yamaoka, H. 2003d, PASJ, 55, L31

Uemura, M., et al. 2003e, Nature, 423, 843

Uemura, M., et al. 2004b, PASJ, in press

Uemura, M., et al. 2002e, PASJ, 54, 599

Uemura, M., et al. 2002f, PASJ, 54, L15

Uemura, M., et al. 2002g, PASJ, 54, 285

Uemura, M., et al. 2000b, PASJ, 52, L9

Uemura, M., et al. 2000c, PASJ, 52, L15

Uemura, M., Kato, T., Novak, R., Jensen, J. T., Takamizawa, K., Schmeer, P., Yamaoka, H., & Henden, A. 2000d, IAU Circ., 7349

Uemura, M., Kato, T., Pavlenko, E., Baklanov, A., & Pietz, J. 2001c, PASJ, 53, 539

Uemura, M., Kato, T., Pavlenko, E., Shugarov, S., Mitskevich, M., Fried, R. E., & Sano, Y. 2004c, PASJ, submitted

Uemura, M., Kato, T., & Watanabe, M. 2000e, Inf. Bull. Variable Stars, 4831

Uemura, M., Kato, T., Watanabe, T., Stubbings, R., Monard, B., & Kawai, N. 2002h, PASJ, 54, 95

Uemura, M., Kato, T., & Yamaoka, H. 2000f, IAU Circ., 7390

Uemura, M., Kato, T., & Yamaoka, H. 2001d, GCN, 948

Uemura, M., Kato, T., & Yamaoka, H. 2001e, IAU Circ., 7628

Uemura, M., Kato, T., Yamaoka, H., Fujii, M., Quinn, J., Garnavich, P., & Takamizawa, K. 2000g, IAU Circ., 7382

Uemura, M., Yamaoka, H., Ishioka, R., & Kato, T. 2003f, GCN, 1994

van der Woerd, H., van der Klis, M., van Paradijs, J., Beuermann, K., & Motch, C. 1988, ApJ, 330, 911

van Driel, W., et al. 1993, PASJ, 45, L59

van Genderen, A. M. 1994, A&A, 284, 465

van Genderen, A. M., & Gautschy, A. 1995, A&A, 294, 453

van Paradijs, J., et al. 1994, MNRAS, 267, 465

van Teeseling, A., Hessman, F. V., & Romani, R. W. 1999, A&A, 342, L45

Vanderspek, R., et al. 2003, GCN, 1997

Vanmunster, T. 1997, Odessa Astron. Publ., 10, 47

Vanmunster, T. 2001, Inf. Bull. Variable Stars, 5031

Vanmunster, T., & Howell, S. B. 1996, in IAU Colloq. 158, Cataclysmic Variables and Related Objects, ed. A. Evans, & J. H. Wood (Dordrecht: Kluwer Academic Publishers), 63

Vanmunster, T., Skillman, D. R., & Fried, R. 2000a, Inf. Bull. Variable Stars, 4940

Vanmunster, T., Velthuis, F., & McCormick, J. 2000b, Inf. Bull. Variable Stars, 4955

Veen, P. M., van Genderen, A. M., van der Hucht, K. A., Li, A., Sterken, C., & Dominik, C. 1998, A&A, 329, 199

Verbunt, F. 1984, MNRAS, 209, 227

Villata, M., et al. 2002a, A&A, 390, 407

Villata, M., et al. 2002b, Mem. Soc. Astron. Ita., 73, 1191

Vogt, N. 1980, A&A, 88, 66

Vogt, N. 1983, A&A, 118, 95

Vogt, N. 1993, in 2nd Technion-Haifa Conference on Cataclysmic Variables and Related Physics, ed. O. Regev, & G. Shaviv (Jerusalem: Israel Physical Society), 63

Vrielmann, S., Horne, K., & Hessman, F. V. 1999, MNRAS, 306, 766

Vrielmann, S., & Offutt, W. 2003, MNRAS, 338, 165

Vrielmann, S., Stiening, R. F., & Offutt, W. 2002, MNRAS, 334, 608

Waagen, E., Royer, R., & Sventek, P. 1992, IAU Circ., 5609

Wagner, R. M., Halpern, J. P., & Jackson, M. 2002, IAU Circ., 7785

Wagner, R. M., Schmidt, G. D., Smith, P., Hines, D., & Starrfield, S. G. 2000, IAU Circ., 7417

Wagner, R. M., et al. 2003, IAU Circ., 8176

Wagner, R. M., et al. 1998, AJ, 115, 787

Wakuda, M. 1988, VSOLJ Variable Star Bull., 5, 17

Walker, M. F. 1954, PASP, 66, 230

Walker, M. F. 1955, PASP, 67, 262

Wang, L., Howell, D. A., & Wheeler, J. C. 1998, IAU Circ., 6802

Warner, B. 1979, Inf. Bull. Variable Stars, 1707

Warner, B. 1985, in Interacting Binaries, ed. P. P. Eggelton, & J. E. Pringle (Dordrecht: D. Reidel Publishing Company), 367

Warner, B. 1995a, Ap&SS, 225, 249

Warner, B. 1995b, Cataclysmic Variable Stars (Cambridge: Cambridge University Press)

Warner, B. 1995c, Ap&SS, 226, 187

Warner, B. 1998, in ASP Conf. Ser. 137, Wild Stars in the Old West, ed. S. Howell, E. Kuulkers, & C. Woodward (San Francisco: ASP), 2

Warner, B., & Cropper, M. 1983, MNRAS, 203, 909

Warner, B., Livio, M., & Tout, C. A. 1996, MNRAS, 282, 735

Warner, B., & McGraw, J. T. 1981, MNRAS, 196, 59

Warner, B., & Nather, R. E. 1971, MNRAS, 152, 219

Warner, B., & O'Donoghue, D. 1987, MNRAS, 224, 733

Warner, B., & van Citters, G. W. 1974, Observatory, 94, 116

Warner, B., & Woudt, P. A. 2002, MNRAS, 335, 84

Watanabe, T. 1998a, VSOLJ Variable Star Bull., 32, 2

Watanabe, T. 1998b, VSOLJ Variable Star Bull., 32, 1

Watanabe, T. 1999a, VSOLJ Variable Star Bull., 34, 3

Watanabe, T. 1999b, VSOLJ Variable Star Bull., 34, 3

Watanabe, T. 2000, VSOLJ Variable Star Bull., 37, 1

Watanabe, T. 2001, VSOLJ Variable Star Bull., 38, 1

Watanabe, T., Kiyota, S., & Hasegawa, N. 2000, VSOLJ Variable Star Bull., 35-36, 3

Watson, M. G., King, A. R., & Osborne, J. 1985, MNRAS, 212, 917

Webbink, R. F. 1984, ApJ, 277, 355

Wenzel, W. 1983a, Mitteil. Veränderl. Sterne, 9, 141
Wenzel, W. 1983b, Inf. Bull. Variable Stars, 2262
Wenzel, W. 1984, Inf. Bull. Variable Stars, 2481
Wenzel, W., & Fürtig, W. 1971, Inf. Bull. Variable Stars, 541
West, D., Yamaoka, H., & Masi, G. 2002, IAU Circ., 7975
West, J. D., Liller, W., Yamaoka, H., Pereira, A., Bedient, J., & Hornoch, K. 2001, IAU Circ., 7708
Wheatley, P. J., Kuulkers, E., Drake, J. J., Kaastra, J. S., Mauche, C. W., Starrfield, S. G., & Wagner, R. M. 2001, IAU Circ., 7677
Wheeler, J. C., et al. 1993, ApJL, 417, L71
Whitehurst, R., & King, A. 1991, MNRAS, 249, 25
Whyte, C. A., & Eggleton, P. P. 1980, MNRAS, 190, 801
Williams, D. B. 2000, Inf. Bull. Variable Stars, 4904
Williams, P. M. 1995, in Wolf-Rayet Stars: Binaries, Colliding Winds, Evolution, ed. van der Hucht K. A., & P. M. Williams (IAU Symp. No. 163, Kluwer, Dordrecht), 335
Williams, P. M., van der Hucht, K. A., Pullock, A. M. T., Florkowski, D. R., van der Woerd, H., & Wamsteker, W. M. 1990, MNRAS, 243, 662
Wisniewski, J. P., Morrison, N. D., Bjorkman, K. S., Miroshnichenko, A. S., Gault, A. C., Hoffman, J. L., Meade, M. R., & Nett, J. M. 2003, ApJ, 588, 486
Wolf, S., Mantel, K. H., Horne, K., Barwig, H., Schoembs, R., & Baernbantner, O. 1993, A&A, 273, 160
Wood, J. H., Horne, K., Berriman, G., & Wade, R. A. 1989a, ApJ, 341, 974
Wood, J. H., et al. 1989b, MNRAS, 239, 809
Wood, J. H., Naylor, T., Hassall, B. J. M., & Ramseyer, T. F. 1995, MNRAS, 273, 772
Wood, M. A., Casey, M. J., Garnavich, P. M., & Haag, B. 2002, MNRAS, 334, 87
Wood, M. A., Montgomery, M. M., & Simpson, J. C. 2000, ApJL, 535, L39
Woodward, C. E., Cole, J., Hodge, T., & Gehrz, R. D. 1992, IAU Circ., 5619
Woodward, C. E., Lawrence, G. F., Gehrz, R. D., Jones, T. J., Kobulnicky, H. A., Cole, J., Hodge, T., & Thronson, H. A. J. 1993, ApJL, 408, L37
Woudt, P. A., & Warner, B. 2003a, MNRAS, 343, 313
Woudt, P. A., & Warner, B. 2003b, MNRAS, in press (astro-ph/0307473)
Wu, X., Li, Z., Gao, W., & Leung, K. 2001, ApJL, 549, L81
Wyckoff, S., & Wehinger, P. A. 1979, PASP, 91, 173
Yamaoka, H. 2000, IAU Circ., 7419
Yamaoka, H. 2002, IAU Circ., 7943
Yamaoka, H., & Ayani, K. 2003, IAU Circ., 8142
Yamaoka, H., Ayani, K., & Shirakami, K. 2000, IAU Circ., 7411
Yamaoka, H., Kato, T., Filippenko, A. V., & Van Dyk, S. D. 1998, IAU Circ., 6859
Yamaoka, H., Kato, T., Mashitani, Y., Fukuhara, N., Kereszty, Z., & Sano, Y. 2002a, IAU Circ., 8001
Yamaoka, H., Kato, T., Uemura, M., & Ishioka, R. 2003a, GCN, 2147
Yamaoka, H., Kato, T., Uemura, M., Ishioka, R., & Naito, H. 2003b, in ASP Conf. Ser. 289, The Proceeding of the IAU 8th Asian-Pacific Regional Meeting, Volume II, ed. S. Ikeuchi, J. Hearnshaw, & T. Hanawa (San Francisco: ASP), 363
Yamaoka, H., Kiyota, S., & Nelson, P. 2002b, IAU Circ., 8006
Yamaoka, H., Llapasset, J.-M., Kinugasa, K., Kawakita, H., West, D., Watanabe, J., Tsuda, H., & Sano, Y. 2002c, IAU Circ., 7836
Yamaoka, H., Monard, B., Buil, C., Kato, T., Schmeer, P., Shida, R. Y., & Pearce, A. 2002d, IAU Circ., 7972
Yamaoka, H., & Rogers, W. 2002, IAU Circ., 7836
Yamaoka, H., & Sanders, E. 2002, IAU Circ., 7923
Yamaoka, H., & Yamamoto, M. 2000, IAU Circ., 7344
Yamasaki, T., Kato, S., & Mineshige, S. 1995, PASJ, 47, 59
Yi, I., Kim, S.-W., Vishniac, E. T., & Wheeler, J. C. 1992, ApJL, 391, L25
Yoshida, S., Kadota, K., & Kato, T. 1999a, Inf. Bull. Variable Stars, 4792
Yoshida, S., Kadota, K., & Kato, T. 1999b, Inf. Bull. Variable Stars, 4814
Yoshida, S., Kadota, K., & Kato, T. 2000a, Inf. Bull. Variable Stars, 4841
Yoshida, S., Kadota, K., & Kato, T. 2000b, Inf. Bull. Variable Stars, 4843
Yoshida, S., Kadota, K., Kato, T., & Schmeer, P. 2000c, Inf. Bull. Variable Stars, 4845
Yoshida, S., Ohkura, N., Smelcer, L., & Samus, N. 2003, Inf. Bull. Variable Stars, 5396
Yungelson, L. R., Nelemans, G., & van den Heuvel, E. P. J. 2002, A&A, 388, 546
Zhang, E.-H., Robinson, E. L., & Nather, R. E. 1986, ApJ, 305, 740
Zijlstra, A. A., & Bedding, T. R. 2002, MNRAS, 334, 498
Zimmermann, H. U., et al. 1993, IAU Circ., 5750